

Figure 1: Map of the studied sites

Code	Taxa
AST	Asterionella+Asterionellopsis+Asteroplanus
CHA	Chaetoceros
CRY	Cryptophytes
DIT	Ditylum
EUG	Euglenophytes
GUI	Guinardia
GYM	Gymnodinium+Gyrodinium
LEP	Leptocylindrus
NIT	Nitzschia+Hantzschia
PLE	Pleurosigma+Gyrosigma
PRO	Prorocentrum
PRP	Protoperidinium+Archaeoperidinium+Peridinium
PSE	Pseudo-nitzschia
RHI	Rhizosolenia+Neocalyptrella
SCR	Scrippsiella+Enciculifera+Pentapharsodinium+Bysmatrum
SKE	Skeletonema
THL	Thalassionema+Lioloma
THP	Thalassiosira+Porosira

Table 1: Name and composition of the phytoplanktonic groups used in the paper, based on¹

Name of site	Location	Region	Number of points? ¹	Temperature	Salinity
Men Er Roue	47°32'5" N / 3°5'37" W	Brittany	503	3.8-22.2 (14.4 +/- 3.7)	20.1-38 (33)
Loscolo	47°27'27" N / 2°32'18" W	Brittany	463	5.7-22.4 (14.9 +/- 4.0)	14.0-36.8 (33)
Croisic	47°18'0" N / 2°30'51" W	Brittany	500	4.8-28.9 (14.7 +/- 3.9)	14.7-37.6 (33)
L'Eperon	46°16'13" N / 1°14'16" W	Oléron	460	3.0-26.0 (15.3 +/- 4.8)	13.0-36.6 (33)
Cornard	46°3'19" N / 1°7'50" W	Oléron	491	3.1-29.2 (15.6 +/- 4.8)	19.0-38.1 (33)
Auger	45°47'59" N / 1°12'19" W	Oléron	524	3.0-24.5 (15.4 +/- 4.4)	23.9-36.0 (33)
Buoy7	44°32'32" N / 1°15'49" W	Arcachon	311	7.2-23.9 (15.2 +/- 3.8)	31.8-36.1 (33)
Teychan	44°40'25" N / 1°9'31" W	Arcachon	494	5.5-25.2 (15.5 +/- 4.6)	20.6-35.8 (33)
Antoine	43°22'41" N / 4°50'45" E	Mediterranean Sea	539	4.6-30.0 (16.8 +/- 5.1)	26.8-38.9 (33)
Lazaret	43°5'14" N / 5°54'23" E	Mediterranean Sea	512	8.7-29.2 (17.4 +/- 4.2)	21.6-39.6 (33)

Table 2: Attempt of summary for our locations ; should we add the species for each region ?




Figure 2: Comparison of BIC with different interaction matrices, compared to the null model (diagonal interaction matrix, allowing only intragroup interactions), for four different sites separated by dashed lines (Brittany, Oléron, Arcachon and Mediterranean Sea) and 10 different subsites. Different interaction matrices may allow all interactions between taxa (unconstrained), only interactions within pennate diatoms, centric diatoms, dinoflagellates, or other phytoplanktonic taxa (pence), only interactions within diatoms, dinoflagellates or other taxa (diatdin), or only interactions between taxa belonging to these different groups. As model structures (length of the times series taken into account) are different between sites and subsites, groups of bars should not be compared.

¹From 1996, without linear interpolation




Figure 3: Interaction matrices estimated in Arcachon (a) and in the Mediterranean Sea (b). Only interactions between clades (pennate and centric diatoms, dinoflagellates, other planktonic taxa) are allowed. The figure should be read as taxon i having effect e_{ji} on taxon j . The scale for the coefficient values is given at the bottom left of panel a). 95% significance of coefficients was determined by bootstrapping and is marked by asterisks (*). The identity matrix was subtracted to the interaction matrix ($\mathbf{B}-\mathbf{I}$) in order to make effects on growth rates comparable. Composition of planktonic groups is given in Table 1.




Figure 4: Effect of abiotic variables (temperature, TEMP or salinity, SAL) on phytoplankton group in Brittany (a, b), Oléron (c, d), Arcachon (e, f) and in the Mediterranean Sea (g, h). Each color corresponds to a different site. Error bar corresponds to the 95% confidence interval around the estimated coefficient. All variables were normalized before estimation.




Figure 5: Variance of the coefficient in the interaction matrix ($\mathbf{B-I}$), as a function of their mean, for 10 sites in 4 regions, with a model allowing interactions only within clads (see above). The mean-variance relation was either computed with raw values of intergroup interactions (a) or absolute values of the intergroup coefficients (b). Intragroup coefficients were not modified.

References for the meta-analysis




Figure 6: Ratio of intra-to-intergroup interaction strength, when taking into account only the significant values.

Code on the plot	Ref	Dimension	Type of organisms	System
1a	[2]	12	Phytoplankton	Outside a bay
1b	[2]	12	Phytoplankton	Inside a bay
2a	[3]	7	Phytoplankton	Coastal site
2b	[3]	7	Phytoplankton	Offshore site
3a	[4]	14	Plankton	Lake
3b	[4]	14	Plankton, growing season	Lake
4a	[5]	13	Plankton	Lake
4b	[5]	7	Simpler web, plankton	Lake
5a	[6]	10	Ciliates	Lake
5b	[6]	10	Phytoplankton	Lake
6a	[7], conditional least square estimate	9	Zooplankton	Lake
6b	[7], total least square estimate	9	Zooplankton	Lake
7a	[8]	4	Plankton	Lake
7b	[8]	4	Plankton	Lake with high planktivory
7c	[8]	4	Plankton	Lake with low planktivory
8a	[9]	2	Phytoplankton	Lake
8b	[9]	3	Zooplankton	Lake
9a	[10]	4	Functional groups of plankton	Lake
9b	[10]	5	Taxonomic groups of plankton	Lake
10a	[11]	3	Fish	Baltic Sea
11a	[12]	2	Lynx/Hare	Terrestrial
12a	[13]	3	Insects	Terrestrial

Table 3: References used [TO COMPLETE]

References

- [1] T. Hernández Fariñas, C. Bacher, D. Soudant, C. Belin, and L. Barillé. Assessing phytoplankton realized niches using a French national phytoplankton monitoring network. *Estuarine, coastal and shelf science*, 159:15–27, 2015.
- [2] F. Barraquand, C. Picoche, D. Maurer, L. Carassou, and I. Auby. Coastal phytoplankton community dynamics and coexistence driven by intragroup density-dependence, light and hydrodynamics. *Oikos*, 127(12):1834–1852, 2018.
- [3] J.R. Griffiths, S. Hajdu, A.S. Downing, O. Hjerne, U. Larsson, and M. Winder. Phytoplankton community interactions and environmental sensitivity in coastal and offshore habitats. *Oikos*, 125(8):1134–1143, 2015.
- [4] Stephanie E. Hampton and Daniel E. Schindler. Empirical evaluation of observation scale effects in community time series. *Oikos*, 113(3):424–439, 2006.
- [5] Stephanie E. Hampton, Mark D. Scheuerell, and Daniel E. Schindler. Coalescence in the lake washington story: Interaction strengths in a planktonic food web. *Limnology and oceanography*, 51(5):2042–2051, 2006.
- [6] V. Huber and U. Gaedke. The role of predation for seasonal variability patterns among phytoplankton and ciliates. *Oikos*, 114(2):265–276, 2006.
- [7] A.R. Ives, S.R. Carpenter, and B. Dennis. Community interaction webs and zooplankton responses to planktivory manipulations. *Ecology*, 80(4):1405–1421, 1999.
- [8] A. R. Ives, B. Dennis, K. L. Cottingham, and S. R. Carpenter. Estimating community stability and ecological interactions from time-series data. *Ecological monographs*, 73(2):301–330, 2003.
- [9] J.L. Klug, J.M. Fischer, A.R. Ives, and B. Dennis. Compensatory dynamics in planktonic community responses to pH perturbations. *Ecology*, 81(2):387–398, 2000.
- [10] J.L. Klug and K.L. Cottingham. Interactions among environmental drivers: Community responses to changing nutrients and dissolved organic carbon. *Ecology*, 82(12):3390–3403, 2001.
- [11] M. Lindegren, C. Möllmann, A. Nielsen, and N.C. Stenseth. Preventing the collapse of the Baltic cod stock through an ecosystem-based management approach. *Proceedings of the national academy of sciences*, 106(34):14722–14727, 2009.

- [12] J.O. Vik, C.N. Brinch, S. Boutin, and N.C. Stenseth. Interlinking hare and lynx dynamics using a century's worth of annual data. *Population ecology*, 50(3):267–274, 2008.
- [13] K. Yamamura, M. Yokozawa, M. Nishimori, Y. Ueda, and T. Yokosuka. How to analyze long-term insect population dynamics under climate change: 50-year data of three insect pests in paddy fields. *Population ecology*, 48(1):31–48, 2006.