

Stabilizing niche differences are still required to maintain species-rich communities in temporally variable environments

Coralie Picoche¹, Alix Sauve¹, Frédéric Barraquand^{1,2}

July 2, 2018

¹Integrative and Theoretical Ecology Chair, Labex COTE, University of Bordeaux, ²Institute of Mathematics of Bordeaux, CNRS

Abstract

Explaining coexistence in species-rich communities of primary producers (e.g., tropical trees, phytoplankton) remains a challenge for ecologists because of the likely competition for shared basal resources. Following Hutchinson's seminal suggestion, many theoreticians have tried to create diversity through a fluctuating environment, which impairs or slows down competitive exclusion. There are now several fluctuating-environments models that allow coexistence, but they often produce only a dozen of coexisting species at best. Here, we investigate what it would take to create even richer communities in fluctuating environments using an empirically parameterized model. Building on the forced multi-species Lotka-Volterra model of Scranton and Vasseur (2016), we have investigated the effect of two coexistence mechanisms, namely the storage effect and stabilizing niche differences (SNDs). SNDs occur when intraspecific competition is stronger than interspecific (as opposed to neutral models where competition is symmetric), and we tuned the competition ratio based on empirical phytoplankton data. Although SNDs maintained more species than the storage effect (50 % w.r.t 25%), we show that none of the two coexistence mechanisms considered could, by itself, ensure the

coexistence of all species present at the beginning of our simulations. Realistic seasonal environments only aggravated that picture, as they decreased persistence relative to white noise. Our results suggest that combining different mechanisms for biodiversity maintenance into community models might be more fruitful than trying to find which explains best the observed diversity levels. Our results on community composition (formation of clumps, biomass distribution) in the different scenarios give further hints on how to empirically weight the influence of coexistence mechanisms.

1 Introduction

There has been a rich debate in theoretical ecology on how to reconcile niche and neutral perspectives on species coexistence (Gravel et al, 2006; Carmel et al, 2017). Under the neutral perspective, all species have equal birth and death rates and compete equally (for space) whilst under the niche perspective, birth and death rates can vary between species and various mechanisms contribute to increasing intraspecific over interspecific competition (Hubbell, 2001). But as it has been pointed out repeatedly, niche and neutral processes are not mutually exclusive, and may actually act together to produce observed species coexistence (Gravel et al, 2006; Mutshinda et al, 2009; Götzenberger et al, 2012).

For instance, Scheffer and van Nes (2006) put forward the concept of 'clumpy coexistence', whereby niche and neutral community assembly may be simultaneously at work, creating several clumps of similar species along a single trait axis. Classical stabilizing niche differences enable coexistence of multiple clumps through stronger net intraspecific competition (Chesson, 2000), while within-clump coexistence occurs through neutral processes (Hubbell, 2001; Munoz and Huneman, 2016), as species that differ too little in their fitnesses cannot exclude themselves out on reasonable timeframes. Indeed, clumps on the trait axis eventually thin out in absence of immigration, but transient coexistence can last for extended periods of time. This 'emergent neutrality' within groups (Holt, 2006) has been proposed as a unifying concept for the niche and neutral theories. However, the findings of Scheffer and van Nes (2006) have been disputed due to hidden niches in the original model (Barabás et al, 2013). Hidden niches emerge through stronger intraspecific competition mediated by a an additional intraspecific predation-like term (Barabás et al, 2013). This makes coexistence in the Scheffer and van Nes model more similar to that of the classical Lotka-Volterra model (Barabás et al, 2016), so that coexistence within clumps is not exactly

49 neutral.

50 Still, the idea that niche and neutral assembly can mould communities stays potent (Vernon et al,
51 2013). Since then, several studies have suggested that ‘clumpy coexistence’ can occur in theoretical
52 models, most notably models incorporating temporal variation (Scranton and Vasseur, 2016; Sakavara
53 et al, 2018). In these temporal-variation models, equal competitive strengths are combined with other
54 mechanisms like the storage effect that increase the possibility of coexistence by making the interaction
55 strength covary with a fluctuating environment (see also Barabás et al, 2012).

56 Here, we build on the work of Scranton and Vasseur (2016) to investigate the possibility of coex-
57 istence through species response to fluctuating environments (the so-called storage effect or temporal
58 niche partitioning, that are equivalent concepts for forced Lotka-Volterra models, Barabás et al, 2012;
59 Scranton and Vasseur, 2016). Our enthusiasm for the Scranton and Vasseur (2016) model stems from our
60 interest in phytoplankton communities, that were the key inspiration for the thermal preferences curves
61 modelling intrinsic growth rates in Scranton and Vasseur (2016). However, Scranton and Vasseur (2016)
62 described temperature as a white noise, i.e. independent and identically distributed Gaussian random
63 variates over time. This appeared to us as a key assumption to relax. Under most latitudes, tempera-
64 ture is indeed a seasonal signal, and seasonal forcing can strongly affect the dynamics of the community
65 considered (Vesipa and Ridolfi, 2017). Over short timescales, random temporal variation therefore often
66 only add noise to a largely deterministic seasonal trend. Our present work can therefore be seen as an
67 attempt to blend Scranton and Vasseur (2016)’s framework with the periodic environments of Barabás
68 et al (2012), to better represent the mixture of stochastic and deterministic environmental forces affecting
69 phytoplankton community dynamics.

70 Because many phytoplankton species or genera respond in similar ways to temperature despite hav-
71 ing different optima (Moisan et al, 2002), we hypothesized that a large seasonal variation might not
72 necessarily foster species coexistence. In fact, similar responses to seasonal fluctuations should lead to
73 an increased synchrony of species abundances which, in turn, should theoretically mitigate the expected
74 temporal partitioning. How seasonality affects coexistence (as opposed to a purely randomly fluctuating
75 environment) is therefore a key feature of this paper. In particular, we contrast cases where the storage
76 effect is present vs. absent, which conveniently maps to two different parameterizations of the forced

77 Lotka-Volterra model. Moreover, we realized while running Scranton and Vasseur (2016)’s model that
78 the overall diversity obtained after 5000 years was relatively low compared to what we usually observe in
79 phytoplankton communities (several dozens to hundreds of species). We have therefore sought out which
80 mechanisms to add to attain a truly species-rich community for extended periods of time.

81 In an empirical study combining phytoplankton community-level time series and multivariate au-
82 toregressive models Barraquand et al, 2017, we found that despite a large influence of the environment
83 (including temperature, irradiance, and other factors), intraspecific (or intragenus) competition was rather
84 large and most likely the key driver of species coexistence. In other words, stabilizing niche differences
85 had a large role to play in maintaining species diversity in coastal phytoplankton (Barraquand et al, 2017)

86 Here, we therefore try to establish what are the relative contributions of the storage effect vs intraspe-
87 cific density-dependence to coexistence in a phytoplankton-like theoretical community model. This led
88 us to cross different combinations of seasonality in the forcing signal, presence of the storage effect or
89 not, and intra- vs interspecific competition, in order to disentangle the contributions of all these factors
90 to biodiversity maintenance.

91 **2 Methods**

92 *Models description*

93 The model described in Scranton and Vasseur (2016) is based on a variant of the Lotka-Volterra com-
94 petition model. Fluctuations in the environment are introduced in the model by temperature-dependent
95 intrinsic growth rates (see eq. 1-2, all coefficients are defined in Tab. 1) so that species growth rates can
96 be expressed as:

$$\frac{dN_i}{dt} = r_i(\tau)N_i \left(1 - \sum_{j=1}^S \alpha_{ij}N_j\right) - mN_i \quad (1)$$

$$r_i(\tau) = a_r(\tau_0)e^{E_r \frac{(\tau-\tau_0)}{k\tau\tau_0}} f_i(\tau) \quad (2)$$

$$\text{where } f_i(\tau) = \begin{cases} e^{-|\tau-\tau_i^{opt}|^3/b_i}, & \tau \leq \tau_i^{opt} \\ e^{-5|\tau-\tau_i^{opt}|^3/b_i}, & \tau > \tau_i^{opt} \end{cases} \quad (3)$$

$$\text{and } b_i \text{ defined by numerically solving } \int r_i(\tau)d\tau = A \quad (4)$$

Model parameters are detailed in Table 1, and we set their values to match the features of phytoplankton communities as in Scranton and Vasseur's work (2016). The niche of each species is defined by its thermal optimum τ_i^{opt} . Thermal performance curves defined in eq. 3 are parameterized so that all species share the same niche area (eq. 4), which sets a trade-off between maximum growth rates and niche width.

Table 1: Parameter definition and values of the model described in eq. 1-4

Name	Definition	Value (unit)
S	Number of species	60
N_i	Biomass density of the i th species	(kg/area)
τ	Temperature	K
$r_i(\tau)$	Growth rate of species i as a function of temperature	$\frac{\text{kg}}{\text{kg*year}}$
α_{ij}	Strength of competition of species j on species i	0.001 area/kg
b_i	Normalization of the thermal decay rate	
m	Mortality rate	$15 \frac{\text{kg}}{\text{kg*year}}$
τ_0	Reference temperature	293 K
$a_r(\tau_0)$	Growth rate at reference temperature	$386 \frac{\text{kg}}{\text{kg*year}}$
E_r	Activation energy	0.467 eV
k	Boltzmann's constant	$8.6173324 \cdot 10^{-5} \text{ eV.K}^{-1}$
$f_i(\tau)$	Fraction of the maximum rate achieved for the i th species	
μ_τ	Mean temperature	293 K
σ_τ	Standard deviation for temperature	5 K
τ_{\min}	Minimum thermal optimum	288K
τ_{\max}	Maximum thermal optimum	298 K
A	Niche breadth	$10^{3.1} \frac{\text{kg}}{\text{kg*year}}$
τ_i^{opt}	Thermal optimum for growth of the i th species	K
θ	Scaling between white noise to seasonal signal	$[0, \sqrt{(2)}]$
ρ	Ratio of intra-to-intergroup competition strengths	(1;10)

101 We kept the mean and standard deviation of the forcing signal but included a lower-frequency com-
 102 ponent using a sinusoidal function with a period of 365 days (1 time unit being one day, eq.5). We tune
 103 the ratio of low-to-high frequency with the variable θ so as to keep the same energy content - equal total
 104 variance - in the forcing signal.

$$\tau(t) = \mu_\tau + \theta \sigma_\tau \sin\left(\frac{2\pi t}{365}\right) + \varepsilon_t, \text{ where } \varepsilon_t \sim \mathcal{N}(0, \sigma_\tau \sqrt{1 - \frac{\theta^2}{2}}) \quad (5)$$

105 Note that the upper limit for θ , $\sqrt{2}$, corresponds to a completely determinist model which we do not
 106 explore here (but see Zhao, 1991 proving bounded coexistence). We chose to keep the stochasticity in
 107 the signal and to model a plausible temperature signal with $\theta = 1.3$ with temperature ranging from ## to
 108 ## over one year (see Fig. 1b, Moisan et al, 2002) when considering a seasonal forcing of the dynamics.-

109 The formulation of the forced Lotka-Volterra model of Scranton and Vasseur (2016) implies a storage
 110 effect, as the competition strengths covary positively with the growth rate value $r_i(\tau)$ (Ellner et al, 2016).
 111 To test for the effect of an explicit storage effect in the model, we formulated a new version of this model,
 112 where we removed this assumption by using the mean value of a species' growth rate (\bar{r}_i) to weight the
 113 interaction coefficients.

$$\frac{dN_i}{dt} = N_i \left(r_i(\tau) - \sum_{j=1}^S \bar{r}_i \alpha_{ij} N_j \right) - mN_i \quad (6)$$

114 In this way, competition strengths remain unaffected by the environmental conditions, in contrast to
 115 growth rates (eq. 6), while preserving the same average magnitude as in eq. 1.

116 Stabilizing niche differences are ensured by the addition of the coefficient ρ , which is the ratio of
 117 intra-to-intergroup competition strength. We can therefore re-write the interaction coefficients α_{ij} in eq.

118 7

$$\alpha_{ij} = \alpha (1 + (\rho - 1) \delta_{ij}) \quad (7)$$

119 where δ_{ij} is the Kronecker symbol, equal to 1 if $i = j$ and to 0 otherwise. The value of the parameter
 120 $\rho = 10$ was chosen from analyses of phytoplanktonic data (Barraquand et al, 2017).

121 In addition to two types of environmental forcings (white noise with $\theta = 0$, and seasonal forcing

122 with $\theta = 1.3$), we therefore compare the results for four formulations of the model: with and without
 123 an explicit storage effect (eq. 1 and eq. 6, respectively) ; with and without stabilizing niche differences
 124 ($\rho = 10$ or $\rho = 1$, respectively). These are summed up in Table 2.

$\frac{dN_i}{dt} + mN_i$	Storage effect	No storage effect
Stabilizing niche differences	$r_i(\tau)N_i \left(1 - \sum_{j=1}^S \alpha (1 + 9\delta_{ij}) N_j\right)$	$N_i \left(r_i(\tau) - \sum_{j=1}^S \bar{r}_i \alpha (1 + 9\delta_{ij}) N_j\right)$
No stabilizing niche differences	$r_i(\tau)N_i \left(1 - \sum_{j=1}^S \alpha N_j\right)$	$N_i \left(r_i(\tau) - \sum_{j=1}^S \bar{r}_i \alpha N_j\right)$

Table 2: Growth rate of species i in the four formulations of the model we present

125

126 Set-up

127 We replicate the 'Species sorting' experiment of Scranton and Vasseur (2016) so as to investigate on
 128 how synthetic phytoplankton communities structure under the various scenarios we described above.
 129 We focused on the dynamics of a community initialized with 60 species with thermal optima uniformly
 130 spaced along the interval $[15^\circ\text{C}, 25^\circ\text{C}]$, and with the same initial density $(\frac{1}{\alpha S})$. Each simulation was run
 131 for 5000 years in 1-day intervals. When the density of a species dropped below 10^{-6} , it was considered
 132 extinct. For each combination of the parameters of the model (type of environmental signal, storage
 133 effect and stabilizing niche differences), we ran 100 simulations.

134 All simulations were ran with Matlab's ode45 algorithm, an explicit Runge-Kutta (4,5) algorithm
 135 with an absolute error tolerance of 10^{-8} , and relative error tolerance 10^{-3} .

136 3 Results

137 Typical dynamics of the community following Eq. 1 (the model of Scranton and Vasseur, 2016), with
 138 both the environmental signals described in Eq. 1 (original choice of Scranton and Vasseur, 2016) and
 139 5 (our variant), are shown in Fig. 1. Forcing by a sinusoidal signal produces the strongly seasonally
 140 structured dynamics that are typical of phytoplankton. Even though only 5 species can be seen in Fig.
 141 1c), there were 14 species still present at the end of the simulation forced by a white noise, with large

142 disparities in the range of variation of their biomasses. A third of the species kept a biomass above 10
143 kg/area while 6 out of the 14 remained below the unit. All species in the white noise simulations were
144 clustered along a 3.2°C-range of thermal optima (Fig. 5). No obvious temporal patterns (e.g., cycles)
145 could be seen in the communities forced by white noise. On the other hand, seasonal cycles were clear
146 in the seasonally-forced case of Fig. 1 d). Only 4 species coexisted at the end of the simulation with
147 seasonal forcing, gathered in two groups with large thermal optimum differences (5.7°C between the
148 maximum thermal optimum of the first group and the minimum thermal optimum of the second group).
149 When temperatures are high, the group with a higher thermal optima reach maximum biomass, then as
150 temperature decreases through the season, they leave room for the growth of the low-temperature group.




Figure 1: Times series of temperature (top) and extant species (bottom) for the 2 last years of a 5000 year simulation, with storage effect and no stabilizing niche differences. The forcing temperature is either a white noise (left) or a noisy seasonal signal (right). Line colors of species biomasses correspond to their thermal optima (from blue, corresponding to low thermal optima, to red, corresponding to high thermal optima. The whole range of the biomass distribution is shown on Fig. 3).

151 The decrease in persistence due to seasonality observed in Fig. 1 was confirmed in all our simulations
 152 (Fig. 2). In cases where final species richness varied from one simulation to another (namely, the two
 153 middle cases in Fig. 2): with storage effect but without stabilizing niche differences, or without storage
 154 effect but with stabilizing niche differences), seasonality reduced the number of extant species to, on
 155 average, 27% and 48% of its original value, respectively (Fig. 2). A seasonal signal therefore led to a
 156 much smaller average persistence. There was also less variance in persistence between seasonally forced
 157 simulations when compared to white noise simulations.

Both the stabilizing niche differences and the storage effect have a marked effect on increasing persistence. Without any of these coexistence mechanisms, only one species persisted at the end of the simulations. When only the storage effect was present, the number of extant species varied between 8 and 20 with a white noise, or 2 and 6 with a seasonal signal. On the other hand, when only stabilizing niche differences were present, the number of extant species nearly doubled, varying between 20 and 32, or 12 and 15, with a white noise or a seasonal signal respectively. Remarkably, when the storage effect and stabilizing niche differences (SNDs) both affected the community dynamics, all species persisted in the community, while neither the storage nor SNDs were able to produce that result alone, for both white noise and seasonal forcing.




Figure 2: Number of species still present at the end of 5000-year simulations, initialized with 60 species, with a white noise forcing signal (blue) or a noisy seasonal signal (red). For each model type (with and without storage effect, + / - storage, respectively; with and without Stabilizing Niche Differences, + / - SND, respectively), 100 simulations were launched. Community compositions are stable in the cases -storage-SND and +storage+SND (1 or 60 species are still present at the end of all simulations, respectively).

When the richness of the community was stable (either 1 or 60 species at the end of the simulation, Fig. 3), there were still large differences in the structure of the community due to both stochasticity and the type of forcing (Fig. 3). Without storage effect and SNDs, a white noise forcing favoured species with intermediate thermal optima, with two thirds of the simulations ending with a species with a thermal

171 optima between 18.9°C and 21.4°C (corresponding to 25% of the range of thermal optima present at
172 the beginning of the simulation) and reaching a maximum average biomass in this range (Fig. 3 a).
173 This distribution can be related to a selection for the highest long-term growth rates, averaged over time
174 (see normalized growth rates in Fig. 4). On the contrary, seasonality tended to favour species with
175 larger maximum growth rates, with thermal optima above 22°C, their frequency of persistence and mean
176 biomass increasing with the thermal optimum (Fig. 3). 38% of the simulations therefore ended with the
177 species having the highest temperature optimum, 25°C.

178 When both coexistence mechanisms were present, the 60 initial species coexisted with small varia-
179 tions in biomasses over the 50 simulations (mean CV=0.008 across simulations with either a white noise
180 or a seasonal forcing signal, Fig. 3b and d). The forcing signal modified only the distribution of biomasses
181 resulting in contrasted community structures despite equal richness in both simulation types. With a white
182 noise, the distribution was unimodal with a maximum biomass reached for the second best long-term av-
183 erage growth rate (corresponding to a thermal optimum of 20.2°C). On the contrary, a seasonal signal
184 led to a bi-modal distribution (centered on 17.0°C and 24.4°C), each corresponding to one season, with
185 highest biomass for higher thermal optima (Fig. 3 d). The minimum biomass was reached for the best
186 long-term average growth rate at an intermediate temperature (20.4°C), one species apart from the max-
187 imum biomass in the white noise case ~~., thus there are quite broad changes in community composition~~
188 ~~even though the richness is equal for the two forcing signals. The two modes in the seasonal biomass~~
189 ~~distribution correspond to the two modes in the seasonal signal (winter and summer).~~




Figure 3: Mean biomass distribution over the last 200 years of 100 simulations, as a function of thermal optima, for the two stable-composition cases: without storage effect nor stabilizing niche differences (left, only one species is extant at the end of simulation, its mean value is represented by a dot for one simulation. There can be several dots for the same species, corresponding to multiple simulations ending with this species alone) and with storage effect and stabilizing niche differences (right, all species are present at the end of the simulation, their mean biomass is represented by boxplots of the mean value for 50 simulations). The forcing signal is either a white noise (top) or a seasonal signal (bottom). Each species is identified by its thermal optimum and its color code. Mean (filled circles) and maximum (open circles) growth rates for each species are indexed on the right axis [CP: Not sure about showing GR on this figure, though. I have another one without GR if necessary Fred: if you do this, you need to make it a yy-plot, with a right-hand y-axis labelled “Growth rate”].

190 In cases where the richness of the community varied, the overall shape (multimodal vs. unimodal)
 191 of the marginal distribution of extant species with respect to the trait axis were similar for both types
 192 of environmental forcings (Fig. 4). By contrast, the type of coexistence mechanism generated different
 193 shapes. Indeed, the storage effect led to a multi-modal biomass distribution with respect to thermal
 194 optima. We always observed 3 modes with a white noise and 3 modes in 95% of the seasonal simulations,
 195 Fig. 4a. With a white noise, extant species are grouped in rather similar clumps regarding species thermal
 196 optimum (between 18.8°C and 22.2°C whereas clumps tended to be further apart in the seasonal case,
 197 covering a total range of 7.7°C, with species grouping in the higher part of the thermal range, above
 198 22°C. On the other hand, stabilizing niche differences led to a quasi-uniform biomass distribution (Fig. 4
 199 b). Species characterising communities forced by a white noise stayed in the lower range of temperatures
 200 (in 96% of the simulations, the highest thermal optimum was 22.4°C) while they were filtered out in

communities subjected to a seasonal fluctuations of their environment, for which species with thermal optima above 20.5°C persisted.




Figure 4: Mean biomass distribution over the last 200 years of 100 simulations, as a function of thermal optima, with storage effect and no stabilizing niche differences (top) and without storage effect, with stabilizing niche differences (bottom). The forcing signal is either a white noise (blue) or a seasonal signal (red). Shades of the same colour corresponds to the 50th, 90th and 100th percentiles of the distributions while coloured lines correspond to one representative simulation. Scaled average (whose optimum is indicated by the dashed line) and maximum growth rates are shown as filled and open circles, respectively, and indexed on the left y-axis. [F: Add a right-handed y-axis labelled “Growth rates”]

4 Discussion

We have simulated competitive Lotka-Volterra dynamics forced by a fluctuating environment (e.g., temperature fluctuations) under a range of scenarios allowing more or less coexistence. Two coexistence mechanisms, the storage effect and stabilizing niches differences, could be either present or absent, which led to four scenarios. These four scenarios were crossed with two possibilities for the forcing signal, a

208 white noise and a stochastic yet seasonal signal, both with equal temporal variance. Our investigation
209 therefore built on the model of Scranton and Vasseur (2016), which included a white noise forcing and a
210 storage effect, but considered seven additional combinations of mechanisms. This was motivated by our
211 wish to include two observed features of phytoplankton dynamics: seasonal cycles (Winder and Cloern,
212 2010) and stabilizing niche differences (Barraquand et al, 2017).

213 We have found that first, forced Lotka-Volterra dynamics cannot sustain any diversity with our
214 phytoplankton-based set of parameters, unless the structure is geared to include either a storage effect
215 or SNDs. Although this absence of diversity-enhancing effect of “pure” environmental variation has al-
216 ready been stated by other authors (Barabás et al, 2012; Fox, 2013; Scranton and Vasseur, 2016), this is
217 not always intuitive (Fox, 2013), so we feel compelled to stress it once more: temporal variation in growth
218 rate alone cannot help coexistence within competitive communities. A nice point made by Scranton and
219 Vasseur (2016) was that a built-in storage effect in a forced Lotka-Volterra model, parameterised for phy-
220 toplankton communities, could lead to some degree of coexistence. Our investigation reproduced these
221 results, using the white noise forcing considered by Scranton and Vasseur (2016), but an arguably more
222 realistic seasonal forcing by temperature (thinking of phytoplankton) considerably lessened the number
223 of species coexisting after 5000 timesteps, decreasing from 15 on average to 4 on average. Even imagin-
224 ing that groups represented here are genera or classes rather than species, this is a fairly low diversity for
225 a phytoplankton-like community (Reynolds, 2006).

226 We have therefore sought out whether stabilizing niche differences (SNDs) could produce a higher
227 diversity maintenance, using field-based intra- vs inter-group (species or genera) competition coefficient
228 ratio (Barraquand et al, 2017), where the intra-group density-dependence is approximately 10 times
229 stronger. On their own, SNDs produced a higher level of diversity than the storage effect (almost double
230 for white noise).

231 However, the seasonal forcing still considerably reduced diversity when SNDs were considered, espe-
232 cially the “neutral” kind of diversity, i.e., diversity within clumps of similar traits. This diversity reduction
233 occurs because within a season, the signal autocorrelation gives long, contiguous chunks of time to the
234 best competitor to exclude its less adapted heterospecifics, while white noise generates large temperature
235 shifts more frequently, and thereby forbid such competitive exclusion. In a seasonal setting, even though

236 a species has the best long-term averaged growth rate, it can disappear as a result of a strong competition
237 from both low- and high-temperature tolerant species, with and without storage effect.

238 Our results may appear at odds with recent proposals that seasonal forcing in itself would help main-
239 tain diversity (Sakavara et al, 2018). However, we compared the effect of seasonal forcing to that of
240 other forcing signals while controlling total variance. Thus, the contrast between our results and those
241 of Sakavara et al (2018) may be due to the role of forcing variance over time. Overall, seasonality
242 may be slightly better than no forcing at all in maintaining diversity, but on its own, seasonal forcing of
243 parameters does little to improve coexistence and it does not really foster diversity within clumps here.

244 In our previous empirical study of coastal phytoplankton dynamics (Barraquand et al, 2017), we
245 have not found any storage effect (which certainly does not mean that it could not be observed in other
246 systems). Given the results on species richness and composition presented here, we are skeptical that
247 even if it was present, the storage effect alone could help explaining phytoplankton diversity. This is all
248 the more likely that SNDs seem fairly strong when estimated from phytoplankton time series (Barraquand
249 et al, 2017). However, our results suggest that in phytoplankton-like seasonal environments, even though
250 empirically-based SNDs produce more diversity than the storage effect when considered in isolation,
251 the storage effect can help diversity maintenance when combined to other mechanisms. Indeed, the
252 combination storage effect + SNDs is non-additive: the cases where both SNDs and the storage effect
253 were present showed more diversity than generated by any mechanism on its own.

254 The above results suggest the very exciting idea that multiple coexistence mechanisms might com-
255 bine superadditively, thus helping us to better understand the astounding diversity of primary producers.
256 This logic could, in principle, be extended to mechanisms that we have not considered here (e.g., spa-
257 tial structure, specialized natural enemies, that could be as important here for plankton as they are for
258 tropical trees Bagchi et al; Comita et al; Stump and Chesson; Barraquand et al). Previous research has
259 however demonstrated that generalist seed predation could weaken the storage effect (Kuang and Ches-
260 son, 2009, 2010) thus different mechanisms might not always combine superadditively as we found here.
261 That said, other models have shown that pathogens could enhance the storage effect (Mordecai, 2015)
262 or that equalizing mechanisms due to optimally foraging predators are not greatly affected by environ-
263 mental variation (Stump and Chesson, 2017). Better explaining plant or microbial diversity would then

264 not be about selecting the best unique mechanism susceptible to explain the observed diversity, but rather
 265 better combining those mechanisms together. This may obviously be an annoyance for those who like to
 266 sharpen Occam's razor, but it clearly holds opportunities for theoreticians wishing to investigate syner-
 267 gies between coexistence mechanisms in highly diverse communities. Aside from the synergies between
 268 predator diversity-enhancing effects and SNDs or storage effects (on the temporal axis), one obvious
 269 follow-up of this research would be interactions with spatial structure. Spatial structure occurs both en-
 270 dogeneously, through spatially restricted movements and interactions, and exogeneously, through spatial
 271 variation in environmental covariates [Bolker \(2003\)](#) (~~[Bolker-TPB-2003](#)~~). Numerous studies (~~[Bolker and](#)~~
 272 ~~[Pacala Am-Nat-2000](#)~~[Bolker and Pacala \(1999\)](#), Murrell and Law Ecology 2002) have shown that spa-
 273 tially restricted movements and interactions - very small-scale spatial structure - can help coexistence,
 274 which we believe would be especially important for phytoplankton since many species form colonies
 275 (Reynolds, 2006; see discussion in Barraquand et al, 2017). Moreover, although temperature is usually
 276 relatively spatially homogeneous over space, other drivers (e.g., rainfall, pH in terrestrial ecosystems;
 277 salinity in aquatic ones) may exhibit spatial variation which is a main factor for coexistence (Snyder,
 278 2008). The odds that different (resource) niches, natural enemies, spatial limits to competition and tem-
 279 poral niche partitioning all interact to promote the very high-dimensional coexistence observed in the
 280 field seem much higher than for any of those mechanisms alone. Whether the diversity-enhancing effects
 281 of these mechanisms combine sub-additively (as in Kuang and Chesson, 2010) or super-additively like
 282 here is therefore worthy of further research.

283 **References**

- 284 Bagchi R, Gallery RE, Gripenberg S, Gurr SJ, Narayan L, Addis CE, Freckleton RP, Lewis OT
 285 (2014) Pathogens and insect herbivores drive rainforest plant diversity and composition. *Nature*
 286 506(7486):85–88, doi:10.1038/nature12911
- 287 Barabás G, Meszéna G, Ostling A (2012) Community robustness and limiting similarity in periodic
 288 environments. *Theoretical Ecology* 5(2):265–282, doi:10.1007/s12080-011-0127-z

- 289 Barabás G, D’Andrea R, Rael R, Meszéna G, Ostling A (2013) Emergent neutrality or hidden niches?
290 *Oikos* 122(11):1565–1572, doi:10.1111/j.1600-0706.2013.00298.x
- 291 Barabás G, Michalska-Smith MJ, Allesina S (2016) The Effect of Intra- and Interspecific Competition on
292 Coexistence in Multispecies Communities. *The American Naturalist* pp 000–000, doi:10.1086/686901
- 293 Barraquand F, Picoche C, Maurer D, Carassou L, Auby I (2017) Weak interactions between groups and
294 physical drivers of community dynamics in coastal phytoplankton. *bioRxiv* doi:10.1101/171264
- 295 Bolker B, Pacala S (1999) Spatial Moment Equations for Plant Competition: Understanding Spatial
296 Strategies and the Advantages of Short Dispersal. *The American Naturalist* 153(6):575–602
- 297 Bolker BM (2003) Combining endogenous and exogenous spatial variability in analytical population
298 models. *Theoretical Population Biology* 64(3):255–270, doi:10.1016/S0040-5809(03)00090-X
- 299 Carmel Y, Suprunenko YF, Kunin WE, Kent R, Belmaker J, Bar-Massada A, Cornell SJ (2017) Us-
300 ing exclusion rate to unify niche and neutral perspectives on coexistence. *Oikos* 126(10):1451–1458,
301 doi:10.1111/oik.04380
- 302 Chesson P (2000) Mechanisms of maintenance of species diversity. *Annual review of Ecology and Sys-*
303 *tematics* pp 343–366
- 304 Comita LS, Queenborough SA, Murphy SJ, Eck JL, Xu K, Krishnadas M, Beckman N, Zhu Y (2014)
305 Testing predictions of the Janzen-Connell hypothesis: a meta-analysis of experimental evidence for
306 distance- and density-dependent seed and seedling survival. *Journal of Ecology* 102(4):845–856,
307 doi:10.1111/1365-2745.12232
- 308 Ellner SP, Snyder RE, Adler PB (2016) How to quantify the temporal storage effect using simulations
309 instead of math. *Ecology Letters* doi:10.1111/ele.12672
- 310 Fox JW (2013) The intermediate disturbance hypothesis should be abandoned. *Trends in Ecology &*
311 *Evolution* 28(2):86–92, doi:10.1016/j.tree.2012.08.014
- 312 Götzenberger L, de Bello F, Bråthen KA, Davison J, Dubuis A, Guisan A, Lepä J, Lindborg R,
313 Moora M, Pärtel M, Pellissier L, Pottier J, Vittoz P, Zobel K, Zobel M (2012) Ecological assembly

314 rules in plant communities-approaches, patterns and prospects. *Biological Reviews* 87(1):111–127,
315 doi:10.1111/j.1469-185X.2011.00187.x

316 Gravel D, Canham CD, Beaudet M, Messier C (2006) Reconciling niche and neutrality: the contin-
317 uum hypothesis: Reconciling niche and neutrality. *Ecology Letters* 9(4):399–409, doi:10.1111/j.1461-
318 0248.2006.00884.x

319 Holt R (2006) Emergent neutrality. *Trends in Ecology & Evolution* 21(10):531–533,
320 doi:10.1016/j.tree.2006.08.003

321 Hubbell SP (2001) *The Unified Neutral Theory of Biodiversity and Biogeography* (MPB-32). Princeton
322 University Press

323 Kuang JJ, Chesson P (2009) Coexistence of annual plants: generalist seed predation weakens the storage
324 effect. *Ecology* 90(1):170–182

325 Kuang JJ, Chesson P (2010) Interacting coexistence mechanisms in annual plant communities:
326 frequency-dependent predation and the storage effect. *Theoretical population biology* 77(1):56–70

327 Moisan JR, Moisan TA, Abbott MR (2002) Modelling the effect of temperature on the maximum growth
328 rates of phytoplankton populations. *Ecological Modelling* 153(3):197–215, doi:10.1016/S0304-
329 3800(02)00008-X

330 Mordecai EA (2015) Pathogen impacts on plant diversity in variable environments. *Oikos* 124(4):414–
331 420

332 Munoz F, Huneman P (2016) From the neutral theory to a comprehensive and multiscale theory of eco-
333 logical equivalence. *The Quarterly Review of Biology* 91(3):321–342

334 Mutshinda CM, O’Hara RB, Woiwod IP (2009) What drives community dynamics? *Proceedings of the*
335 *Royal Society B: Biological Sciences* 276(1669):2923–2929, doi:10.1098/rspb.2009.0523

336 Reynolds CS (2006) *The ecology of phytoplankton*. Cambridge University Press

337 Sakavara A, Tsirtsis G, Roelke DL, Mancy R, Spatharis S (2018) Lumpy species coexistence arises
338 robustly in fluctuating resource environments. *Proceedings of the National Academy of Sciences*
339 115(4):738–743, doi:10.1073/pnas.1705944115

340 Scheffer M, van Nes EH (2006) Self-organized similarity, the evolutionary emergence of groups of simi-
341 lar species. *Proceedings of the National Academy of Sciences* 103(16):6230–6235

342 Scranton K, Vasseur DA (2016) Coexistence and emergent neutrality generate synchrony among com-
343 petitors in fluctuating environments. *Theoretical Ecology* doi:10.1007/s12080-016-0294-z

344 Snyder RE (2008) When does environmental variation most influence species coexistence? *Theoretical*
345 *Ecology* 1(3):129–139

346 Stump SM, Chesson P (2015) Distance-responsive predation is not necessary for the Janzen-Connell
347 hypothesis. *Theoretical population biology* 106:60–70

348 Stump SM, Chesson P (2017) How optimally foraging predators promote prey coexistence in a variable
349 environment. *Theoretical population biology* 114:40–58

350 Vergnon R, van Nes EH, Scheffer M (2013) Interpretation and predictions of the Emergent neutrality
351 model: a reply to Barab  s et al. *Oikos* 122(11):1573–1575, doi:10.1111/j.1600-0706.2013.00790.x

352 Vesipa R, Ridolfi L (2017) Impact of seasonal forcing on reactive ecological systems. *Journal of Theo-*
353 *retical Biology* 419:23–35, doi:10.1016/j.jtbi.2017.01.036

354 Winder M, Cloern JE (2010) The annual cycles of phytoplankton biomass. *Philosophical Transactions of*
355 *the Royal Society B: Biological Sciences* 365(1555):3215–3226, doi:10.1098/rstb.2010.0125

356 Zhao XQ (1991) The qualitative analysis of n-species lotka-volterra periodic competition systems. *Math-*
357 *ematical and Computer Modelling* 15(11):3–8




Figure 5: Mean biomasses over the last 200 years of 5000-year simulations as a function of the thermal optimum defining each species, for a white noise (left) or a seasonal signal (right) with storage effect and no stabilizing niche differences. This simulation is the one described in Fig. 1.




Figure 6: Mean biomass distribution over the last 200 years of 9 representative simulations, as a function of thermal optima, without storage effect with stabilizing niche differences, in which temperature is either a seasonal signal (red) or a white noise (blue). The distribution induced by a white noise overlaps the one obtained with a seasonal signal in only 2 simulations out of 50.

359 ~~NOTE: figures below are here for discussion between us. I am not sure we should keep them in the paper.~~

360 ~~I think this illustrates an issue (?) in the model: when I removed the storage effect, I must have changed~~

361 ~~the total carrying capacity of the environment. Should we correct it? Note from Fred: there is clear underyielding v~~

362 ~~I also computed the evenness, not as an explanation but as a further exploration of the community.~~




Figure 7: Sum of the average biomasses of all species for the last 200 years of a 5000-year simulation, initialized with 60 species with total initial biomass 10^3 , with a white noise forcing signal (blue) or a noisy seasonal signal (red). For each model type (with and without storage effect, + / - storage, respectively; with and without Stabilizing Niche Differences, + / - SND, respectively), 100 simulations were launched.




Figure 8: Normalized species evenness of the community, initialized with 60 species with total initial biomass 10^3 , with a white noise forcing signal (blue) or a noisy seasonal signal (red). For each model type (with and without storage effect, + / - storage, respectively; with and without Stabilizing Niche Differences, + / - SND, respectively), 100 simulations were launched.

White noise	Storage effect	No storage effect	Season+White noise	Storage ef
Stabilizing niche differences	60	28 [20-32]	Stabilizing niche differences	60
No stabilizing niche differences	15 [8-20]	1	No stabilizing niche differences	4 [2-6]

Table 3: Table corresponding to Fig. 2. Which one to choose?