



[Re] Reproductive pair correlations and the clustering of organisms

Coralie Picoche^{1,2, } and Frederic Barraquand^{1,2, }

¹Institute of Mathematics of Bordeaux, CNRS University of Bordeaux, Talence, France – ²Integrative and Theoretical Ecology, LabEx COTE, University of Bordeaux, Pessac, France

Edited by
(Editor)

Received
01 November 2018

Published
19 February 2021

DOI
—

Introduction

This article is a reproduction of [1].

Methods

Analytical solution of G

Derivation of $G(r,t)$ – Finding back Eq. (2) in the original paper?

$$\frac{\partial G}{\partial t} = 2Dr^{1-d} \frac{\partial}{\partial r} \left(r^{d-1} \frac{\partial G}{\partial r} \right) + 2(\lambda - \mu)G + \gamma r^{1-d} \frac{\partial}{\partial r} \left(r^{d+1} \frac{\partial G}{\partial r} \right) + 2\lambda C\delta(r) \quad (1)$$

We will focus on the case $d = 2$ and $\lambda = \mu$, which means Eq. (1) can be reduced to

$$\frac{\partial G}{\partial t} = \frac{2D}{r} \frac{\partial}{\partial r} \left(r \frac{\partial G}{\partial r} \right) + \frac{\gamma}{r} \frac{\partial}{\partial r} \left(r^3 \frac{\partial G}{\partial r} \right) + 2\lambda C\delta(r) \quad (2)$$

Analytical solution with advection – In the presence of advection ($\gamma \neq 0$), a steady-state solution can be found.

$$\begin{aligned} 0 &= \frac{2D}{r} \frac{\partial}{\partial r} \left(r \frac{\partial G}{\partial r} \right) + \frac{\gamma}{r} \frac{\partial}{\partial r} \left(r^3 \frac{\partial G}{\partial r} \right) + 2\lambda C\delta(r) \\ \Leftrightarrow 0 &= 2\pi r \left(\frac{2D}{r} \frac{\partial}{\partial r} \left(r \frac{\partial G}{\partial r} \right) + \frac{\gamma}{r} \frac{\partial}{\partial r} \left(r^3 \frac{\partial G}{\partial r} \right) + 2\lambda C\delta(r) \right) \\ \Leftrightarrow 0 &= 2\pi \left(2D \frac{\partial}{\partial r} \left(r \frac{\partial G}{\partial r} \right) + \gamma \frac{\partial}{\partial r} \left(r^3 \frac{\partial G}{\partial r} \right) \right) + 2\pi r 2\lambda C\delta(r) \end{aligned} \quad (3)$$

We can then integrate Eq. (2) over a small area centered on a particle, with radius ρ . Let us first note that

Copyright © 2021 C. Picoche and F. Barraquand, released under a Creative Commons Attribution 4.0 International license.
Correspondence should be addressed to Coralie Picoche (coralie.picoche@u-bordeaux.fr)
The authors have declared that no competing interests exist.
Code is available at https://github.com/CoraliePicoche/brownian_bug_fluid/code..

$$\begin{aligned}
\int_{R^2} \delta(\mathbf{r}) d^2\mathbf{r} &= 1 \\
\Leftrightarrow \int_0^{2\pi} \int_0^\rho \delta(\mathbf{r}') r' dr' d\theta &= 1 \\
\Leftrightarrow 2\pi \int_0^\rho \delta(\mathbf{r}') r' dr' &= 1
\end{aligned} \tag{4}$$

Using Eq. (3) and (4), we can integrate between 0 and ρ ,

$$\begin{aligned}
0 &= 2\pi \left(2D\rho \frac{\partial G}{\partial r} + \gamma\rho^3 \frac{\partial G}{\partial r} \right) + 2\lambda C \\
\Leftrightarrow \frac{\partial G}{\partial r} &= -\frac{1}{2\pi} \frac{2\lambda C}{2D\rho + \gamma\rho^3}
\end{aligned} \tag{5}$$

We can integrate between ρ and ∞ , knowing that $G(\infty) = C^2$.

$$C^2 - G(\rho) = -\frac{1}{2\pi} \int_\rho^\infty \frac{2\lambda C}{2Dr + \gamma r^3} dr \tag{6}$$

Using the variable change $u = 2Dr + \gamma r^3$, the integral is equivalent to $\int \frac{u'}{u} du$

$$C^2 - G(\rho) = -\frac{\lambda C}{2\pi} \frac{1}{4D} [\log(\gamma) - \log(\frac{2D}{r^2} + \gamma)] \tag{7}$$

$$\Leftrightarrow G(\rho) = \frac{\lambda C}{8\pi D} \log\left(\frac{2D + \gamma r^2}{\gamma r^2}\right) + C^2 \tag{8}$$

Finally, the pair correlation function $g = G/C^2$ is defined as

$$g = \frac{\lambda}{8\pi DC} \log\left(\frac{2D + \gamma r^2}{\gamma r^2}\right) + 1 \tag{9}$$

Analytical solution without advection – When $U = 0$, $\gamma = 0$ and there is no steady solution. We can get back to Eq. (2).

$$\frac{\partial G}{\partial t} = \frac{2D}{r} \frac{\partial}{\partial r} \left(r \frac{\partial G}{\partial r} \right) + 2\lambda C \delta(\mathbf{r}) \tag{10}$$

Assuming an isotropic environment, this means

$$\frac{\partial G}{\partial t} - 2D\Delta G = 2\lambda C \delta(\mathbf{r}) \tag{11}$$

where $\Delta = \nabla^2$ is the Laplacian operator.

We therefore have

$$\mathcal{L}G(\mathbf{r}, t) = 2\lambda C \delta(\mathbf{r}) \tag{12}$$

where \mathcal{L} is the linear differential operator $\partial_t - 2D\Delta$.

Let's use the Green's function H , defined with $\mathcal{L}H = \delta(\mathbf{r}, t) = \delta(\mathbf{r})\delta(t)$.

By definition, we know that $G(y) = \int H(y, s) 2\lambda C \delta(s) ds$ (where $y = (\mathbf{r}, t)$) is a solution to Eq.(12).

$$\begin{aligned}
G(\mathbf{r}, t) &= 2\lambda C \int_{\mathbb{R}^2} \int_0^t H(\mathbf{r} - \mathbf{r}', t') \delta(\mathbf{r}') d\mathbf{r}' dt' \\
\Leftrightarrow &= 2\lambda C \int_0^t H(\mathbf{r}, t') dt'
\end{aligned} \tag{13}$$

Let's substitute Eq.(13) in Eq. (10):

$$\begin{aligned}
\frac{\partial}{\partial t} \left(2\lambda C \int_0^t H(\mathbf{r}, t') dt' \right) &= 2D2\lambda C \Delta \int_0^t H(\mathbf{r}, t') dt' + 2\lambda C \delta(\mathbf{r}) \\
\Leftrightarrow \int_0^t \left(\frac{\partial H(\mathbf{r}, t')}{\partial t'} - 2D\Delta H(\mathbf{r}, t') \right) dt' &= \delta(\mathbf{r}) \\
\Leftrightarrow \int_0^t \delta(\mathbf{r}) \delta(t') dt' &= \delta(\mathbf{r})
\end{aligned}$$

which is true.

A solution for the Green's function using $\mathcal{L} = \partial_t - 2D\Delta$ in 2 dimensions is $H(r, t) = \frac{1}{4\pi 2Dt} \exp(\frac{-r^2}{4 \times 2Dt})$. $G(r, t)$ can then be computed:

$$G(r, t) = 2\lambda C \left[\frac{E1\left(\frac{r^2}{8Dt}\right)}{8D\pi} \right]_0^t \tag{14}$$

where $E1$ is the exponential integral. Using $G(r, 0) = C^2$ and $\lim_{x \rightarrow +\infty} E1 = 0$ in Eq. (14),

$$G(r, t) = 2\lambda C \frac{E1\left(\frac{r^2}{8Dt}\right)}{8D\pi} + C^2 \tag{15}$$

$$\Leftrightarrow g(r, t) = \frac{2\lambda}{C} \frac{E1\left(\frac{r^2}{8Dt}\right)}{8D\pi} + 1 \tag{16}$$

Results

Discussion

References

1. W. R. Young, A. J. Roberts, and G. Stuhne. "Reproductive pair correlations and the clustering of organisms." In: **Nature** 412.6844 (2001), pp. 328–331.