INCH-POUND

MIL-N-246B
INTERIM AMENDMENT 3 (OS)
21 June 1990

MILITARY SPECIFICATION

NITROGLYCERIN

This interim amendment is approved for use within the Naval Sea Systems Command(OS), with MIL-N-246B, dated 19 February 1962.

This Navy interim amendment 3(OS), replaces unpublished Navy interim amendment 2(OS), dated 30 April 1990. Army interim amendment 2(AR), dated 8 March 1990 already existed.

PAGE 6

Add new paragraph 4.3.3.4:

"4.3.3.4 Alternate method-determination of nitrogen. The titration method of assay in MIL-STD-286, method T209.11 may be used except the following points of technique shall be incorporated:

a. The sample shall be 0.40 ± 0.05 gm nitroglycerin

b. Place a 2-ounce-per-square-inch nitrogen purge on the titrant reservoir.

c. Set the controller functions for a potentiometric deadstop titration; the pH meter range is 1000 millivolts (mV), the endpoint for nitroglycerin is 200 mV, and for KNO_3, 350 mV.

d. Place the reaction beaker in an external bath with coolant maintained between 2° to 5°C by a refrigeration-circulation unit.

e. Add the sample dropwise into the vortex of the acid. Lower the electrodes and dispensing tip into the acid and immediately begin titrating. Adjust the titrant addition rate to obtain an endpoint within 6 to 8 minutes.

DISTRIBUTION STATEMENT A. Approved for public release; distribution is unlimited.

Source: https://assist.dla.mil -- Downloaded: 2026-10-16T06:49Z
Check the source to verify that this is the current version before use.

f. Percent nitrogen = $\frac{V \times F}{W}$ where,

V = volume of titrant, ml
F = factor, %N/ml
W = moisture free weight of nitroglycerin, gm.

g. The standard shall be 0.50 ± 0.05 gm KNO_3, *NIST, ground to the consistency of table salt. It shall be dried at 135°C for 4 hours or redried at 105°C.

h. Add the KNO_3 into the vortex of acid which has been warmed by the use of a hot beaker. Allow the KNO_3 to dissolve for 5 to 10 minutes before titrating. Adjust the titrant addition rate to obtain an endpoint within 8 to 10 minutes.

i. The factor for the formula weight of $FeSO_4$ is 13.855.

*NIST= National Institute of Standards and Technology"

Preparing Activity:
Navy-OS
(Project 1376-N391)

Source: https://assist.dla.mil -- Downloaded: 2026-10-16T06:49Z
Check the source to verify that this is the current version before use.