MILITARY SPECIFICATION

PROMETHAZINE HYDROCHLORIDE, CHLOROFORM, IPECAC FLUID-EXTRACT, AND POTASSIUM GUAIACOLSULFONATE SYRUP

This specification is approved for use by all Departments and Agencies of the Department of Defense.

1. SCOPE

1.1 Scope. This specification covers Promethazine Hydrochloride, Chloroform, Ipecac Fluidextract and Potassium Guaiacolsulfonate Syrup.

1.2 Classification. The syrup covered by this specification shall be the following sizes (see 6.2):

Size 1 - 1 gal (3.78 liters)
Size 2 - 4 fl oz (118 ml)

2. APPLICABLE DOCUMENTS

2.1 Issues of documents. The following documents, of the issue in effect on date of invitation for bids or request for proposals, form a part of this specification to the extent specified herein:

SPECIFICATIONS

Federal

PPP-C-186 - Containers, Packaging and Packing for Drugs, Chemicals, and Pharmaceuticals (See 6.3)

STANDARDS

Military

MIL-STD-105 - Sampling Procedures and Tables for Inspection by Attributes.

(Copies of specifications and standards required by contractors in connection with specific procurement functions should be obtained from the procuring activity or as directed by the contracting officer.)

FSC 6505

Source: https://assist.dla.mil -- Downloaded: 202640-21T10:25Z
Check the source to verify that this is the currenversion before use.

2.2 Other publications. The following documents form a part of this specification to the extent specified herein. Unless otherwise indicated, the issue in effect on date of invitation for bids or request for proposals shall apply.

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE,
FOOD AND DRUG ADMINISTRATION

Federal Food, Drug, and Cosmetic Act and
Regulations Promulgated Thereunder.

(Application for copies should be addressed to the Food and Drug Administration, Washington, DC 20204.)

U. S. PHARMACOPEIAL CONVENTION, INC.

The United States Pharmacopeia.

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

National Formulary.

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

3. REQUIREMENTS

3.1 Material. Shall be a clear, colored, palatable syrup containing in each 5 ml:

Promethazine Hydrochloride	5 mg
Fluidextract Ipecac	0.17 minims
Potassium Guaiacolsulfonate	44 mg
Chloroform	0.25 minims
Citric Acid, Anhydrous	60 mg
Sodium Citrate	197 mg

3.1.1 Alcohol. Shall contain 7 percent alcohol by volume.

3.2 Additives. The coloring, flavoring, sweetening agents and preservatives used in the syrup shall be in the amounts approved by the Federal Food and Drug Administration.

3.3 Finished product.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:25Z
Check the source to verify that this is the current version before use.

3.3.1 Assay.

3.3.1.1 Promethazine hydrochloride. The syrup shall assay to contain not less than 90.0 percent and not more than 110.0 percent of the required amounts of promethazine hydrochloride when determined as specified in 4.5.1.

3.3.1.2 Alcohol content. The finished preparation shall assay to contain not less than 6.3 percent and not more than 7.7 percent of alcohol by volume when determined as specified in 4.5.3.

3.3.1.3 Chloroform content. The finished preparation shall assay to contain not less than 90.0 percent and not more than 125.0 percent of the required amount of chloroform when determined as specified in 4.5.4.

3.3.1.4 Potassium guaiacolsulfonate. The finished preparation shall contain not less than 90.0 percent and not more than 110.0 percent potassium guaiacolsulfonate when determined as specified in 4.5.5.

3.3.1.5 Citric acid,anhydrous, fluidextract ipecac, and sodium citrate. The citric acid, anhydrous,fluidextract ipecac, and sodium citrate in the syrup shall be determined from the batch formulation record kept on file by the manufacturer. The contents of the four ingredients shall be in accordance with the requirements specified in 3.1.

3.3.2 Specific gravity. The specific gravity of the preparation shall be not less than 1.28 and not more than 1.30 at 25° C., when determined, using a suitable, accurate, and reproducible method.

3.3.3 pH. The pH of the undiluted preparation shall be not less than 4.70 and not more than 5.20 at 25° C., when determined potentiometrically, using the U.S.P. method.

3.3.4 Identification. The promethazine in the preparation shall be identified by its ultraviolet absorption spectrum between 380 nm and 280 nm and by yellow crystal formation when determined as specified in 4.5.2.

3.3.5 Flavor and palatability. The finished syrup shall be palatable and pleasant to the taste with no unpleasant after-taste. Not later than the time specified for opening of bids or receipt of proposals, the offeror shall submit to the contracting officer six (6) individually packaged samples (each containing 4 fl oz) of the finished syrup, representative of the product which the offeror proposes to furnish. Two (2) samples will be subjected to panel testing for a determination of palatability (see 4.6 - Palatability test). The remaining samples will be used by cognizant Government inspection and quality assurance activities for determining compliance of supplies furnished

Source: https://assist.dla.mil -- Downloaded: 202610-21T10:25Z
Check the source to verify that this is the currentversion before use.

hereunder with the palatability requirement. Approval as to palatability of any sample submitted by the offeror will not constitute approval of the sample as to any other requirement of this specification. The requirement for submission of samples for use in determining compliance with the palatability requirement may be waived, provided the offeror states, in his bid or proposal, that the product he proposes to furnish is the same product he has offered to the purchasing activity on a previous procurement and the contracting officer determined that such product was previously procured and/or tested by the purchasing activity and found to comply with the palatability requirement.

3.4 Ingredients.

3.4.1 Ipecac fluidextract. The ipecac fluidextract entering into the manufacture of the finished preparation shall be in accordance with the tests, standards, and requirements of the U.S.P. XVI, which was official from 1 October 1960 to 31 August 1965.

3.4.2 Potassium guaiacolsulfonate. The potassium guaiacolsulfonate entering into the manufacture of the finished preparation shall be in accordance with the tests, standards, and requirements of the N.F. XIII, which was official from 1 September 1970 to 30 June 1975.

3.4.3 Other ingredients. All other ingredients entering into the manufacture of the finished preparation shall be of U.S.P. or N.F. quality, or if not included in either of these compendia, the ingredients shall be suitable for use in the preparation.

3.5 Delivery. Not more than 4 months shall have elapsed from the date of manufacture of the syrup to the date of delivery to the Government.

3.6 Workmanship. The material and its containers shall be free from defects which detract from their appearance or may impair their serviceability.

4. QUALITY ASSURANCE PROVISIONS

4.1 Supplier responsibility for inspection. Unless otherwise specified in the contract or purchase order, the supplier is responsible for the performance of all inspection requirements as specified herein. Except as otherwise specified in the contract or order, the supplier may use his own or any other facilities suitable for the performance of the inspection requirements specified herein, unless disapproved by the Government. The Government reserves the right to perform any of the inspections set forth in the specification where such inspections are deemed necessary to assure supplies and services conform to prescribed requirements.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:25Z
Check the source to verify that this is the current version before use.

4.1.1 Records of examinations and tests performed by or for the contractor shall be maintained by the contractor and made available to the Government, upon the Government's request, at any time, or from time to time, during the performance of the contract and, (i) as to any expiration dated item, for the expiration dating period (potency period) specified by the contractor for such item, or such longer period as may be required by regulation of any federal agency; or (ii) as to non-dated items, for not less than 3 years after delivery of the items to the Government.

4.1.2 No company supplying any ingredient(s) to the contractor will be considered an acceptable facility for the performance of any inspection requirements specified herein.

4.2 Lot. For purposes of this specification, a lot, batch, or control is that single, uniform, and homogeneous quantity of syrup produced from one formulation, subjected to the same compounding and manufacturing operation, and filled into final containers.

4.3 Sampling. Sampling shall be conducted in accordance with MIL-STD-105.

TABLE I. Sampling

For visual examination	Inspection level	AQL (percent defective)	Unit of product
Major	S-2	1.0	Filled immediate container
Minor	II	2.5	Filled immediate container

4.3.1 For end item testing. Noncompliance with any of the following is cause for rejection.

TABLE II. For end item testing

Characteristic	Requirement paragraph	Inspection level 1/	Minimum sample
Assay:			
Promethazine hydrochloride	3.3.1.1	S-2	5 } Composite
Alcohol content	3.3.1.2	S-2	5 } Composite
Chloroform content	3.3.1.3	S-2	5 } Composite
Potassium guaiacolsulfonate	3.3.1.4	S-2	5 } Composite

Source: https://assist.dla.mil -- Downloaded: 202640-21T10:25Z
Check the source to verify that this is the current version before use.

Table II. For end item testing (cont'd)

Characteristic	Requirement paragraph	Inspection level 1/	Minimum sample
Specific gravity	3.3.2	S-2	5
pH	3.3.3	S-2	5 Composite
Identification	3.3.4	S-2	5
Palatability	3.3.5	-	-

1/ Where possible, the sample shall be used for two or more tests.

4,4 Examination. The finished syrup shall be examined to determine compliance with all requirements of this specification. Nonconformance will be permitted to the extent indicated in 4.3.

4.4.1 Classification of defects. Examination will be conducted in accordance with the following classification of defects:

TABLE III. Classification of defects

Categories	Defects 1/
Major	
101	Syrup not free from foreign matter.
102	Syrup has crystal formation.
103	Syrup not homogeneous.
104	Syrup not clear.
105	Syrup not colored.
Minor	
201	Exterior of immediate container is not free from contents and foreign matter.
202	Label not free from defacing marks and tears.

1/ Inspection is not restricted to the classified possible defects listed above.

4.5 Tests.

4.5.1 Assays for promethazine hydrochloride.

a. Palladium Chloride Solution - Transfer 500 mg of palladium chloride, $PdCl_2$ to a 500 ml beaker, add 5 ml of hydrochloric acid, and warm on a steam bath until solution is complete. Cool, add 300 ml of purified water and 41 gram of sodium acetate, mix to dissolve and adjust the solution's pH to 3.5 with concentrated hydrochloric acid. Dilute the solution to 500 ml with purified water and store in a glass stoppered bottle.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:25Z
Check the source to verify that this is the current version before use.

b. Standard Preparation - Dissolve a suitable quantity of U.S.P. Promethazine Hydrochloride Reference Standard, accurately weighed, in purified water to obtain a solution having a known final concentration of about 1 mg per ml (Protect the solution from light).

c. Procedure - Transfer 2.0 ml of the sample, equivalent to about 2 mg of promethazine hydrochloride, and 2.0 ml of Standard Preparation to separate 125 ml separators containing 10 ml of Palladium Chloride Solution. Shake well for 30 seconds and add 10 ml of a saturated sodium chloride solution to each separator. Extract the palladium-promethazine complex from each one with three 15 ml portions of ethylene dichloride collecting the extracts in 50 ml volumetric flasks through ethylene dichloride-wetted cotton overlaid with anhydrous sodium sulfate (granular). Wash the cotton with ethylene dichloride, using the washings to dilute the extracts to 50.0 ml. Concomitantly measure the absorbances of the extracts at 415 nm with a suitable spectrophotometer in 1 cm cells, and using ethylene dichloride as the blank. Retain the sample extract for Identification Test 4.5.2a., and calculate the assay as follows:

$$\frac{Au}{As} \times 100C = \text{percent of required amount}$$

where,

Au = sample absorbance

As = standard absorbance

C = concentration of Standard Preparation, mg/ml

4.5.2 Identification for Promethazine Hydrochloride.

a. Dilute the sample extract retained from test 4.4.1 with an equal volume of ethylene dichloride. Record its ultraviolet absorption spectrum between 380 and 280 nm, with a suitable recording spectrophotometer, in 1 cm cells, and using ethylene dichloride as the blank. The spectrum exhibits a minimum at 350 + 2 nm and a maximum at 305 $\pm$ 2 nm.

Source: https://assist.dla.mil -- Downloaded: 202640-21T10:25Z
Check the source to verify that this is the current version before use.

b. Transfer 50 ml of sample to a 250 ml separatory funnel, add 50 ml of chloroform and 10 ml of anhydrous methanol. Stopper and shake vigorously for 5 minutes. Allow the two phases to separate and transfer the bottom layer to a 200 ml beaker. Evaporate to dryness on a steam bath with the aid of a current of air. Add 5 ml of 3.5 percent anhydrous methanol picric acid solution and heat on a steam bath for 5 minutes. Cool in an ice bath until yellow crystals appear.

4.5.3 Alcohol content. Adjust the temperature of the sample to 25° C and transfer exactly 100 ml to a distilling flask. Dilute with 150 ml purified water and distill, collecting about 90 ml distillate in a 100 ml volumetric flask immersed in an ice bath. Rinse the distillate into a separator, add 10 grams NaCl and 30 ml petroleum ether. Shake the mixture to extract volatile oils and $CHCl_3$. Draw off the water-alcohol layer into a distilling flask. Wash the petroleum ether solution in the separator with two 40 ml portions of saturated NaCl solution, adding the aqueous washings to the distilling flask. Distill, collecting about 90 ml distillate in a 100 ml volumetric flask immersed in an ice bath. Adjust the temperature of the distillate to 25°C and dilute to the mark with purified water at the same temperature. Add about 5 gram talc, mix, and filter. Determine the specific gravity of the filtrate at 25°C in a pycnometer. Calculate percent required amount of alcohol. As an alternate, the alcohol content may be determined by a suitable gas chromatography method yielding accurate and reproducible results.

4.5.4 Chloroform assay.

REAGENTS:

Cyclohexane - B.P. 80° - 81° C
Chloroform - Suitable for Spectrophotometric use.

STANDARD CHLOROFORM SOLUTION:

Transfer exactly 5 ml of chloroform to a 250 ml volumetric flask containing about 150 ml of cyclohexane. Dilute to the mark with cyclohexane and mix well. Transfer exactly 20 ml of this solution to a 100 ml volumetric flask containing 50 ml cyclohexane. Dilute to the mark with cyclohexane and mix well (Chloroform concentration at this dilution is 0.40 ml/100 ml).

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:25Z
Check the source to verify that this is the current version before use.

PROCEDURE: ALL TRANSFERS SHALL BE MADE AS RAPIDLY AS POSSIBLE.

Transfer about 20 ml of sample accurately measured to a suitable glass stoppered centrifuge tube (or other suitable vessel), add 4 glass beads and exactly 20 ml of cyclohexane. Stopper securely and shake vigorously for 5 minutes. Centrifuge at 3000 rpm for 5 minutes. Determine the absorbance of the cyclohexane solution of the sample and standard in 0.1 mm NaCl cells, using a suitable infrared spectrophotometer.

Calculate percent required amount of chloroform.

NOTE: Set the instrument for optimal operating conditions. Determine the absorbances of sample and standard at 763 cm-1.

4.5.5 Assay for potassium guaiacolsulfonate.

a. Reagents and Solutions:

1. 4-Aminoantipyrene Hydrochloride Solution: Prepare a 1 percent solution of the reagent in purified water. Store the solution in amber glassware and use within one week.

2. Potassium Ferricyanide Solution: Prepare a fresh 4 percent solution in purified water. (protect from light).

3. Buffer Solution: Dissolve 121.1 gram of tris (hydroxymethyl) aminomethane in about 900 ml of purified water. Adjust the pH to 9.0 with hydrochloric acid and dilute to 1000 ml with purified water.

b. Standard Preparation:

Accurately weigh a suitable quantity of potassium guaiacolsulfonate reference standard and dilute with purified water to obtain a known final concentration of approximately 0.088 mg per ml.

c. Assay Preparation:

Dilute 5.0 ml of the sample (equivalent to about 44 mg of potassium guaiacolsulfonate hemihydrate) to 500.0 ml with purified water.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:25Z
Check the source to verify that this is the current version before use.

d. Procedure:

Into three separate 50 ml volumetric flasks, pipet 3.0 ml portions of Standard Preparation, Assay Preparation, and purified water (blank). To each flask, add 10.0 ml of Buffer Solution, 15.0 ml isopropyl alcohol, and 2.0 ml of Potassium Ferricyanide Solution. Mix well and then add 5.0 ml of 4-Aminoantipyrene Hydrochloride Solution to each flask; mix and dilute to volume with purified water. After 30 minutes, concomitantly measure the absorbances of the sample and standard at 500 nm, with a suitable spectrophotometer in 1-cm cells, and using the blank in the reference cell. Calculate as follows:

$$\frac{Au}{As} \times \frac{500C}{44} \times 100 = \frac{Au}{As} \times 1136C = \text{percent of required amount}$$

where,

Au = sample absorbance

As = standard absorbance

C = concentration of Standard Preparation, mg of potassium guaiacolsulfonate hemihydrate per ml.

4.6 Palatability test. A taste panel consisting of 10 members will be used to determine acceptability of samples. Samples will be prepared for testing (samples will be tested undiluted), coded, and served to panel members under controlled serving conditions, e.g., all samples will be of the same amount, and served at the same temperature; each panel member will receive an equal number of samples; the order of serving will be varied among panel members; an interval of at least five (5) minutes will elapse between successive samples and panel members will rinse their mouths with water (room temperature) after each sample; panel members will test without interference either from each other or from outsiders. The product offered shall be rated equal to or better than the Palatability Standard[1/], when determined by the taste panel, using the following 9-point hedonic rating scale. The average rating of the sample shall be equal to or greater than the average rating of the standard similarly prepared and tested.

1	2	3	4	5	6	7	8	9
Dislike extremely	Dislike very much	Dislike moderately	Dislike slightly	Neither like nor dislike	Like slightly	Like moderately	Like very much	Like extremely

1/ Palatability Standard is available, upon separate request to the contracting officer, Defense Personnel Support Center.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:25Z
Check the source to verify that this is the current version before use.

4.7 Examination for preparation for delivery. An examination shall be made to determine whether the packaging, packing, and marking comply with the requirements of Section 5. Defects shall be scored as specified in table IV. Sampling shall be in accordance with MIL-STD-105. The sample unit shall be one container fully prepared for delivery. The lot shall be the number of containers offered for inspection at one time. The inspection level shall be S-2 and the acceptable quality level shall be 4.0 defects per hundred units. For labeling, acceptance number shall be zero.

TABLE IV. Examination for preparation for delivery 1/

Examination	Defect
Containers	Not as specified.
Quantity	Quantity in interior packages and shipping containers not as specified.
Packaging and Packing Materials	Any component missing or damaged.
Workmanship	Inadequate application of components, such as incomplete closure of container flaps, tape improperly applied, or containers distorted.
Markings	Omitted, incorrect, illegible, improper size, location, sequence, or method of application.

1/ The above examination is only applicable to the unit, intermediate and exterior containers.

5. PACKAGING

5.1 Packaging requirements. The requirements for packaging shall be in accordance with all applicable requirements of PPP-C-186 (see 6.3) and as specified herein.

5.1.1 Unit of issue. One bottle containing one gallon (3.78 liters) or one bottle containing four fl. oz. (118 ml.) as specified constitutes one unit of issue.

5.2 Packaging. Packaging shall be level A or C as specified (see 6.1).

5.2.1 Level A.

Source: https://assist.dla.mil -- Downloaded: 202640-21T10:25Z
Check the source to verify that this is the current version before use.

5.2.1.1 Immediate container. The immediate container for size 1 and size 2 shall be a commercially available bottle considered acceptable by the Food and Drug Administration (FDA) for the product contained therein and shall be of such compatibility that neither the contents nor the container are altered in any way by each other.

5.2.1.2 Unit package.

5.2.1.2.1 Size 1. One gallon. Each glass bottle shall be packaged in a regular slotted corrugated, double-wall box having a minimum bursting strength of 200 pounds. The box shall be designed with a bottom pad and a top air-cell pad constructed of double-wall corrugated fiberboard, 200-pound test. The liner shall be not less than 8-1/2 inches in height and fabricated of nontest double-wall, C flute corrugated board. The liner shall be two thicknesses on all sides of the bottle. The top air-cell shall be folded and fitted over the closure end of the bottle and rest upon the liner, and then bent up toward the top of the box. The top air-cell shall be 27-7/16 inches long and 7-1/8 inches wide prior to folding. The inside dimensions of the box shall be 7-1/2 inches long by 7-3/8 inches wide and 14-1/2 inches in depth. Closure of the box shall be adequate to prevent spilling of contents under normal handling. As an alternate, each bottle shall be packaged in a 200 lb. test double wall B/C flute corrugated box having a two-piece molded or die-cut polystyrene insert. Closure of the box shall be adequate to prevent accidental opening.

5.2.1.2.2 Size 2. At the option of the contractor, a unit package may be furnished for each 4 fl. oz. bottle. If a unit package is furnished, each bottle shall be packaged as specified in paragraph 5.2.5 of PPP-C-186.

5.2.1.3 Intermediate package. An intermediate package shall not be required for size 1. Twelve size 2 units of issue shall be packaged as specified in paragraph 5.2.8 of PPP-C-186.

5.2.2 Level C. Units shall be packaged in standard commercial containers of the size and kind commonly used, which will afford the degree of protection required for shipment and use of the product for its intended purpose.

5.3 Packing. Packing shall be level A, B or C as specified (see 6.1).

5.3.1 Level A. Four unit packages containing size 1 or four intermediate packages (48 units) of size 2 shall be packed in an exterior container as specified in 5.4.4 of PPP-C-186.

Source: https://assist.dla.mil -- Downloaded: 202640-21T10:25Z
Check the source to verify that this is the currentversion before use.

5.3.2 Level B. Four unit packages, containing size 1 material, shall be packed in an exterior container as specified in 5.4.3.4 of PPP-C-186. Four intermediate packages (48 units) of size 2 material shall be packed in an exterior container as specified in 5.4.3.2 of PPP-C-186.

5.3.3 Level C. The subject commodity shall be packed in substantial commercial containers of the type, size, and kind commonly used for the purpose, so constructed as to insure acceptance and safe delivery by common or other carriers, at the lowest rate, to point of delivery called for in the contract or purchase order.

5.4 Marking.

5.4.1 Labeling. Labeling shall be in accordance with the requirements of the Federal Food, Drug and Cosmetic Act and shall include the information required below:

5.4.1.1 Immediate container. Each immediate container label shall bear the following information:

(a) label information in accordance with commercial practice.

(b) the date of manufacture prefixed by "MFD".

5.4.1.2 Unit package. Each unit package label shall bear the following:

(a) labeling information in accordance with commercial practice.

(b) the National Stock Number.

Notes: (1) The National Stock Number shall not be preceded by any prefix.

(2) Strip labeling may be used.

(3) Label adhesion test shall not apply.

(4) If a unit package is not supplied for size 2, the National Stock Number shall appear on the immediate container label.

5.4.2 Intermediate package. Each intermediate package shall be marked in accordance with 5.5.3 of PPP-C-186 except that the date of manufacture shall be shown in lieu of the date packed. Marking shall include the legend:

"STORE AT CONTROLLED ROOM TEMPERATURE (59° - 86° F)".

Source: https://assist.dla.mil -- Downloaded: 202640-21T10:25Z
Check the source to verify that this is the current version before use.

5.4.3 Exterior container. Exterior container shall be marked in accordance with 5.5.4 of PPP-C-186 except that the date of manufacture shall be shown in lieu of the date packed. Marking shall include the legend:

"STORE AT CONTROLLED ROOM TEMPERATURE (59° - 86° F)".

6. NOTES

6.1 Ordering data. Procurement documents should specify the following:

(a) Number and date of this specification.
(b) National Stock Number.
(c) Selection of applicable levels of packaging and packing (See 5.2 and 5.3).

6.2 This specification covers the following item(s):

National Stock No.	Item Identification	Specification Designation
6505-00-890-2010	PROMETHAZINE HYDROCHLORIDE, CHLOROFORM, IPECAC FLUIDEXTRACT, AND POTASSIUM GUAIACOLSULFONATE SYRUP, 1 gal (3.78 liters)	Size 1
6505-00-926-9026	PROMETHAZINE HYDROCHLORIDE, CHLOROFORM, IPECAC FLUIDEXTRACT, AND POTASSIUM GUAIACOLSULFONATE SYRUP, 4 fl oz (118 ml)	Size 2

6.3 All references to PPP-C-186 refer to Interim Federal Specification PPP-C-00186B which is mandatory for use until superseded by a revision.

6.4 This specification does not cover all types, classes, grades, or sizes of the commodity indicated by the title of this specification, or those which are commercially available, but is intended to cover the types which are normally procured to meet Military requirements.

Custodians:
Army - MD
Navy - MS
Air Force - 03

Preparing activity:
Defense Supply Agency - DM

Project No. 6505-1459

CS

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:25Z
Check the source to verify that this is the current version before use.