MILITARY SPECIFICATION

DESONIDE CREAM

This specification is approved for use by all Departments and Agencies of the Department of Defense.

1. SCOPE

1.1 This specification covers Desonide Cream.

1.2 Classification. The cream covered by this specification shall be the following sizes (see 6.2):

Size	1	-	15 Gram
Size	2	-	60 Gram

2. APPLICABLE DOCUMENTS

2.1 The following documents, of the issue in effect on date of invitation for bids or request for proposals, form a part of this specification to the extent specified herein:

SPECIFICATIONS

Federal

PPP-C-186	Containers, Packaging and Packing for Drugs, Chemicals, and Pharmaceuticals.

STANDARDS

Military

MIL-STD-105	Sampling Procedures and Tables for Inspection by Attributes.
MIL-STD-129	Marking for Shipment and Storage.

(Copies of specifications and standards required by contractors in connection with specific procurement functions should be obtained from the procuring activity or as directed by the contracting officer.)

FSC 6505

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:07Z
Check the source to verify that this is the current version before use.

2.2 Other publications. The following documents form a part of this specification to the extent specified herein. Unless otherwise indicated, the issue in effect on date of invitation for bids or request for proposals shall apply.

AMERICAN PHARMACEUTICAL ASSOCIATION PUBLICATION

National Formulary.

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE,
FOOD AND DRUG ADMINISTRATION

Federal Food, Drug, and Cosmetic Act and
Regulations Promulgated Thereunder.

(Application for copies should be addressed to the Food and Drug Administration, Washington, D. C. 20204.)

U. S. PHARMACOPEIAL CONVENTION, INC.

Pharmacopeia of the United States.

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

3. REQUIREMENTS

3.1 Material. Shall be Desonide Cream 0.05%.

3.1.1 Description. The desonide cream shall be smooth, clean, semi-solid, and no contamination by foreign material shall be noted.

3.1.2 Identification. The cream shall comply with the identity test as specified in 4.5.2.

3.1.3 pH. Using sufficient sample to cover the electrode tips, the cream shall have a pH of 4.3 to 5 using the USP method.

3.1.4 Assay. The cream shall assay to contain not less than 90 percent and not more than 110.0 percent of the required amount of desonide content as determined in 4.5.1.

3.2 Ingredients.

3.2.1 Desonide powder. The desonide powder used in the manufacture of the cream shall comply with the following requirements:

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:07Z
Check the source to verify that this is the current version before use.

3.2.1.1 Identification. The infrared absorbance spectrum (graph) of a potassium bromide dispersion of a sample of desonide exhibts maxima only at the same wavelengths as that of a similarly prepared reference sample, as kept on file by the manufacture . A copy of the standard infrared spectrum shall be submitted by the contractor, and shall be kept on file in the Technical & Quality Assurance Division, Directorate of Medical Material, Defense Personnel Support Center, for use if samples are submitted.

3.2.1.2 Loss on Drying. Shall not exceed 3.0 percent when 1 gram is dried in a vacuum oven where the vacuum is maintained for 3 hours at 60°C.

3.2.1.3 Residue on Ignition. Shall not be more than 1.0 percent when determined using the USP method.

3.2.1.4 Related Foreign Steroids. Shall not be more than 3 percent when determined as specified in 4.5.3.

3.2.1.5 Assay. Shall assay not less 97.0 and not more than 102.0 percent when determined as specified in 4.5.4.

3.2.2. Other ingredients. All other ingredients entering into the finished product shall be U.S.P. or N.F. quality, or, if not included in either of these compendia, the ingredients shall be of the highest pharmaceutical grade.

3.3 Contents. Filled immediate containers shall contain not less than 95 percent of the label content.

3.4 Potency Period. The potency period (Expiration Dating Period) shall be 36 months.

3.5 Delivery. Not less than 33 months of the potency period shall remain at the time of delivery to the Government.

3.6 Workmanship. Workmanship shall be first class throughout. The material and its containers shall be free from any defects which detract from their appearance or may impair their serviceability.

4. QUALITY ASSURANCE PROVISIONS

4.1 Supplier Responsibility for inspection. Unless otherwise specified in the contract or purchase order, the supplier is responsible for the performance of all inspection requirements as specified herein. Except as otherwise specified in the contract or order, the supplier may use his own or any other facilities suitable for the performance of the inspection requirements specified herein, unless disapproved by the Government. The Government reserves the right to perform any of the inspections set forth in the specification where such inspections are deemed necesssary to assure supplies and services conform to prescribed requirements.

Source: https://assist.dla.mil -- Downloaded: 20260-21T10:07Z
Check the source to verify that this is the current version before use.

4.1.1 Records of examinations and tests performed by or for the contractor shall be maintained by the contractor and made available to the Government upon the Government's request, at any time, or from time to time, during the performance of the contract and for a period of 3 years after delivery of the supplies to which such records relate.

4.1.2 No company supplying any ingredient(s) to the contractor will be considered an acceptable facility for the performance of any inspection requirements specified herein.

4.2 Samples. If (i) the quality assurance representative submits samples to the laboratory at the Defense Personnel Support Center, or (ii) the bidder or offeror is required by the terms of the Schedule or otherwise to submit samples to said laboratory, a quantity of 1 gram of the desonide powder used in the manufacture of each lot of the cream and 0.5 gram of the Reference Standard, accompanied by a certificate of analysis, listing all test results shall be submitted with the finished product to the Technical and Quality Assurance Division, Directorate of Medical Materiel, Defense Personnel Support Center, Attention: Quality Assurance Branch, 2800 South 20th Street, Philadelphia, PA 19101.

4.3 Sampling. Sampling shall be conducted in accordance with MIL-STD-105.

Table I. Sampling

For visual examination	Inspection level	AQL(percent) defective	Unit of product
Major	S-2	1.0	Filled tube

4.3.1 For raw material and end item testing. Noncompliance with any of the following is cause for rejection.

Table II. Raw Material & End Item Testing

Characteristic	Requirement paragraph	Test paragraph	Inspection level*	Minimum sample
Identifiction	3.1.2	4.5.2	S-2	5
pH	3.1.3	--	S-2	5
Assay	3.1.4	4.5.1	S-2	5
Contents	3.3	--	S-2	10
Ingredients			**	
Identification	3.2.1.1	--		
Loss on Drying	3.2.1.2			
Residue on Ignition	3.2.1.3	--		
Related Foreign Steroids	3.2.1.4	4.5.3		
Assay	3.2.1.5	4.5.4		

*Where possible, the same sample shall be used for too or more tests.
** Sufficient quantities of each of the raw materials shall be selected from each lot to perform tests indicated in applicable test methods. Failure of any sample to comply with test requirements shall be cause for rejection of the lot represented.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:07Z
Check the source to verify that this is the current version before use.

4.4 Examination. The Desonide Cream shall be examined to determine compliance with all requirements of this specification. Nonconformance will be permitted to the extent indicated in 4.3 and 4.3.1.

4.4.1 Classification of defects. Examination will be conducted in accordance with the following classification of defects:

Table III. Classification of defects

Categories	Defects*
Major	
101	Cream not homogeneous, not free from separation.
102	Cream not smooth.
103	Cream not free from foreign matter.
104	Cream not semi-solid.

*Inspection is not restricted to the classified possible defects listed above.

4.5 Tests.

4.5.1 Assay for desonide in the cream.

Preparation of Solvents and Reagents:

50% Ethanol/Chloroform v/v: Mix equal volumes of ethanol (2B) and chloroform.

Tetramethylammonium Hydroxide 2% v/v: Dilute 5.0 ml of the 10% aqueous reagent to 25.0 ml with ethanol (2B). Prepare fresh daily.

Tetrazolium Blue Solution. 0.34% w/v: Dissolve 85 mg. tetrazolium blue in about 15 ml ethanol (2B) in a 25 ml volumetric flask by gentle warming (warm water, or steam bath). Cool the concentrate to room temperature, dilute to volume with ethanol (2B) and filter. Prepare fresh daily and protect from light.

Preparation of Solutions for Assay

Stock Standard Solution

Dissolve about 25 mg (accurately weighed) desfluorotriamcinolone acetonide (desonide) in 50% ethanol (2B)/chloroform and dilute to volume in a 50 ml volumetric flask. Prepare fresh daily.

Source: https://assist.dla.mil -- Downloaded: 20261O-21T10:07Z
Check the source to verify that this is the current version before use.

Working Standard Preparation.

Dilute 10.0 ml of the stock standard solution with 50% ethanol (2B)/chloroform to volume in a 50 ml volumetric flask.

Standard Sample Preparation.

Weigh about one gram of Desonide Cream, which does not contain desfluorotriamcinolone acetonide, into a 125 ml Erlenmeyer flask. Pipette 5.0 ml of the working standard solution into the flask and add about 25 ml 50% ethanol (2B)/chloroform. Carefully swirl the flask to disperse the cream, filter quantitatively through Whatman #40 filter paper into a 50 ml volumetric flask and dilute to volume with 50% ethanol (2B)/chloroform. . . . conc., ca. 10 mcg/ml.

Sample Preparation

Accurately weigh about one gram of the .05% desfluorotriamcinolone acetonide cream into a 125 ml Erlenmeyer flask. Add about 30 ml of 50% ethanol (2B)/chloroform solution and carefully swirl the flask to disperse the cream. Filter quantitatively through Whatman #40 filter paper into a 50 ml volumetric flask and dilute to volume with 50% ethanol (2B)/chloroform.

Vehicle Blank Preparation

Weight about 1 gram of Desonide Cream, which does not contain desfluorotriamcinolone acetonide, into a 125 ml Erlenmeyer flask and work up as described above for the sample.

Procedure

Into each of three 25 ml volumetric flasks pipette 20.0 ml aliquots of:

a. The standard sample solution (Std.)
b. the sample solution (spl.) and
c. the vehicle solution (V. Blk.).

To each of the flasks add 1.0 ml of the tetrazolium blue solution and mix thoroughly by gently swirling the flask. Then:

1. To the first flask add 1.0 ml of the tetramethylammonium hydroxide solution, immediately dilute to volume with chloroform, stopper and mix well.

2. Repeat step (1) with the remaining flasks so that dilution to volume for each is carried out at successive two minute intervals.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:07Z
Check the source to verify that this is the current version before use.

After exactly 15 minutes of color development in each of the solutions, read A_{520} (vs. ethanol (2B)) on a suitable colorimeter or spectrophotometer.

Calculation:
% (w/w) desfluorotriamcinolone acetonide (Desonide) in formulation =

$$\frac{(A_{520}\text{Spl} - A_{520}\text{ V. Blk.})\text{ (Wt. Standard, in mg.)}}{(A_{520}\text{std} - A_{520}\text{ V. Blk.})\text{ (Wt. Sample, in grams) (500)}}$$

4.5.2 Identification of desonide in the cream

Materials

Silica Gel: 70/325 mesh for column chromatography, E. Merck AG 7734 or equivalent of activity grade 2-3 (according to Brockmann & Schodder).

Miniature chromatographic column: Glass, disposable Pasteur capillary pipette or equivalent.

Chloroform: Reagent Grade (ACS), bottled.

Methanol (anhydrous): Reagent Grade (ACS).

Thin-Layer Plates: Glass (5 or 10 x 20 cm.) coated with silica gel (250 μ) containing a fluorescent indicator (254 nm).

Thin-Layer Chromatography Developing Chamber: For accommodating 10 x 20 cm plates.

Ultraviolet Light Source (254 nm): Chromato-Vue CC20, or equivalent.

Tetrazolium Blue ("B.T."): Nutritional Biochemicals Corp. (Cleveland, Ohio) or other source of good purity material.

Solutions and Reagents

Desonide Standard Solution - To a small vial accurately weigh about one milligram reference standard desonide. Dissolve in chloroform by the addition (graduated pipette) of a quantity of the solvent such that the final concentration of desonide is 500 ug/ml (i.e., 1.0 mg in 2/0 ml). Alternatively, accurately weigh 12.5 mg. desonide and make to 25 ml. with chloroform in a 25 ml volumetric flask.

Source: https://assist.dla.mil -- Downloaded: 20260-21T10:07Z
Check the source to verify that this is the current version before use.

Blue Tetrazolium Spray Reagent - One volume of 20% sodium hydroxide solution (aqueous) is combined with two volumes of 0.2% (2 mg/ml) blue tetrazolium (methanolic) solution just prior to use. The stock blue tetrazolium solution maintains adequate stability for long periods if stored under refrigeration.

Procedure

A. Miniature Chromatographic Column - Into a vertically supported Pasteur capillary pipette (disposable) place a small pledget of glass wool then a slurry consisting of silica gel (500 mg) and chloroform (sufficient for transfer). The column is now ready to receive the test cream/silica gel blend. It is not crucial that a solvent head be maintained.

B. Isolation - Weigh about 100 mg of the test cream and about 200 mg. of silica gel into a small beaker and blend to a free flowing powder with a stainless steel spatula. Slurry the blend with chloroform (5 ml) and transfer the slurry to the miniature chromatographic column with a disposable Pasteur pipette. Make the transfer complete using chloroform (5 ml in small portions), then wash the column with an additional 5 ml of chloroform. Discard the chloroform effluents. Wash the column with 10% methanol in chloroform (3 ml) collecting the effluent in a small vial. Evaporate the solution to dryness (stream of air or nitrogen) then dissolve the residue to 0.10 ml chloroform.* This is the test solution.

* Use a Mohr measuring pipette (0.1, 0.2 or 0.5 ml.) for this addition or equivalent.

C. Thin-Layer Chromatography- Apply 10 ul aliquots of the test solution and of the desonide reference standard solution at least 2 cm apart on an imaginary line 3 cm from and parallel to the end of the chromatographic plate. Score a line 18 cm from and parallel to the same end of the plate. Irrigate the plate with 5% methanol in chloroform in a developing chamber (tlc tank) until the solvent has advanced to the scored line. Remove the plate and allow it to air dry.

D. Detection - Either or both of the following detection methods may be used. Upon exposure to short wave length (254 nm) ultraviolet light desonide appears as a blue spot on a light background. Upon spraying the plate with blue tetrazolium reagent, desonide appears as a purple-blue spot on a light background.

Source: https://assist.dla.mil -- Downloaded: 202610-21T10:07Z
Check the source to verify that this is the currentversion before use.

E. Specification The detected ingredient of the test solution must appear at the same Rf as that of the reference standard desonide.

4.5.3 Related foreign steroids (ingredients).

Reagents and Apparatus

A Chromatographic Plates - Uniplates, 20cm by 20cm, silica gel BF, 250 microns thick (Analtech, Inc., Wilimington, Delaware) or equivalent.

B Developing Solvent
Mixture A - Mix 93 ml of methylene chloride with 7 ml of methanol.
Mixture B - Add 50 ml cyclohexane, 50 ml dioxane, and 10 ml purified water to a separatory funnel. Shake and allow the layers to separate.
Mix 75 ml of Mixture A and 75 ml of the upper layer of Mixture B.

C Chromatographic Chamber - Line the chromatographic chamber with suitable adsorbent paper. Pour 150 ml of the developing solvent into the chamber and allow to equlibrate 20-30 minutes.

Procedure

1. Preparation of the standard - Accurately weigh exactly 30 mg of the desonide standard and transfer to a 100 ml volumetric flask. Add acetone to dissolve, bring up to volume with acetone and mix well. Standard A. Dilute 10 ml of Standard A to 15 ml with acetone and mix well. Standard B. Dilute 5 ml of Standard A to 15 ml with acetone and mix well. Standard C.

2. Preparation of Sample - Make a solution containing exactly 20 mg per ml in acetone. If necessary, use gentle heat to dissolve sample.

3. Thin Layer Chromatography - At the line of origin, at about 2.5 cm from the bottom of the plate, spot the prepared Standards and Sample solutions at 2 cm intervals.
where: -

1 - 5 ul	Standard A (1.5 mcg)
2 - 5 ul	Standard B (1.0 mcg)
3 - 5 ul	Standard C (0.5 mcg)
4 - 5 ul	Sample (100 mcg)
5 - 10 ul	Sample (200 mcg)

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:07Z
Check the source to verify that this is the current version before use.

Place the glass plate into the developing chamber immediately after applying the solutions to the plates. Develop the plate until the solvent front has moved to a point 1 cm from the top of the plate. Air dry the developed plate for 30 minutes.

Results

View the dried plate under shortwave ultraviolet light and compare the intensity of any foreign related steroid spot with the standard spots. Since the sample is chromatographed at 100 and 200 mcg levels and the standards at 0.5, 1.0 and 1.5 mcg, an estimation of the intensity of any foreign related steroid present at a level of 0.25% to 1.5% can be made. If the intensity of any foreign related steroid is greater than the 1.5 mcg (Standard A spot), repeat the chromatograph using lower concentrations of the sample, unitl the intensity of the spot falls within the 0.5 to 1.5 mcg range.

4.5.4 Desonide Assay (Ingredients)

Material

1. Chromatographic Plate: Prepare a slurry from 30 gram of chromatographic silica gel containing a fluorescent indicator and calcum sulfate by mixing about 65 ml of a mixture of 5 volumes of water and 2 volumes of alcohol and spreading the slurry 0.25 mm thick on 20 x 20 em plate and allow the plate to dry at room temperature and store in a desiccator.

2. Developing Solvent: Mix 10 ml of methanol with 190 ml of chloroform.

3. Chromatographic Chamber: Line a standard thinlayer chromatographic chamber with suitable absorbent paper, pour 150 ml of developing solvent into the chamber, equilibrate for 30 minutes.

Standard Preparation

Weigh accurately about 25 mg of Desonide Reference Standard into a 50 ml volumetric flask and dilute to volume with chlorofrom. Pipette 5.0 ml of this solution into a 50 ml volumetric flask and dilute to volume with chlorofrom. Pipette 5.0 ml of this diluted solution into a centrifuge cone and evaporate to dryness with a stream of nitrogen.

Sample Preparation

Weight accurately about 25 mg of Desonide into a 50 ml volumetric flask and then proceed exactly as for the Standard preparation.

Source: https://assist.dla.mil -- Downloaded: 2026l0-21T10:07Z
Check the source to verify that this is the currentversion before use.

Procedure

Divide the area of the Chromatographic plate into three equal sections, the left and right sections to be used for the Sample preparation and Standard preparation respectively, and the center section for the blank. Remove about 0.5 cm of absorbant from each of the two sides of the plate and score the plate across its entire width at a point 15 cm from the origin. Quantitatively apply the residue from the Standard preparation, as a spot, 3.0 cm from the bottom of the appropriate section of the plate according to the following procedure. Add about 0.1 ml of methylene chloride to the centrifuge cone, reflux gently for about one-half minute in a warm water bath (about 45° C.), then cool, thus concentrating the residue to the tip of the cone. If necessary, employ a stream of dry nitrogen to further reduce the volume of the concentrate to about 25 µl. Transfer the concentrate to the origin spot with the aid of a disposable Pasteur pipette (9" long) using capillary action for withdrawal and discharge, taking care not to disturb the silica gel layer. Rinse the Pasteur pipette with about 50 µl of methylene chloride, collecting the rinse in the centrifuge cone. Reflux, cool and employ, as necessary, dry nitrogen evaporation to keep the volume of the rinse small. Apply the rinse solution, as before, to the same spot. Repeat the rinse procedure one more time. Quantitatively apply the residue from the Sample preparation to the appropriate section. Irrigate the plate in the Chromatographic chamber until the solvent has moved 15 cm above the origin points. Remove the plate, allow the solvent to exaporate and locate the principal spot of the Standard preparation by viewing under a short-wavelength (254nm.) ultraviolet light. Mark this area as well as corresponding areas of the Sample preparation and blank section of the chromatogram.

Remove the silica gel from each area separately, either by scraping onto glazed weighing papers or by using a suitable vacuum collecting device. Quantitatively transfer the silica gel to a 25 ml volumetric flask with methanol and dilute to volume with methanol. Shake each flask for not less than two minutes. Filter a portion of each solution and concomitantly determine the absorbances of the clear filtrates from the Sample preparation, the Standard preparation and the silica gel blank at the wavelength of maximum absorbance at about 244 mm., with a suitable spectrophotometer, against methanol as the blank.

Calculations:

Calculate the percentage of Desonide taken by the formula

$$\% \text{ Desonide (w/w)} = \frac{Ws}{Wa} \times \frac{Au}{As} \times 100$$

in which Ws is the amount weighted in mg of Desonide Reference Standard in the Standard preparation, Wu is the amount weighed in mg of Desonide in the Sample preparation and Au and As are the absorbances of the solutions from the Sample preparation and the Standard preparation, respectively, both being corrected for the absorbance of the silica gel blank.

Source: https://assist.dla.mil -- Downloaded: 202610-21T10:07Z
Check the source to verify that this is the currentversion before use.

4.6 Examination for preparation for delivery. An examination shall be made to determine whether the packaging, packing, and marking comply with the requirements of Section 5. Defects shall be scored as specified in Table IV. Sampling shall be in accordance with MIL-STD-105. The sample unit shall be one container fully prepared for delivery. The lot shall be the number of containers offered for inspection at one time. The inspection level shall be S-2 and the acceptable quality level shall be 4.0 defects per hundred units.

Table IV. Examination for Preparation for Delivery

Examination	Defect
Containers	Not as specified.
Quantity	Quantity in interior packages and shipping containers not as specified.
Materials	Any component missing or damaged.
Workmanship	Inadequate application of components, such as incomplete closure of container flaps, tape improperly applied, or containers distorted.
Markings	Omitted, incorrect, illegible, improper size, location, sequence, or method of application.

5. PREPARATION FOR FELIVERY

5.1 Preparation for Delivery shall be in accordance with all applicable requirements of PPP-C-186, and as specified herein:

5.2 Packaging. Packaging shall be level A or C as specified (see 6.1).

5.2.1 Unit of issue. One tube containing 15 grams or 60 grams of cream as specified, constitutes one unit of issue.

5.2.2 Level A.

5.2.2.1 Immediate container. Immediate container shall be a commercially available ointment tube considered acceptable by the Food and Drug Administration (FDA) for the product contained therein and shall be of such compatibility that neither the contents nor the container are altered in any way by each other.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:07Z
Check the source to verify that this is the current version before use.

5.2.2.2 Unit package. Each tube shall be packaged in a carton or sleeve of appropriate size.

5.2.2.3 Intermediate package. Twelve (12) units of issue of size 1 shall be packaged in accordance with 5.2.12.1 of PPP-C-186 or twelve (12) units of issue of size 2 shall be packaged as specified in 5.2.13 of PPP-C-186.

5.2.3 Level C. Units shall be packaged in standard commercial containers of the size and kind commonly used, which will afford the degree of protection required for shipment and use of the product for its intended purpose.

5.3 Packing. Packing shall be level A, B or C as specified (see 6.1).

5.3.1 Level A. Four hundred thirty-two (432) units of issue of size 1 or one hundred forty-four (144) units of issue of size 2 shall be packed as specified in 5.4.4 of PPP-C-186.

5.3.2 Level B. Four hundred thirty-two (432) units of issue of size 1 or one hundred forty-four (144) units of issue of size 2 shall be packed as specified in 5.4.3.5.1 of PPP-C-186.

5.3.3 Level C. The subject commodity shall be packed in substantial commercial containers of the type, size, and kind commonly used for the purpose, so constructed as to insure acceptance and safe delivery by common or other carriers, at the lowest rate, to point of delivery called for in the contract or purchase order.

5.4 Marking.

5.4.1 Labeling. Labeling shall be in accordance with the requirements of the Federal Food, Drug and Cosmetic Act, and shall include the information required below.

5.4.1.1 Immediate container. Each immediate container label shall bear the following information:

(a) label information in accordance with commercial practice.

(b) the expiration date.

5.4.1.2 Unit package. Each unit package label shall bear the following:

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:07Z
Check the source to verify that this is the current version before use.

(a) labeling information in accordance with commercial practice.

(b) the national stock number.

NOTE: The National Stock Number shall not be preceded by any prefix.

Strip labeling may be used.

Label adhesion test shall not apply

5.4.2 Intermediate package. Intermediate package shall be marked as specified in 5.5.3 of PPP-C-186. Type 1 shelf life marking shall be shown as specified in MIL-STD-129. Marking shall include the legend: "STORE AT CONTROLLED ROOM TEMPERATURE (59° - 86°F.)"

5.4.3 Exterior container. Exterior container shall be marked as specified in 5.5.4 of PPP-C-186. Type I shelf like marking shall be shown as specified in MIL-STD-129. Marking shall include the legend: "STORE AT CONTROLLED ROOM TEMPERATURE (59° - 86°F.)"

6. NOTES

6.1 Ordering data Procurement documents should specify the following:

(a) Title, number and date of this specification.
(b) the National Stock No.
(c) Selection of applicable levels of packaging and packing (see 5.2 and 5.3).

6.2 This specification covers the following item:

National Stock Number	Item Identification
6505-00-148-6968	DESONIDE CREAM, 0.05%, 60 Gram
6505-00-148-6969	DESONIDE CREAM, 0.05%, 15 Gram

6.3 This specification does not cover all types, classes, grades, or sizes of the commodity indicated by the title of this specification, or those which are commercially available, but is intended to cover the types which are normally procured to meet Military requirements.

Custodians:
Army - MD
Navy - MS
Air Force - 03

Preparing activity:
Defense Supply Agency - DM

Project No. 6505-1349

Source: https://assist.dla.mil -- Downloaded: 202640-21T10:07Z
Check the source to verify that this is the current version before use.

STANDARDIZATION DOCUMENT IMPROVEMENT PROPOSAL

OMB Approval
No. 22-R255

INSTRUCTIONS: The purpose of this form is to solicit beneficial comments which will help achieve procurement of suitable products at reasonable cost and minimum delay, or will otherwise enhance use of the document. DoD contractors, government activities, or manufacturers/vendors who are prospective suppliers of the product are invited to submit comments to the government. Fold on lines on reverse side, staple in corner, and send to preparing activity. Comments submitted on this form do not constitute or imply authorization to waive any portion of the referenced document(s) or to amend contractual requirements. Attach any pertinent data which may be of use in improving this document. If there are additional papers, attach to form and place both in an envelope addressed to preparing activity.

DOCUMENT IDENTIFIER AND TITLE
MIL-D-37177 DESONIDE CREAM

NAME OF ORGANIZATION AND ADDRESS

CONTRACT NUMBER

MATERIAL PROCURED UNDER A

☐ DIRECT GOVERNMENT CONTRACT ☐ SUBCONTRACT

1. HAS ANY PART OF THE DOCUMENT CREATED PROBLEMS OR REQUIRED INTERPRETATION IN PROCUREMENT USE?
   A. GIVE PARAGRAPH NUMBER AND WORDING.

   B. RECOMMENDATIONS FOR CORRECTING THE DEFICIENCIES

2. COMMENTS ON ANY DOCUMENT REQUIREMENT CONSIDERED TOO RIGID

3. IS THE DOCUMENT RESTRICTIVE?

   ☐ YES ☐ NO *(If "Yes", in what way?)*

4. REMARKS

SUBMITTED BY *(Printed or typed name and address - Optional)*

TELEPHONE NO.

DATE

DD FORM 1 JAN 72 1426 REPLACES EDITION OF 1 JAN 66 WHICH MAY BE USED

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:07Z
Check the source to verify that this is the current version before use.

FOLD

DEFENSE PERSONNEL SUPPORT CENTER
2800 South 20th Street
Philadelphia, Pa. 19101

POSTAGE AND FEES PAID
DEFENSE SUPPLY AGENCY
DoD - 304

U.S.MAIL

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE $300

HEADQUARTERS, DEFENSE PERSONNEL SUPPORT CENTER
ATTN: DIRECTORATE OF MEDICAL MATERIEL, CODE ATT
2800 SOUTH 20TH STREET
PHILADELPHIA, PA. 19101

FOLD

Source: https://assist.dla.mil -- Downloaded: 202610-21T10:07Z
Check the source to verify that this is the current version before use.