MIL-B-36420A
6 October 1972
SUPERSEDING
MIL-B-36420
29 March 1965

MILITARY SPECIFICATION

BROMPHENIRAMINE MALEATE, PHENYLEPHRINE HYDROCHLORIDE, AND PHENYLPROPANOLAMINE HYDROCHLORIDE ELIXIR

This specification is mandatory for use by all Departments and Agencies of the Department of Defense

1. SCOPE

1.1 Scope. This specification covers two sizes (quantity of contents) of Brompheniramine Maleate, Phenylephrine Hydrochloride, and Phenylpropanolamine Hydrochloride Elixir.

1.2 Classification. The Brompheniramine Maleate, Phenylephrine Hydrochloride, and Phenylpropanolamine Hydrochloride Elixir shall be furnished in the size (quantity of contents) as specified for the appropriate Federal Stock No. (FSN). See 6.2.

Size I.- 1 gal (3.78 liters).

Size II.- 4 fl oz (118 cc).

2. APPLICABLE DOCUMENTS

2.1 The following documents, of the issue in effect on date of invitation for bids or request for proposals, form a part of this specification to the extent specified herein:

SPECIFICATION

PPP-C-00186a Containers, Packaging and Packing for Drugs, Chemicals, and Pharmaceuticals.

STANDARDS

MIL-STD-105 Sampling Procedures and Tables for Inspection by Attributes.

MIL-STD-129 Marking for Shipment and Storage.

(Copies of specification and standards required by contractors in connection with specific procurement functions should be obtained from the procuring activity or as directed by the contracting officer.)

FSC 6505

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

2.2 Other publications. The following documents form a part of this specification to the extent specified herein. Unless otherwise indicated, the issue in effect on date of invitation for bids or request for proposals shall apply.

AMERICAN PHARMACEUTICAL ASSOCIATION PUBLICATION

National Formulary.

(Application for copies should be addressed to the Mack Publishing Company, Easton, Pa. 18042.)

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE,
FOOD AND DRUG ADMINISTRATION

Federal Food, Drug, and Cosmetic Act and Regulations Promulgated Thereunder.

(Application for copies should be addressed to the Food and Drug Administration, Washington, D. C. 20402.)

U. S. PHARMACOPEIAL CONVENTION, INC.

Pharmacopeia of the United States.

(Application for copies should be addressed to the Mack Publishing Company, Easton, Pa. 18042.)

3. REQUIREMENTS

3.1 Material. Shall be a grape flavored, purple-colored elixir containing in each 5 cc, 4.0 mg of Brompheniramine Maleate, 5.0 mg of Phenylephrine Hydrochloride, 5.0 mg of Phenylpropanolamine Hydrochloride, and, in addition, shall contain 2.3 percent, by volume, of Alcohol. Shall be suitable for use as an antihistaminic nasal decongestant.

3.2 Assay. The elixir shall assay to contain not less than 93.0 percent and not more than 107.0 percent of the required amounts of Brompheniramine Maleate, Phenylephrine Hydrochloride, and Phenylpropanolamine Hydrochloride, when assayed as specified in 4.4.

3.2.1 The assay limits for the alcohol shall be not less than 2.19 percent and not more than 2.42 percent, by volume, when determined using the U.S.P. Alcohol Determination Test.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

3.3 Identification. The Brompheniramine Maleate and Phenylpropanolamine Hydrochloride in the elixir shall comply with the spectrophometric identification specified in 4.5. The Phenylephrine Hydrochloride in the elixir shall comply with the color identity test as specified in 4.5.

3.4 pH. The pH of the finished elixir shall be not less than 2.5 and not more than 3.5, when determined using a Beckman Model G or its equivalent.

3.5 Specific gravity. Specific gravity of the finished elixir shall be not less than 1.12 and not more than 1.15, using a hydrometer.

3.6 Refractive index. Refractive index of the finished elixir shall not be less than 1.390 and not more than 1.400 when determined using an Abbe Refractometer or its equivalent.

3.7 Color. The elixir shall be purple colored. The coloring agents used shall be approved by the Federal Food and Drug Administration (F.D.A.).

3.8 Flavor and palatability. The elixir shall be grape flavored, and shall be palatable and pleasant to the taste with no unpleasant aftertaste. Not later than the time specified for opening of bids or receipt of proposals, the offeror shall submit to the contracting officer six (6) individually packaged samples (each containing 4 fl oz) of Brompheniramine Maleate, Phenylephrine Hydrochloride, and Phenylpropanolamine Hydrochloride Elixir, representative of the product which the offeror proposes to furnish. Two (2) samples will be subjected to panel testing for a determination of palatability (see 4.6 Palatability test). The remaining samples will be used by cognizant Government inspection and quality assurance activities for determining compliance of supplies furnished hereunder with the palatability requirement. Approval as to palatability of any sample submitted by the offeror will not constitute approval of the sample as to any other requirement of this specification. The requirement for submission of samples for use in determining compliance with the palatability requirement may be waived, provided the offeror states, in his bid or proposal, that the product he proposes to furnish is the same product he has offered to the purchasing activity on a previous procurement and found to comply with the compatibility requirement.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

3.9 Active ingredients.

3.9.1 Brompheniramine maleate. The brompheniramine maleate, used in the preparation of the elixir, shall be in accordance with the tests, standards, and requirements of the N.F., including any supplements or revisions thereto, and, in addition, shall comply with the following:

3.9.1.1 Description. Shall be a white, odorless, crystalline powder.

3.9.2 Phenylpropanolamine hydrochloride. The phenylpropanolamine hydrochloride, used in the preparation of the elixir, shall be in accordance with the tests, standards, and requirements of the N.F., including any supplements or revisions thereto, and, in addition, shall comply with the following:

3.9.2.1 Description. Shall be a white, crystalling powder, having a slight aromatic odor.

3.9.3 Phenylephrine hydrochloride and alcohol. The phenylephrine hydrochloride and alcohol, used in the preparation of the elixir, shall be in accordance with the tests, standards, and requirements of the U.S.P., including any supplements or revisions thereto.

3.9.4 Other ingredients. Ingredients necessary to insure a suitable, stable product meeting all requirements of this specification shall be employed. All ingredients, used in the manufacture of the elixir, shall be of U.S.P. or N.F. quality or, if not included in either of these compendia, the ingredients shall be of the highest pharmaceutical grade.

3.10 Delivery. Not more than 6 months shall have elapsed from the date of manufacture of the product, to the date of delivery to the Government.

3.11 Workmanship. Workmanship shall be first class throughout. The material and its containers shall be free from defects which detract from their appearance or may impair their serviceability.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

4. QUALITY ASSURANCE PROVISIONS

4.1 Supplier responsibility for inspection. Unless otherwise specified in the contract or purchase order, the supplier is responsible for the performance of all inspection requirements as specified herein. Except as otherwise specified in the contract or order, the supplier may use his own or any other facilities suitable for the performance of the inspection requirements specified herein, unless disapproved by the Government. The Government reserves the right to perform any of the inspections set forth in the specification where such inspections are deemed necessary to assure supplies and services conform to prescribed requirements.

4.1.1 Records of examinations and tests performed by or for the contractor shall be maintained by the contractor and made available to the Government, upon the Government's request, at any time, or from time to time during the performance of the contract and for a period of 3 years after delivery of the supplies to which such records relate.

4.1.2 No company supplying any ingredient(s) to the contractor will be considered an acceptable facility for the performance of any inspection requirements specified herein.

4.2 Lot. For purposes of this specification, a lot, batch, or control is that single, uniform, and homogeneous quantity of the elixir, produced from one formulation, subjected to the same compounding and manufacturing operation, and filled into final containers.

4.3 Sampling. Sampling shall be conducted in accordance with the procedures set forth in MIL-STD-105, with an acceptable quality level (AQL) of 1.0 percent defective for major defects and 2.5 percent defective for minor defects.

4.4 Tests.

4.4.1 Assay for active ingredients in finished product.

4.4.1.1 Brompheniramine maleate.

Test reagents.

Aniline Solution, 4 percent: Mix 4 ml of aniline, reagent grade (colorless) with sufficient denatured alcohol to make 100 ml.

Potassium Acid Phthalate Solution, 2 percent: Dissolve 2 grams of potassium acid phthalate in sufficient purified water to make 100 ml and adjust to pH 6 with 0.1N sodium hydroxide.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

Brompheniramine standard: Dissolve 200 mg of N.F. Reference Standard Brompheniramine Maleate in sufficient 0.1N hydrochloric acid to make 100 ml. Adjust to pH 6 with 0.1N sodium hydroxide. Mix and transfer 10 ml (20 mg) of this solution to a 50-ml volumetric flask. This standard contains 0.4 mg of brompheniramine maleate per ml.

Cyanogen bromide reagent: Dissolve 25 grams of cyanogen bromide in 250 ml of purified water in a mixing cylinder. (Caution: Prepare this solution under the fume hood. Do not inhale the fumes).

Blank solution: Dissolve 50 mg of phenylephrine hydrochloride and 50 mg of phenylpropanolamine hydrochloride in 40 ml of potassium acid phthalate solution and adjust to pH 6 with 0.1N sodium hydroxide. Make to 50 ml with purified water.

Procedure:

Transfer 25 ml of the elixir to a separatory funnel using a calibrated pipet. Make the elixir basic with 2 ml of 25 percent sodium hydroxide and extract five times using 20-ml portions of chloroform. Combine all the chloroform extracts into one funnel and wash twice with 5 ml of saturated sodium chloride solution. Filter the chloroform into a 150-ml beaker; evaporate the chloroform to approximately 25 ml on a steam bath. Add to the flask 25 ml of 0.1N hydrochloric acid and continue evaporation until all the chloroform is evaporated. Adjust the solution to pH 6 using 0.1N sodium hydroxide. Quantitatively transfer the solution to a 50-ml volumetric flask and make to exactly 50 ml with purified water. To one Klett colorimeter tube, transfer 5 ml of brompheniramine standard. Transfer 5 ml of the test solution to a second tube. Transfer 5 ml of the blank solution to a third Keltt tube. To all three Klett tubes, add 5 ml of potassium acid phthalate solution and 1 ml of cyanogen bromide solution. Allow to stand exactly 15 minutes. Zero the instrument using purified water and a blue No. 42 Klett filter. Add 1 ml of aniline solution to each tube. Read the optical density of each tube periodically, until a maximum color is obtained.

Calculate the percent of brompheniramine maleate per 5 ml.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

4.4.1.1.1 As an alternate, the brompheniramine maleate in the finished product shall be assayed by the following method:

Test reagents:

4% Sodium hydroxide. Dissolve 4 grams of sodium hydroxide pellets with purified water to make 100 ml.

1N Hydrochloric acid. Dilute 95 ml of reagent grade hydrochloric acid with purified water to make 1000 ml.

Standard preparation. Dissolve exactly 80 mg of brompheniramine maleate Reference Standard, N.F., and 100 mg of phenylpropanolamine hydrochloride powder, N.F., in a separatory funnel with 10 ml of purified water. Add 10 ml of 4 percent sodium hydroxide and extract with eight 20-ml portions of chloroform. Filter the extracts into a 200-ml volumetric flask and make to volume with additional chloroform and mix. Transfer 10 ml to a 100-ml volumetric flask. Make to volume with chloroform and mix. This is the working standard.

Procedure:

Transfer exactly 5 ml of elixir to a separatory funnel and add 5 ml of 1N hydrochloric acid. Extract with five 40-ml portions of chloroform. Discard the chloroform and add 10 ml of 4 percent sodium hydroxide to the aqueous phase. Extract with four 20-ml portions of chloroform and filter the extracts into a 100-ml volumetric flask. Make to volume with additional chloroform. Obtain the absorbance of the sample and standard at 263 mu, using 1 cm quartz cells with chloroform in a reference cell.

Calculate the percent brompheniramine maleate per 5 ml.

4.4.1.2 Phenylephrine hydrochloride.

Test reagents.

4-aminoantipyrine solution: Prepare a 0.5 percent aqueous solution.

Potassium ferricyanide solution: Prepare a 2 percent aqueous solution.

Sodium bicarbonate buffer. Prepare a 0.084 percent aqueous solution.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

Phenylephrine hydrochloride: Prepare a standard of phenylephrine hydrochloride by dissolving 100 mg U.S.P. Reference Standard Phenylephrine Hydrochloride in 100 ml of purified water (1 mg/ml).

Procedure:

Accurately transfer 1 ml of the elixir to a 100-ml volumetric flask. Add to a second 100-ml volumetric flask, 1.0 ml of the phenylephrine hydrochloride standard solution. To both flasks, add 5 ml of potassium ferricyanide solution followed by 5 ml of 4-aminoantipyrine solution. Dilute both solutions to 50 ml with sodium bicarbonate buffer and allow to stand for 15 minutes.

Transfer a portion of these solutions to separate Klett colorimeter tubes and obtain the optical density of the test solution and the standard solution using Klett No. 54 green filter.

Calculate the percent of phenylephrine hydrochloride per 5 ml.

4.4.1.2.1 As an alternate, the phenylephrine hydrochloride in the finished elixir shall be assayed by the following method:

Test reagents.

4-aminoantipyrine solution. Weigh accurately 508 mg of 4-aminoantipyrine and dissolve in sufficient purified water to make 100 ml. Solution must be fresh.

Potassium ferricyanide solution. Dissolve 2 grams of potassium ferricyanide in sufficient purified water to make 100 ml. Solution must be fresh.

Sodium bicarbonate buffer. Dissolve 840 mg of sodium bicarbonate in sufficient purified water to make 1000 ml.

Phenylephrine hydrochloride. Prepare a standard of phenylephrine hydrochloride by dissolving 100 mg U.S.P. Reference Standard Phenylephrine Hydrochloride in 100 ml of purified water (1 mg/ml).

Procedure:

Accurately transfer 1 ml of the elixir to a 200-ml volumetric flask. Add to a second 200-ml volumetric flask 1.0 ml of the phenylephrine hydrochloride standard solution. To both flasks, add 5 ml of potassium ferricyanide solution followed by 5 ml of

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

4-aminoantipyrine solution. Dilute both solutions to 50 ml with sodium bicarbonate buffer and allow to stand for one hour.

Transfer a portion of each solution to separate 1 cm spectrophotometer cells and obtain the absorbance of each at 500 mu.

Calculate the percent phenylephrine hydrochloride per 5 ml.

4.4.1.3 Phenylpropanolamine hydrochloride.

Test reagent. 5 percent phosphomolybdic acid aqueous solution.

Procedure:

Transfer 50 ml of the elixir to a 150-ml beaker and add 10 ml of 5 percent phosphomolybdic acid solution. Mix well and centrifuge. Pour the supernatant liquid into a separatory funnel. Wash the residue once with 50 ml of purified water and centrifuge again. Add the supernatant liquid to the first portion of supernatant contained in the separatory funnel. Make the solution distinctly basic with 25 percent sodium hydroxide and extract four times with 25 ml portions of chloroform. Filter the collected extracts through chloroform saturated filter paper into a 250-ml Erlenmeyer flask. Add 50 ml of 0.01N sulfuric acid and a few drops of methyl red indicator. Evaporate the chloroform, cool, and titrate the sulfuric acid to a salmon pink endpoint with 0.01N sodium hydroxide.

Calculate the percent of Phenylpropanolamine Hydrochloride per 5 ml.

4.4.1.3.1 As an alternate, the phenylpropanolamine hydrochloride in the finished elixir shall be assayed by the following method:

Test reagents.

0.01N Perchloric acid. Mix 3.5 ml of reagent grade perchloric acid with four liters of glacial acetic acid. Standardize according to U.S.P. XVI procedure, page 1080.

Indicator. Dissolve 100 mg of quinaldine red in 10 ml of glacial acetic acid.

25% Sodium hydroxide. Dissolve 25 grams of reagent grade sodium hydroxide in purified water to make 100 ml.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

Procedure:

Pipet 25 ml of elixir into a separatory funnel. Extract with five 50-ml portions of chloroform. Wash chloroform with 10 ml of purified water and add to aqueous phase. Discard the chloroform. Make aqueous phase basic with 25 percent sodium hydroxide and extract with four 40-ml portions of chloroform. Wash with about 10 ml of purified water, filter the chloroform into an Erlenmeyer flask, add 50 ml glacial acetic acid and 5 ml acetic anhydride. Add three drops of quinaldine red indicator and titrate with 0.01N perchloric acid.

Calculate the percent phenylpropanolamine hydrochloride per 5 ml.

4.5 Identity.

4.5.1 Identity (ultraviolet curve) for Phenylpropanolamine Hydrochloride. Pipet 5 ml of elixir into a separatory funnel. Extract with five 40-ml portions of $CHCl_3$. Discard the $CHCl_3$. Add 5 ml of 4 percent NaOH to the sample in the separatory funnel and extract with four 25-ml portions of $CHCl_3$. Combine the extracts by filtering into a clean, dry, separatory funnel. Pipet 15 ml of 0.1N H_2SO_4 into the combined extracts. Shake vigorously. Discard the $CHCl_3$. Transfer the acid extract to a beaker.

Concurrently prepare a standard. Weigh 100 mg of phenylpropanolamine hydrochloride powder, N.F., and transfer to a separatory funnel containing 10 ml of 4 percent NaOH. Extract with five 40-ml portions of $CHCl_3$. Combine the extracts by filtering into a 200-ml volumetric flask. Dilute to volume with $CHCl_3$. Pipet 10 ml of the sample in $CHCl_3$ to a clean, dry, separatory funnel. Pipet 15 ml of 0.1N H_2SO_4 into the separatory funnel. Shake vigorously and discard the $CHCl_3$.

Scan the sample and standard on a recording spectrophotometer as follows:

Light Path - 1 cm silica cell
Scan Range - 300-230 mu
Reference - 0.1N H_2SO_4

For identification, compare the points of maximum absorbance for sample and standard.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

4.5.2 Identity (ultraviolet curve) for brompheniramine maleate. Make sample basic, and extract brompheniramine and phenylpropanolamine bases. Since both amines are extracted, a standard for quantitative and qualitative comparison shall be prepared containing the theoretical concentrations of both. It will be noted that the absorptivity of phenylpropanolamine will be much lower than for brompheniramine at the point of maximum absorbance of brompheniramine.

4.5.3 Phenylephrine hydrochloride identification. The phenylephrine hydrochloride has a phenyl ring with an unsubstituted position para to the CH on the ring and thus will react with 4-aminoantipyrine in the presence of alkaline oxidizing agent to form an antipyrine dye. The color produced shall at first be a dark red cherry color that soon fades to a rust color which shall remain for a period of about 2 hours only.

4.6 Palatability test. A taste panel consisting of 10 members will be used to determine acceptability of samples. Samples will be prepared for testing (samples will be tested undiluted unless otherwise specified in this document), coded, and served to panel members under controlled serving conditions, e.g., all samples will be of the same amount, and served at the same temperature; each panel member will receive an equal number of samples; the order of serving will be varied among panel members; an interval of at least five (5) minutes will elapse between successive samples and panel members will rinse their mouths with water (room temperature) after each sample; panel members will test without interference either from each other or from outsiders. The product offered shall be rated equal to or better than the FSN 6505-890-1892 and FSN 6505-935-4095 Palatability Standard,* when determined by the taste panel, using the following 9-point hedonic rating scale. The average rating of the sample shall be equal to or greater than the average rating of the standard, similarly prepared and tested.

1	2	3	4	5	6	7	8	9
Dislike extremely	Dislike very much	Dislike moderately	Dislike slightly	Neither like nor dislike	Like slightly	Like moderately	Like very much	Like extremely

*The FSN 6505-890-1892 and FSN 6505-935-4095 Palatability Standards are available, upon separate request to the contracting officer, Defense Personnel Support Center.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

5. PREPARATION FOR DELIVERY

5.1 Preparation for delivery shall be in accordance with all applicable requirements of Interim Federal Specification PPP-C-00186a, dated 15 May 1969, and Amendment-1, dated 27 October 1969, and as specified herein:

5.1.1 Immediate containers. Shall comply with the following classification:

GROUP A	CLASS 1	TYPE e	STYLE 1	GRADE 1
CLOSURE A				SEAL A

5.2 Labeling. Labeling shall be in accordance with the requirements of the Federal Food, Drug, and Cosmetic Act, and shall include the information as specified for the appropriate FSN.

For FSN 6505-890-1892 - See page 13.

For FSN 6505-935-4095 - See page 14.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

5.2.1 Immediate containers.

5.2.1.1 For FSN 6505-890-1892 (Size I). Each immediate container label for FSN 6505-890-1892 shall bear the following information. However, the information is not required to appear in the sequence indicated:

(a) the item name designated as "BROMPHENIRAMINE MALEATE, PHENYLEPHRINE HYDROCHLORIDE, AND PHENYLPROPANOLAMINE HYDROCHLORIDE ELIXIR"

(b) the name and quantity of each active ingredient for every 5 cc of elixir designated as "Brompheniramine Maleate 4.0 mg, Phenylephrine Hydrochloride, 5.0 mg, and Phenylpropanolamine Hydrochloride 5.0 mg"

(c) the quantity of contents designated as "1 gal (3.78 liters)"

(d) the Federal Stock No. designated as "6505-890-1892" or "Stock No. 6505-890-1892"

(e) the lot or control number

(f) the name and address of the manufacturer. When the manufacturer is not the contractor, the name and address of the contractor shall also appear.

When both names are placed on the label, the following designations shall precede the names:

"MFR" for the manufacturer and
"CONTR" for the contractor.

(g) the percentage of alcohol

(h) the statement "Caution: Federal law prohibits dispensing without prescription."

(i) the date of manufacture.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

5.2.1.2 For FSN 6505-935-4095 (Size II). Each immediate container for FSN 6505-935-4095 shall bear the following information. However, the information is not required to appear in the sequence indicated:

(a) the item name designated as "BROMPHENIRAMINE MALEATE, PHENYLEPHRINE HYDROCHLORIDE, AND PHENYLPROPANOLAMINE HYDROCHLORIDE ELIXIR"

(b) the name and quantity of each active ingredient for every 5 cc of elixir designated as "Brompheniramine Maleate 4.0 mg, Phenylephrine Hydrochloride, 5.0 mg, and Phenylpropanolamine Hydrochloride 5.0 mg"

(c) the quantity of contents designated as "4 fl oz (118 cc)"

(d) the Federal Stock No. designated as "FSN 6505-935-4095" or "Stock No. 6505-935-4095"

(e) the lot or control number

(f) the name and address of the manufacturer. When the manufacturer is not the contractor, the name and address of the contractor shall also appear.

When both names are placed on the label, the following designations shall precede the names:

"MFR" for the manufacturer and
"CONTR" for the contractor.

(g) the percentage of alcohol

(h) the statement "Caution: Federal law prohibits dispensing without prescription."

(i) the date of manufacture.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

5.3 Packaging and packing.

5.3.1 Unit of issue. One bottle of Size I (FSN 6505-890-1892) or one bottle of Size II (FSN 6505-935-4095), as specified, constitutes one unit of issue.

5.3.2 Unit package. At the option of the contractor, each unit shall be packaged as specified in 5.2.5 of PPP-C-00186a.

5.3.3 Procedure code. Procedure code No. 8 of PPP-C-00186a shall apply for Size I (FSN 6505-890-1892) and Procedure code No. 4 of PPP-C-00186a shall apply for Size II (FSN 6505-935-4095), except that in Table I, under "Column 2", delete "12" and substitute "24" and under column 4, delete "12/96" and substitute "24/96."

5.4 Marking.

5.4.1 Unit package. When furnished, each unit package shall bear the same information as required for the label of its immediate container.

5.4.2 Intermediate package. In paragraph 5.5.3 of PPP-C-00186a, add: "Date of manufacture shall be shown in lieu of date packed."

5.4.3 Exterior container. In paragraph 5.5.4 of PPP-C-00186a, add: "Date of manufacture shall be shown in lieu of date packed."

6. NOTES

6.1 Ordering data. Procurement documents should specify the following:

(a) Number and date of this specification
(b) the Federal Stock No.
(c) applicable levels of packaging and packing (see PPP-C-00186a).

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

6.2 This specification covers the following items:

FSN	Item Identification	Specification designation
6505-890-1892	BROMPHENIRAMINE MALEATE, PHENYLEPHRINE HYDROCHLORIDE, AND PHENYLPROPANOLAMINE HYDROCHLORIDE ELIXIR, 1 gal (3.78 liters)	I
6505-935-4095	BROMPHENIRAMINE MALEATE, PHENYLEPHRINE HYDROCHLORIDE, AND PHENYLPROPANOLAMINE HYDROCHLORIDE ELIXIR, 4 fl oz (118 cc)	II

Custodians:

Army - MD
Navy - MS
Air Force - 03

Preparing activity:
Defense Supply Agency - DM

Project No. 6505-1300

Review activities:

Army - MD
Navy - MS
Air Force - 03

Review information is current as of the date of this document. For future coordination of changes to this document, draft circulation should be based on the information in the current DODISS.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.

STANDARDIZATION DOCUMENT IMPROVEMENT PROPOSAL

OMB Approval
No. 22-R255

INSTRUCTIONS: The purpose of this form is to solicit beneficial comments which will help achieve procurement of suitable products at reasonable cost and minimum delay, or will otherwise enhance use of the document. DoD contractors, government activities, or manufacturers/vendors who are prospective suppliers of the product are invited to submit comments to the government. Fold on lines on reverse side, staple in corner, and send to preparing activity. Comments submitted on this form do not constitute or imply authorization to waive any portion of the referenced document(s) or to amend contractual requirements. Attach any pertinent data which may be of use in improving this document. If there are additional papers, attach to form and place both in an envelope addressed to preparing activity.

DOCUMENT IDENTIFIER AND TITLE

Military Specification MIL-B-36420A

NAME OF ORGANIZATION AND ADDRESS

CONTRACT NUMBER

MATERIAL PROCURED UNDER A

☐ DIRECT GOVERNMENT CONTRACT ☐ SUBCONTRACT

1. HAS ANY PART OF THE DOCUMENT CREATED PROBLEMS OR REQUIRED INTERPRETATION IN PROCUREMENT USE?
   A. GIVE PARAGRAPH NUMBER AND WORDING.

   B. RECOMMENDATIONS FOR CORRECTING THE DEFICIENCIES

2. COMMENTS ON ANY DOCUMENT REQUIREMENT CONSIDERED TOO RIGID

3. IS THE DOCUMENT RESTRICTIVE?

   ☐ YES ☐ NO *(If "Yes", in what way?)*

4. REMARKS

SUBMITTED BY *(Printed or typed name and address - Optional)*

TELEPHONE NO.

DATE

DD FORM 1 JAN 72 1426 REPLACES EDITION OF 1 JAN 66 WHICH MAY BE USED

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T08:34Z
Check the source to verify that this is the current version before use.