MIL-P-37149A
2 June 1975
SUPERSEDING
MIL-P-37149
10 December 1974

MILITARY SPECIFICATION

PROPOXYPHENE NAPSYLATE AND ACETAMINOPHEN TABLETS

This specification is approved for use by all Departments and Agencies of the Department of Defense.

1. SCOPE

1.1 Scope. This specification covers Propoxyphene Napsylate and Acetaminophen Tablets.

1.2 Classification. The tablets covered by this specification shall be the following sizes (See 6.2):

Size 1 - Bottle of 500s
Size 2 - Bottle of 30s
Size 3 - Individually sealed 100s

2. APPLICABLE DOCUMENTS

2.1 The following documents, of the issue in effect on date of invitation for bids or request for proposals, form a part of this specification to the extent specified herein:

SPECIFICATIONS

Federal

PPP-C-186	Containers, Packaging and Packing for Drugs, Chemicals, and Pharmaceuticals. (See 6.3)

STANDARDS

Federal

FED-STD-140	Tablets (for Medicinal Purposes)

FSC 6505

Source: https://assist.dla.mil -- Downloaded: 202610-21T10:04Z
Check the source to verify that this is the current version before use.

Military

MIL-STD-105	Sampling Procedures and Tables for Inspection by Attributes.
MIL-STD-129	Marking for Shipment and Storage.

(Copies of specifications and standards required by contractors in connection with specific procurement functions should be obtained from the procuring activity or as directed by the contracting officer.)

2.2 Other publications. The following documents form a part of this specification to the extent specified herein. Unless otherwise indicated, the issue in effect on date of invitation for bids or request for proposals shall apply.

AMERICAN PHARMACEUTICAL ASSOCIATION PUBLICATION

National Formulary.

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE, FOOD AND DRUG ADMINISTRATION

Federal, Food, Drug, and Cosmetic Act and Regulations Promulgated Thereunder.

(Application for copies should be addressed to the Food and Drug Administration, Washington, DC 20204.)

U.S. PHARMACOPEIAL CONVENTION, INC.

Pharmacopeia of the United States.

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

3. REQUIREMENTS

3.1 Material. Shall be Propoxyphene Napsylate and Acetaminophen Tablets, and shall be in accordance with all applicable requirements of FED-STD-140 and as specified herein:

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:04Z
Check the source to verify that this is the current version before use.

S2. Classification. Shall be type I.

S5.3 The following additional requirements and tests are added to this paragraph:

The tablets shall contain 100 mg Propoxyphene Napsylate and 650 mg of Acetaminophen per tablet, within the designated assay limits for the tablets.

The tablets shall assay to contain not less than 90.0 percent and not more than 110.0 percent of the required amount of Propoxyphene Napsylate and Acetaminophen when assayed as specified in 4.3.1 and 4.3.2.

The tablets shall comply with the identity test specified in 4.3.3.

S6.4.9.1 Disintegration. Shall comply with the following requirements when tested in accordance with the U.S.P. Disintegration Test for Coated Tablets:

Test Fluid	Total time
Purified Water (room temperature)	5 Minutes
Simulated Gastric Fluid, T.S. (37° C.)	30 Minutes
Simulated Intestinal Fluid, T.S. (37° C.)	15 Minutes

S6.9.1 Classification of defects. Major A No. 109 defect in table VI shall not apply. Major A No. 109 in Table IV (applicable to tablet prior to film coating) shall apply.

3.2 Ingredients. All ingredients used in the manufacture of the tablets shall comply with S5.1 of FED-STD-140.

3.3 Potency period. The potency period (Expiration dating period) shall be 36 months.

4. QUALITY ASSURANCE PROVISIONS

4.1 Supplier responsibility for inspection. Unless otherwise specified in the contract or purchase order, the supplier is responsible for the performance of all inspection requirements as specified herein. Except as otherwise specified in the contract or order, the supplier may use his own or any other other facilities suitable for the performance of the inspection

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:04Z
Check the source to verify that this is the current version before use.

requirements specified herein, unless disapproved by the Government. The Government reserves the right to perform any of the inspections set forth in the specification where such inspections are deemed necessary to assure supplies and services conform to prescribed requirements.

4.1.1 Records of examinations and tests performed by or for the contractor shall be maintained by the contractor and made available to the Government, upon the Government's request, at any time, or from time to time, during the performance of the contract and for a period of 3 years after delivery of the supplies to which such records relate.

4.1.2 No company supplying any ingredient(s) to the contractor will be considered an acceptable facility for the performance of any inspection requirements specified herein.

4.2 Sampling and Inspection. The classification of ~~defects as shown in FED-STD-140 shall be applicable.~~

4.2.1 Samples. If (i), the quality assurance representative submits samples to the laboratory at the Defense Personnel Support Center, or (ii) the bidder (offeror) is required by the terms of the Schedule or otherwise to submit samples to said laboratory, then a quantity of 5 grams of each lot of tablets and 1 gram of Propoxyphene Napsylate reference standard, accompanied by a certificate of analysis listing all test results, shall be submitted with the finished product to the Technical & Quality Assurance Division, Directorate of Medical Materiel, Defense Personnel Support Center, Attention: Quality Assurance Branch, 2800 South 20th Street, Philadelphia, PA 19101.

4.3 Tests.

4.3.1 Assay for Propoxyphene Napsylate in tablets.

Reagents

1. Methanol, Anhydrous, A. R.
2. Chloroform, A. R.
3. Sodium Carbonate Solution, 5% (w/v) in water.

Source: https://assist.dla.mil -- Downloaded: 2026-04-21T10:04Z
Check the source to verify that this is the current version before use.

4. Diphenamid Internal Standard. Prepare a solution in chloroform containing 0.4 mg of Diphenamid per ml.
5. Propoxyphene Napsylate Reference Standard. Accurately weigh about 50 mg of Propoxyphene Napsylate Reference Standard and transfer to 100 ml volumetric flask. Accurately weigh about 325 mg of N.F. Acetaminophen reference standard, and transfer into the flasks containing the Propoxyphene Napsylate. Dissolve and dilute to 100 ml with Methanol and mix well.

Apparatus.

A suitable gas chromatograph and column, such as the Hewlett-Packard Model 402, or equivalent instrument, utilizing a flame ionization detector and equipped with a 4 foot by 3 mm inside diameter glass column packed with 0.5 percent OV-17 filter coated on Diatoport S 80/100 mesh. The column shall have been previously conditioned for use.

An electronic integrator.

Sample Preparation.

Weigh five (5) tablets and transfer into a 500 ml volumetric flask. (Prior to placing the tablets in the flask, break each tablet in two.)

Add about 200 ml methanol to the flask and swirl occasionally until the tablets have completely disintegrated. The film coating will not dissolve.

Ultrasonorate for about one minute. Dilute to mark with Methanol and mix well.

Transfer a 5.0 ml aliquot of the sample assay solution to a test tube. Add 5 ml of Na_2CO_3 solution into the tube. Pipet a 5.0 ml aliquot of the diphenamid internal standard into the tube, and stopper tightly. Rotate the tube at approximately 25 r.p.m. for about one hour to extract the propoxyphene into the chloroform phase. Allow the phases to separate, then filter through phase separation paper into a vial or other suitable container.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:04Z
Check the source to verify that this is the current version before use.

Standard Preparation

Extract 5.0 ml aliquot of the propoxyphene napsylate reference standard solution as described for the sample assay solution.

Approximate Gas Chromatographic Operating Conditions.

Column Temperature:	175° C.
Flash Heater Temperature	200° C.
Detector Temperature:	250° C.
Rotameter:	He 3.5, O_2 3.0, H_2 3.0
Range:	10^2
Attenuation:	8X
Sample Size:	3 µl
Retention Times: (Approximate)	
Diphenamid	4.8 minutes
Propoxyphene	6.3 minutes

Procedure.

Withdraw and inject 3 µl of the chloroform phase of the sample into the column in the gas chromatograph utilizing the conditions listed above. Slight adjustments may be made to obtain optimum results.

Similarly, withdraw and inject 3 µl of the chloroform phase of the standard into the column in the gas chromatograph.

Calculation:

$$\text{Peak Area Ratio (PAR) for Propoxyphene} = \frac{\text{Peak Area Propoxyphene}}{\text{Peak Area Diphenamid}}$$

mg Propoxyphene Napsylate per tablet =

$$\frac{\text{PAR Sample}}{\text{PAR Standard}} \times Cs \times \frac{500}{\text{Wt of Sample (Grams)}} \times \text{Average Tablet Wt (grams)}$$

Where Cs is the concentration of Propoxyphene Napsylate Reference standard in mg per ml.

Source: https://assist.dla.mil -- Downloaded: 202610-21T10:04Z
Check the source to verify that this is the currentversion before use.

4.3.2 Assay for Acetaminophen in the tablets.

Standard Preparation.

Accurately weigh about 325 mg of N.F. Acetaminophen reference standard and transfer to a 100 ml volumetric flask. Accurately weigh about 50 mg of Propoxyphene Napsylate and transfer to the same flask. Dissolve and dilute to 100 ml with methanol and mix well.

Accurately weigh five tablets, break each tablet in two, and transfer to a 1000 ml volumetric flask. Add about 400 ml of methanol to the flask and swirl until the tablets have completely disintegrated. Ultrasonerat for about one minute, dilute to mark with methanol and mix well.

Pipet 2.0 ml of the sample solution and 2.0 ml of the standard solution into separate 1000 ml volumetric flasks. Dilute to volume with purified water and mix well. Concomitantly determine the absorbances of the sample and standard solutions at the maximum peak of absorbance at approximately 244 nm, with a suitable spectrophotometer in 1 cm cells versus purified water as a blank.

Calculate the mg of Acetaminophen per tablet as follows:

$$\frac{Au}{As} \times Cs \times 1000 \times \frac{\text{Ave Tab Wt (grams)}}{\text{Sample Wt (grams)}} = \text{mg acetaminophen per tablet}$$

where:

Au = absorbance of sample solution
As = absorbance of standard solution
Cs - exact initial concentration of standard in mg per ml

4.3.3 Identification.

The presence of Propoxyphene Napsylate and Acetaminophen in the finished tablets shall be determined as follows:

(1) The retention time of the Propoxyphene in the assay (see 4.3.1) matches that of the standard.

(2) The maximum absorbance of the assay solution in the acetaminophen assay (see 4.3.2) is 244 $\pm$ 2 nm.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:04Z
Check the source to verify that this is the current version before use.

(3) The final assay solution in the Propoxyphene Napsylate assay (see 4.3.1) is dextrorotatory.

(4) The sample is substantially free of chloride when tested with silver nitrate in the presence of nitric acid.

4.4 Examination of preparation for delivery. An examination shall be made to determine whether the packaging, packing, and marking comply with the requirements of Section 5. Defects shall be scored as specified in Table I. Sampling shall be in accordance with MIL-STD-105. The sample unit shall be one container fully prepared for delivery. The lot shall be the number of containers offered for inspection at one time. The inspection level shall be S-2 and the acceptable quality level shall be 4.0 defects per hundred units.

Table I. Examination of preparation for delivery

Examination	Defect
Containers	Not as specified.
Quantity	Quantity in interior packages and shipping containers not as specified.
Packaging and packing materials	Any component missing or damaged.
Workmanship	Inadequate application of components such as incomplete closure of container flaps, tape improperly applied, or containers distorted.
Markings	Omitted, incorrect, illegible, improper size, location, sequence, or method of application.

5. PREPARATION FOR DELIVERY

5.1 Preparation for Delivery shall be in accordance with all applicable requirements of PPP-C-186, and as specified herein.

5.2 Packaging. Packaging shall be Level A or C as specified (see 6.1).

5.2.1 Unit of issue. One bottle containing 30 tablets or 500 tablets or one box containing 100 individually packaged tablets, as specified, constitutes one unit of issue.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:04Z
Check the source to verify that this is the current version before use.

5.2.2 Level A.

5.2.2.1 Immediate containers.

5.2.2.1.1 Immediate containers for size 1 shall be a commercially available immediate container considered acceptable by the Food and Drug Administration (FDA) for the product contained therein and shall be of such compatibility that neither the contents nor the container are altered in any way by each other.

5.2.2.1.2 Immediate containers for size 2 shall be a commercially available immediate container, with child-proof safety closure, considered acceptable by the Food and Drug Administration (FDA) for the product contained therein and shall be of such compatibility that neither the contents nor the container are altered in any way by each other.

5.2.2.1.3 Immediate containers for size 3 shall be a commercially available strip or blister package considered acceptable to the Food and Drug Administration (FDA) for the product contained therein and shall be of such compatibility that neither the contents nor the container shall be altered in any way by each other.

5.2.2.2 Unit package. At the option of the contractor, a unit package may be furnished for bottles containing 30 tablets or bottles containing 500 tablets. If furnished, each unit shall be packaged as specified in 5.2.5 of PPP-C-186. One hundred individually sealed tablets shall be packaged in a standard commercial container of appropriate size and design. Closure shall be adequate to prevent spilling of contents under normal handling.

5.2.2.3 Intermediate package. Ten unit packages, each containing one hundred individually packaged tablets or twelve bottles, each containing five hundred tablets, shall be packaged as specified in 5.2.8 of PPP-C-186. Twenty-five bottles, each containing thirty tablets, shall be packaged as specified in 5.2.7.1 of PPP-C-186.

5.2.3 Level C. Units shall be packaged in standard commercial containers of the size and kind commonly used, which will afford the degree of protection required for shipment and use of the product for its intended purpose.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:04Z
Check the source to verify that this is the current version before use.

5.3 Packing.

5.3.1 Level A. Four intermediate packages of bottles each containing 500 tablets, twenty intermediate packages of bottles each containing 30 tablets or eight intermediate packages of boxes, each containing 100 individually packaged tablets shall be packed as specified in 5.4.4 of PPP-C-186. Individually packaged tablets shall not be required to be packed Level B prior to being packed Level A.

5.3.2 Level B. Four intermediate packages of bottles each containing 500 tablets or eight intermediate packages of boxes each containing 100 individually sealed tablets shall be packed as specified in 5.2.8 of PPP-C-186. Twenty intermediate packages of bottles each containing 30 tablets shall be packed as specified in 5.2.7.1 of PPP-C-186.

5.3.3 Level C. The subject commodity shall be packed in substantial commercial containers of the type, size, and kind commonly used for the purpose, so constructed as to insure acceptance and safe delivery called for in the contract or purchase order.

5.4 Marking.

5.4.1 Labeling. Labeling shall be in accordance with the requirements of the Federal Food, Drug and Cosmetic Act, and shall include the information required below:

5.4.1.1 The label of the immediate container for size 1 and size 3 shall bear the following information:

(a) Label information in accordance with commercial practice.

(b) the expiration date.

5.4.1.2 The label of the unit package for size 1 and size 3 shall bear the following:

(a) labeling information in accordance with commercial practice.

(b) the National Stock Number.

Source: https://assist.dla.mil -- Downloaded: 202640-21T10:04Z
Check the source to verify that this is the currenversion before use.

NOTE: The National Stock Number shall not be preceded by any prefix.

Strip labeling may be used.

Label adhesion test shall not apply.

If a unit package is not supplied, the National Stock Number shall appear on the immediate container label.

5.4.1.3 The label of the immediate container for size 2 shall be a tri-part label consisting of the following:

(1) a tab portion which shall remain attached to the immediate container. This portion shall bear:

(a) the National Drug Code (NDC)
(b) the lot or control number
(c) the expiration date

NOTE: Other label information may be included.

(2) a tear-off portion adjacent to the tab portion or at the extreme end of the label bearing the following:

(a) the National Drug Code (NDC)
(b) the lot or control number
(c) the expiration date

No other labeling shall appear thereon.
This portion shall be easily removed for attachment to a prescription.

(3) the principal portion connected to the tear-off and/or the tab portion shall bear the following:

(a) the National Stock Number (which shall not be preceded by any prefix)
(b) all other information required by the Federal Food, Drug, and Cosmetic Act.

This portion shall also be easily removed.

Label adhesion test shall not apply to any portion of the tri-part label.

Source: https://assist.dla.mil -- Downloaded: 202610-21T10:04Z
Check the source to verify that this is the current version before use.

5.4.1.4 The label of the unit package for size 2 shall bear the following:

(a) labeling information in accordance with commercial practice.

(b) the National Stock Number.

NOTE: The National Stock Number shall not be preceded by any prefix.

Strip labeling may be used.

Label adhesion test shall not apply.

If a unit package is not supplied, the National Stock Number shall appear on the principal portion connected to the tear off and/or the tab portion of the label on each immediate container.

5.4.2 Intermediate package. Each intermediate package shall be marked in accordance with paragraphs 5.5.3 and 5.5.8.1 of PPP-C-186. Marking shall include the legend: "STORE AT CONTROLLED ROOM TEMPERATURE (59° - 86° F)".

5.4.3 Exterior container. Exterior container shall be marked in accordance with paragraphs 5.5.4 and 5.5.8.1 of PPP-C-186. Marking shall include the legend: "STORE AT CONTROLLED ROOM TEMPERATURE (59° - 86° F)".

66. NOTES

6.1 Ordering data. Procurement documents should specify the following:

(a) Number and date of this specification.
(b) the National Stock Number.
(c) Selection of applicable levels of packaging and packing (See 5.2 and 5.3).

6.2 This specification covers the following item(s):

National Stock Number	Item Identification
6505-00-111-8359	PROPOXYPHENE NAPSYLATE AND ACETAMINOPHEN TABLETS, 500s
6505-00-111-8364	PROPOXYPHENE NAPSYLATE AND ACETAMINOPHEN TABLETS, 30s
6505-00-111-8373	PROPOXYPHENE NAPSYLATE AND ACETAMINOPHEN TABLETS, INDIVIDUALLY SEALED, 100s

Source: https://assist.dla.mil -- Downloaded: 202640-21T10:04Z
Check the source to verify that this is the currenversion before use.

6.3 All references to PPP-C-186 refer to Interim Federal Specification PPP-C-00186B which is mandatory for use until superceded by a revision.

6.4 This specification does not cover all types, classes, grades, or sizes of the commodity indicated by the title of this specification, or those which are commercially available, but is intended to cover the types which are normally procured to meet Military requirements.

Custodians:
Army - MD
Navy - MS
Air Force - 03

Preparing activity:
Defense Supply Agency - DM

Project No. 6505-1379
cs

Source: https://assist.dla.mil -- Downloaded: 202610-21T10:04Z
Check the source to verify that this is the currentversion before use.