O-R-500C
August 18, 1970
SUPERSEDING
Int. Fed. Spec. O-R-00500B(GSA-FSS)
October 10, 1968 and
Fed. Spec. O-R-500
January 23, 1957

FEDERAL SPECIFICATION

RODENTICIDE, BAIT, ANTICOAGULANT

This specification was approved by the Commissioner, Federal Supply Service, General Services Administration, for the use of all Federal agencies.

1. SCOPE

1.1 Scope. This specification covers a rodenticide consisting of a bait and an anticoagulant which could be: diphacinone, fumarin, Pival, or warfarin.

2. APPLICABLE DOCUMENTS

2.1 The following documents, of the issue in effect on date of invitation for bids or request for proposal, form a part of this specification to the extent specified herein:

Federal Specifications:

N-O-41	- Oats; Rolled.
RR-S-366	- Sieve, Test.
JJJ-S-791	- Sugar, Refined and Brown; Beet or Cane.
PPP-B-636	- Box, Fiberboard.
PPP-C-96	- Cans, Metal, 28 Gage and Lighter.

Federal Standard:

Fed. Std. No. 123 - Marking for Domestic Shipment (Civil Agencies).

(Activities outside the Federal Government may obtain copies of Federal Specifications, Standards, and Handbooks as outlined under General Information in the Index of Federal Specifications and Supplements as issued, is for sale on a subscription basis by the Superintendent of Documents, U. S. Government Printing Office, Washington, D. C. 20402.

(Single copies of this specification and other Federal Specifications required by activities outside the Federal Government for bidding purposes are available without charge from Business Service Centers at the General Services Administration Regional Offices in Boston, New York, Washington, D. C., Atlanta, Chicago, Kansas City, Mo., Fort Worth, Denver, San Francisco, Los Angeles, and Seattle, Washington.

(Federal Government activities may obtain copies of Federal Specifications, Standards, and Handbooks and the Index of Federal Specifications and Standards from established distribution points in their agencies.)

Military Standards:

MIL-STD-105 - Sampling Procedures and Tables for Inspection by Attributes.
MIL-STD-129 - Marking for Shipment and Storage.

(Copies of Military Specifications and Standards required by contractors in connection with specific procurement functions should be obtained from the procuring activity or as directed by the contracting officer)

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T11:14Z
Check the source to verify that this is the current version before use.

Laws and Regulations:

Federal Insecticide, Fungicide and Rodenticide Act of 1947.

(Copies may be obtained upon application, accompanied by money order, coupon, or cash to the Superintendent of Documents, U. S. Government Printing Office, Washington, D. C. 20402.)

2.2 Other publications. The following documents form a part of this specification to the extent specified herein. Unless otherwise indicated, the issue in effect on date of invitation for bids or request for proposal shall apply:

Nongovernmental:

Association of Official Analytical Chemists (AOAC):

Official Methods of Analysis, 10th Edition, 1965.

(Application for copies should be addressed to the Association of the Official Analytical Chemists, P. O. Box 540, Benjamin Franklin Station, Washington, D. C. 20036.)

Uniform Classification Committee:

Uniform Motor Freight Classification.

(Application for copies should be addressed to the Uniform Classification Committee, 202 Union Station, Chicago, Illinois 60606.)

National Classification Board:

National Motor Freight Classification.

(Application for copies should be addressed to the American Trucking Association, Inc., Attn: Tariff Order Section, 1616 P Street, N.W. Washington, D. C. 20036.)

3. REQUIREMENTS

3.1 Material. The rodenticide shall consist of a chemical anticoagulant which may be diphacinone, fumarin, Pival, or warfarin as specified (see 6.2 (e)), suitably compounded with petrolatum, sugar, rolled oats and a dye, the admixture conforming to the compositional requirements given in table I.

TABLE I. Composition

Ingredient	Percent	
	Minimum	Maximum
Anticoagulant:		
As specified (see 6.2 (e)).		
1. Diphacinone	0.005	0.0055
2. Fumarin	.025	.028
3. Pival	.025	.028
4. Warfarin	.025	.028
Liquid petrolatum	4.8	5.2
Sugar, refined	4.8	5.2
Dye	0.05	---
Breakfast rolled oats	Remainder	

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T11:14Z
Check the source to verify that this is the current version before use.

3.1.1 Diphacinone. Diphacinone shall be the chemical 2 -(diphenylacetyl)1, 3-indandione, and the percentage shown in the ingredient statement shall represent pure diphacinone. This diphacinone shall be derived from a technical product which shall be a yellow crystalline powder, melting at 145° to 147°C., and containing not less than 96 percent diphacinone. It shall be free of objectionable odor or taste which might limit its suitability for the proposed end use (see 3.2 and 4.5.1).

3.1.2 Fumarin Fumarin shall be the chemical 3 - (alpha-acetonylfurfuryl)-4-hydroxycoumarin, and the percentage shown in the ingredient statement shall represent pure fumarin. This fumarin shall be derived from a technical product which shall be a fine white or creamy white powder, melting at 120° to 123°C., and containing not less than 96 percent fumarin. It shall be free of objectionable odor or taste which might limit its suitability for the proposed end use (see 3.2 and 4.5.2).

3.1.3 Pival. Pival shall be the chemical 2-(pivalyl) 1, 3-indandione, and the percentage shown in the ingredient statement shall represent pure Pival. This Pival shall be derived from a technical product which shall be a fine yellow or brownish-yellow powder, melting at 107° to 110°C., and containing not less than 96 percent Pival. It shall be free of objectionable odor or taste which might limit its suitability for the proposed end use (see 3.2 and 4.5.3).

3.1.4 Warfarin. Warfarin shall be the chemical 3 - (alpha-acetonylbenzyl)-4-hydroxycoumarin, and the percentage shown in the ingredient statement shall represent pure warfarin. This warfarin shall be derived from a technical product which shall be a fine white or creamy white powder melting at 159° to 163°C., and containing not less than 96 percent warfarin.

3.1.5 Liquid petrolatum. The liquid petrolatum shall be medicinal grade mineral oil, with a specific gravity of not less than 0.83 nor more than 0.86 (see 4.5.8).

3.1.6 Sugar. The sugar shall conform to the requirements of type I, as specified in JJJ-S-791 (see 4.5.8).

3.1.7 Dye. The dye shall be an oil-soluble, brilliant red dye, approved by the Fish and Wildlife Service. It shall be equivalent to Flamming Red Dye No. G305, manufactured by the Ox Color Works, New York, New York.

3.1.8 Rolled oats. The rolled oats shall conform to the requirements of type I, Regular, Rolled Oats, as specified in N-O-41.

3.2 Certificate of compliance. For each delivery of finished rodenticide, the manufacturer shall submit a certified copy of the production record, showing:

(a) Certified copies of analyses of chemical anticoagulant, liquid petrolatum, sugar and rolled oats used in formulation.
(b) Exact weights of all ingredients used.
(c) Time spent in mixing or blending.
(d) Weight of finished product.

3.3 Particle size. Not more than 30 percent of the finished rodenticide shall pass through a No. 20 U. S. Standard woven-wire cloth sieve, when tested as described in 4.5.5.

3.4 Moisture. The finished rodenticide shall not lose more than 12 percent of its weight when tested as described in 4.5.6.

3.5 Performance. The toxicity and performance of the finished rodenticide shall be such that not more than 10 percent of the test animals survive and the test ration comprises a minimum of 33 percent of the test animals total diet when the product is tested as described in 4.5.9.

3.6 Federal Insecticide, Fungicide and Rodenticide Act. The finished rodenticide shall comply with the applicable provisions of the Federal Insecticide, Fungicide and Rodenticide Act (see 6.4).

3.7 A mimeographed instruction sheet, containing the information shown below, shall be placed

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T11:14Z
Check the source to verify that this is the current version before use.

in an envelope marked "Important Instructions". These instruction sheets shall be placed in containers in accordance with 5.1.1.2 and 5.1.2.2.

INSTRUCTIONS FOR USING RODENTICIDE, BAIT, ANTICOAGULANT

1. Do not breathe the dust or swallow any of the poison. Do not smoke or eat while handling the poison. Keep personnel contact with the rodenticide at a minimum.

2. Keep all equipment labeled plainly, wash it thoroughly after use, and do not employ it for other purposes.

3. Wash hands thoroughly after handling the poisoned bait.

METHOD FOR USE IN CONTROL OF RATS AND MICE

Rodenticides, containing (insert here:diphacinone, fumarin, Pival, or warfarin) reduce the blood coagulation rate and death is ensured in most cases from internal hemorrhages.

This rodenticide will not give effective control when applied in a single dose, but must be ingested several times (usually five or more) on successive days.

The control of rats, including roof rats, and house mice, has been obtained through the use of (insert here:diphacinone, fumarin, Pival, or warfarin).

BAIT EXPOSURE

A. RATS

Exposure of baits should be made by establishing an adequate number of protected bait stations in locations where rats feed, water or travel. This can be accomplished by placing open trays of bait in rooms or areas about premises where domestic animals and small children can be excluded or through the use of properly constructed bait boxes. In many instances baits can be protected by improvised means with materials available on the premises. This is specially true where it is possible to employ scrap lumber for this purpose.

Feeding stations should be inspected at frequent intervals for the purpose of replenishing the bait supply where necessary. The frequency of inspections and size of bait placements will, of course, depend upon the degree of infestation encountered and in some instances daily servicing of the bait stations may be required during the first four to five days of a treatment. Bait consumption and rat activity usually drop markedly about 5 days after a rat population begins to feed on baits. Control is usually achieved in 10 to 14 days after feeding begins. Baits should be kept dry during use to maintain maximum acceptability. Contaminated or spoiled baits should be replaced promptly.

It is often desirable to maintain properly protected bait stations with semi-permanent baits after control has been attained on premises which are particularly vulnerable to re-invasion.

Four to eight ounces of bait are suggested as the amount to be placed in each feeding station. Subsequent inspection will determine the correct amount to be added to each station.

B. MICE

In general, the same exposure technique should be used in employing baits for the control of mice as outlined for rat control; the principal difference, however, being that a larger number of placements should be made in areas in which the mice are known to feed. Those feeding stations may in some cases consist of only tablespoonful quantities placed in shallow containers 8 to 12 feet apart.

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T11:14Z
Check the source to verify that this is the current version before use.

Inspection should be made frequently enough throughout a treatment to maintain a continuous supply of bait available at each placement. Baiting should continue for at least 15 days or until feeding stops.

PRECAUTIONS

(Insert here: diphacinone, fumarin, Pival, or warfarin) probably will kill any warm blooded animal if consumed in the proper dosage over an adequate period of time. However, the concentration used in rodent control requires that on the basis of a single dose ingestion, relatively large quantities of the bait be consumed before domestic animals are seriously endangered. On the other hand, if daily feeding on diphacinone, fumarin, pival, or warfarin by domestic animals is allowed, much smaller amounts may be fatal. Dead rats and mice should be picked up and destroyed, so that cats and dogs do not have access to them.

ANTIDOTE

(Insert here diphacinone, fumarin, Pival, or warfarin) is an anticoagulant chemical, which if taken by humans or domestic animals, may increase the clotting time of the blood and cause hemorrhage. Such hemorrhages may be either internal or external and often hematoma are visible under the skin of animals affected with these substances.

In the case of ingestion by humans, vomiting should be induced at once and a physician called. Treatment by the physician should include transfusion with whole blood and intravenous and oral administration of vitamin K preparations as in the case of hemorrhage caused by dicumarol.

3.8 Workmanship. The method employed in making the rodenticide shall be in accordance with best commercial practice. The product shall be clean, uniformly blended, and free of defects.

4. QUALITY ASSURANCE PROVISIONS

4.1 Responsibility for inspection. Unless otherwise specified in the contract or purchase order, the supplier is responsible for the performance of all inspection requirements as specified herein. Except as otherwise specified, the supplier may utilize his own facilities or any commercial laboratory acceptable to the Government. The Government reserves the right to perform any of the inspections set forth in the specification where such inspections are deemed necessary to assure that supplies and services conform to prescribed requirements.

4.2 Sampling for lot acceptance.

4.2.1 Inspection lot. For purposes of sampling, a lot shall consist of all rodenticide manufactured as one batch, and offered for delivery at one time.

4.2.2 Sampling for inspection of filled containers. A random sample of filled containers shall be taken from each lot offered for inspection in accordance with MIL-STD-105 at inspection level I, and acceptable quality level (AQL) = 2.5 percent defective to verify compliance with all stipulation of this specification regarding fill, closure, marking and other requirements not involving tests.

4.2.3 Sampling for tests. From each inspection lot offered for testing, enough containers shall be taken at random to form two separate five-pound specimens. The containers shall be sealed, marked, and forwarded to the testing laboratory designated by the bureau or agency concerned. A copy of the certificate of compliance (see 3.2), shall accompany the specimens.

4.2.3.1 Components. The following samples of components shall be taken and forwarded to the Government testing laboratory:

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T11:14Z
Check the source to verify that this is the current version before use.

Diphacinone, fumarin, Pival, or warfarin (as applicable)	4 ounces
Dye	1 ounce
Liquid petrolatum	4 ounces
Sugar	1 pound
Rolled oats	2 pounds

4.3 Inspection.

4.3.1 Inspection of filled containers. Each sample filled container selected in accordance with 4.2.2, shall be examined for defects of the container and the closure, for evidence of leakage, and for unsatisfactory markings; each sample filled container shall also be weighed to determine the amount of the contents. Any container in the sample, having one or more defects, or under required fill, shall be rejected and, if the number of defective containers in any sample exceeds the acceptance number for the appropriate sampling plan of MIL-STD-105, the lot represented by the sample shall be rejected. Rejected lots may be resubmitted for acceptance tests provided that the contractor has removed or repaired all nonconforming containers.

4.4 Lot acceptance tests. The sample specimens selected in accordance with 4.2.3, shall be subjected separately to the tests specified in 4.5. If either specimen fails one or more of these tests, the lot shall be rejected. Rejected lots may be resubmitted for acceptance tests provided the contractor has removed or reworked all nonconforming products.

4.5 Test methods.

4.5.1 Diphacinone determination. Accurately weigh to within 0.1 mg. approximately 100 mg. of diphacinone into a 100 ml. volumetric flask and make to volume with 3-A alcohol (100 volumes of reagent ethanol and 5 volumes of reagent methanol). After solution is achieved, dilute a 10.0 ml. aliquot and make to 1000 mls. in a volumetric flask with 3-A alcohol. Pipette a sufficient amount (approximately 3 ml) of the diluted aliquot into a 1 cm. Quartz cuvette, and determine its maximum absorbance (A_s) found between 286 and 288 mu. in a Beckman spectrophotometer model D. U. or its equivalent, using 3-A alcohol as a blank. The percentage of diphacinone in the bait is calculated by the following formula:

$$\text{Percent diphacinone} = \frac{\text{Max. } A_s \text{ found X } 0.5862}{\text{Wt. of anticoagulant in mgs. corrected for the \% purity*}}$$

This calculation is based on the percent composition of the bait conforming to 3.1 table I calculated from 3.2 (a), (b), and (d).

4.5.2 Fumarin determination. Accurately weigh to within 0.1 mg. approximately 200 mg. of fumarin into a 100 ml. volumetric flask and make to volume with 3-A alcohol (100 volumes of reagent ethanol and 5 volumes of reagent methanol). After solution is achieved dilute a 10.0 ml. aliquot and make to 1000 mls. in a volumetric flask with 3-A alcohol. Pipette a sufficient amount (approximately 3 ml.) of the diluted aliquot into a 1 cm. quartz cuvette, and determine its maximum absorbance (A_s) found between 304 and 306 mu. in a Beckman spectrophotometer, model D. U. or its equivalent, using 3-A alcohol as a blank. The percentage of fumarin in the bait is calculated by the following formula:

$$\text{Percent fumarin} = \frac{\text{Max } A_s \text{ found X } 7.634}{\text{Wt. of anticoagulant in mgs. corrected for the \% purity*}}$$

This calculation is based on the percent composition of the bait conforming 3.1 table I calculated from 3.2 (a), (b), and (d).

4.5.3 Pival determination. Accurately weigh to within 0.1 mg. approximately 50 mg. of Pival from 4.2.3.1 into a 100 ml. volumetric flask and make to volume with 3-A alcohol (100 volumes of reagent ethanol and 5 volumes of reagent methanol). After solution is achieved dilute a 10.0 ml. aliquot to 1000 mls., in a volumetric flask with 3-A alcohol. Pipette a sufficient amount (approximately 3 ml.) of the diluted aliquot into a 1 cm. Quartz cuvette, and determine its maximum absorbance (A_s) found between 283 and 286 mu. in a Beckman spectrophotometer, model D. U. or its equivalent, using 3-A alcohol as a blank. The percentage of Pival in the bait is calculated by the following formula:

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T11:14Z
Check the source to verify that this is the current version before use.

$$\text{Percent Pival} = \frac{\text{Max. } A_s \text{ found X } 1.969}{\text{Wt. of anticoagulant in mgs. corrected for the \% purity*}}$$

This calculation is based on the percent composition of the bait conforming to 3.1 table I calculated from 3.2 (a), (b), and (d).

4.5.4 Warfarin determination. Accurately weigh to within 0.1 mg. approximately 200 mg. of warfarin from 4.2.3.1 into a 100 ml. volumetric flask and make to volume with 3-A alcohol (100 volumes of reagent ethanol and 5 volumes of reagent methanol). After solution is achieved dilute a 10.0 ml. aliquot and make to 1000 mls. in a volumetric flask with 3-A alcohol. Pipette a sufficient amount (approximately 3 ml.) of the diluted aliquot into a 1 cm. quartz cuvette, and determine its maximum absorbance (A_s) found between 305 and 307 mu. in a Beckman spectrophotometer, model D. U. or its equivalent, using 3-A alcohol as a blank. The percentage of warfarin in the bait is calculated by the following formula:

$$\text{Percent Warfarin} = \frac{\text{Max. } A_s \text{ found X } 7.289}{\text{Wt. of anticoagulant in mgs. corrected for the \% purity*}}$$

This calculation is based on the percent composition of the bait conforming to 3.1 table I calculated from 3.2 (a), (b), and (d).

* Percent purity: See paragraph 3.2 (a). Use % purity of chemical anticoagulant stated in the certified copy of analysis.

4.5.5 Sieve test. Transfer 100 grams of the finished rodenticide to a No. 20 U.S. Standard sieve conforming to RR-S-366 and screen in a Ro-tap or equivalent machine for two minutes. Weigh the material which passes through the sieve and calculate the percentage.

4.5.6 Moisture. Accurately weigh approximately 2 grams of the finished rodenticide into a tared moisture dish (Ni, Pt, or Al) and dry for 10 hours under atmospheric pressure at a temperature of 100°C. (± 20°C.); cool in desiccator and weigh. Dry again for one hour, or until change in weight is not more than 2 mg. Report loss in weight as moisture.

4.5.7 Liquid petrolatum. Accurately weigh approximately 100 grams of the finished rodenticide, ground to pass a 20 mesh screen, prepare ether extract and determine percentage of mineral oil according to procedures prescribed in Methods of Analysis of the Association of Official Analytical Chemists.

4.5.8 Sugar. Accurately weigh sample of finished rodenticide powder and determine sugar content according to procedures described in JJJ-S-791.

4.5.9 Performance. The suitability of the rodenticide concentrate and finished bait shall be determined according to the procedures described in 4.5.9.1.1 and 4.5.9.2.2.

4.5.9.1 Materials.

4.5.9.1.1 Bio-assay animals. The bio-assay animals shall consist of mature, healthy, male and female albino rats of Wistar strain weighing between 150 and 250 grams.

4.5.9.1.2 Control diet. The control diet shall consist of pelleted laboratory chow, or dog biscuits, equivalent to that commonly employed for feeding of laboratory rats.

4.5.9.2 Procedure.

4.5.9.2.1 Preliminary treatment of bio-assay animals. Thirty mature, healthy albino rats (15 males and 15 females) are to be placed in individual feeding cages.

4.5.9.2.2 Bio-assay procedure. The test ration, control diet and water are to be supplied ad libitum for a period of 15 consecutive days. Each cage provided with separate water devices and dual feed cups. The correct ration and control diet should each be offered in excess of the daily food requirement (over 30 grams for each feed cup). The gross weight of each cup and its contents to be determined daily and returned to starting weight. The position of the feed cups in each cage, to be reversed every 24 hours to counter any feeding position habit of the rat. The test ration is to be removed at the end of the fifteen day period and the remaining bio-assay animals given ad libitum access to the control diet for an additional observation period of 5 days. The temperature of the bio-assay room shall be kept between 65° and 80°F. throughout the test period.

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T11:14Z
Check the source to verify that this is the current version before use.

Other bio-assay room conditions such as light, humidity, time of feeding and observation periods shall be standardized. If not more than 10 percent of the test animals survive and the test ration comprises a minimum of 33 percent of the test animals total diet; the rodenticide concentrate and the finished rodenticide shall be deemed acceptable.

4.6 Examination of preparation of delivery requirements. An examination shall be made to determine that packaging, packing, labeling and marking comply with the requirements of section 5. Defects shall be as specified in table II. Sampling shall be in accordance with MIL-STD-105. The sample unit shall be one container fully prepared for delivery. The lot shall be the no. of containers offered for inspection at one time. The inspection level shall be S-2 with an AQL of 4.0 defects, expressed in terms of defects per hundred units.

TABLE II. Examination of preparation for delivery defects

Examine	Defects
Marking and labeling	Omitted; incorrect, illegible; improper size, location, sequence or method of application.
Materials	Component missing or damaged.
Workmanship	Bulging or distortion of containers. Cushioning inadequate, improper or missing.
Contents	More or less than required.
Containers	Not as specified.

5. PREPARATION FOR DELIVERY

5.1 Pakaging shall be level A, B, or C, as specified (see 6.2).

5.1.1 Level A.

5.1.1.1 Unit packaging. The rodenticide shall be packaged in quantities or 2-ounce, 1-pound, 1-½ pounds, 3-pounds, 5-pounds, or 6-pounds, as specified (see 6.2), in round hermetically sealed open-top cans conforming to PPP-C-96, type I, plan B coating. The product shall be compressed to a density of at least 1-ounce for each 2.6 cubic inches based on the net inside packing space of the can.

5.1.1.2 Intermediate packaging. Intermediate packaging of 2-ounce cans shall be as specified in the appendix to PPP-C-96 for level A packaging. Forty-eight instruction sheets (see 3.7) shall be placed in each intermediate package.

5.1.2 Level B.

5.1.2.1 Unit packaging. The rodenticide shall be packaged in quantities of 2-ounces, 1-pound, 1-½pounds, 3-pounds, 5-pounds, or 6-pounds, as specified (see 6.2), in round, hermetically sealed open-top cans conforming to PPP-C-96, type I, plan A coating. The product shall be compressed to a density of at least 1-ounce for each 2.6 cubic inches based on the net inside packing space of the can.

5.1.2.2 Intermediate packaging. Intermediate packaging of 2-ounce cans shall be as specified in the appendix to PPP-C-96 for level A packaging, except that the intermediate package shall conform to PPP-B-636, class domestic. Forty-eight instruction sheets (see 3.7) shall be placed in each intermediate package.

5.1.3 Level C. Rodenticide in the quantity specified (see 6.2) shall be packaged to afford protection against deterioration and physical damage during shipment from the supply source to the first receiving activity for immediate use.

5.2 Packing. Packing shall be level A, B, or C, as specified (see 6.2).

5.2.1 Level A. Packing shall be in accordance with the appendix to PPP-C-96 for level A packing. Twenty-four 1-½ pound cans shall be packed in the shipping container. The 6-pound cans shall be packed in the shipping container in the same arrangement designed for the 5-pound cans.

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T11:14Z
Check the source to verify that this is the current version before use.

Four intermediate packages of 2-ounce cans shall be packed in the shipping container.

5.2.2 Level B. Packing shall be in accordance with the appendix to PPP-C-96, for level B packing. Twenty-four 1-½ pound cans shall be packed in the shipping container. The 6-pound cans shall be packed in the shipping container in the arrangement designated for the 5-pound cans. Four intermediate packages of 2-ounce cans shall be packed in the shipping container.

5.2.3 Level C. Rodenticide, packaged as specified (see 6.2), shall be packed in containers acceptable to the common carrier, and which will insure safe delivery at destination in a satisfactory condition at the lowest applicable rate. Containers, packing or method of shipment, shall comply with Uniform Freight or National Motor Freight Classification Rules and Regulations, or other carrier rules as applicable to the mode of transportation.

5.3 Marking.

5.3.1 Civil agencies. In addition to the labeling and the marking specified in 5.3.3 and any special marking required by the contract or order, marking for shipment shall be in accordance with Fed. Std. No. 123.

5.3.2 Military agencies. In addition to the labeling and the marking specified in 5.3.3 and any special marking required by the contract or order, marking for shipment shall be in accordance with MIL-STD-129.

5.3.3 Labeling. The rodenticide shall be registered and labeled in compliance with the Federal Insecticide, Fungicide and Rodenticide Act.

6. NOTES

6.1 Intended use. Rodenticide is intended for use in the control of commensal rats and mice.

6.2 Ordering data. Purchasers should select the preferred options permitted herein and include the following information in procurement documents:

(a) Title, number, and date of this specification.
(b) Whether packaging and packing should be level A, B, or C (see sec. 5).
(c) Quantity to the unit package (see 5.1.1.1 and 5.1.2.1).
(d) Quantity required (see 5.1.3).
(e) Whether diphacinone, fumarin, pival, or warfarin is required (see 3.1).

6.3 Approved laboratory. The Wildlife Research Laboratory, U. S. Fish and Wildlife Service Laboratory, Agricultural Experiment Station, Purdue University, Lafayette, Indiana, is an approved laboratory for testing the rodenticide to determine compliance with the requirements of this specification.

6.4 The Federal Insecticide, Fungicide, and Rodenticide Act of 1947 is under the jurisdiction of the Pesticides Regulation Division, Agricultural Research Service, U. S. Department of Agriculture, Washington, D. C. 20250.

CUSTODIANS:

Army - GL
Navy - SH

User activities:

Army - ME
Navy - MS, MC

Review activities:

Army - GL, MD, MU, CE
Navy - SH, YD
Air Force - 68

Preparing activity:

GSA-FSS

CIVIL AGENCIES INTEREST:

AGR
HEW
DC GOV
VA

Orders for this publication are to be placed with [illegible] acting as an agent for the Superintendent of Documents. See section 2 of this specification to obtain extra copies and other documents referenced herein.

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T11:14Z
Check the source to verify that this is the current version before use.

GENERAL SERVICES ADMINISTRATION - FEDERAL SUPPLY SERVICE

SPECIFICATION COMMENT SHEET

BUDGET BUREAU NO. 29-R0175

INSTRUCTIONS

This form provides a way for users of this specification to inform the originator of problems encountered in its use. It is not to be used to request changes to accommodate proprietary features. All comments will be considered and appreciated, but please do not expect a reply. To comment: detach, complete, and mail to: General Services Administration, FSS (FMSC), Wash., DC 20406.

NOTE: Comments on this form do not constitute or imply authorization to waive any part of the document or serve to amend contractual requirements.

1. SPECIFICATION

O-R-500C Rodenticide, Bait, Anticoagulant

2. CONTRACT NO. *(If any)*	3. QUANTITY ON CONTRACT *(Optional)*	4. DOLLAR VALUE *(Optional)*

5. GENERAL NATURE OF PROBLEM *(e.g., inspection difficulties, manufacturers unable to meet tolerances, containers collapse under normal warehousing conditions, etc.)*

6. SPECIFIC REQUIREMENTS AFFECTED *(Include paragraph number and lines of wording)*

7. SPECIFIC PROBLEMS *(e.g. tests in 4.2.2 will not assure that the battery will last required time; temperature ranges in table 2 do not conform to commercially available items.)*

8. RECOMMENDATIONS

9. NAME OF MANUFACTURER, ASSOCIATION, GOVT., AGENCY, ETC.	10. ADDRESS *(Number, Street, City, State and Zip Code)*

11. NAME AND TITLE OF SUBMITTER	12. DATE

GSA DC 69-9761

FORM GSA FEB 69 2200

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T11:14Z
Check the source to verify that this is the current version before use.