MILITARY SPECIFICATION

# PANCURONIUM BROMIDE INJECTION

This specification is approved for use by all Departments and Agencies of the Department of Defense

## 1. SCOPE

1.1 Scope. This specification covers Pancuronium Bromide Injection.

1.2 Classification. The injection covered by this specification shall be of the following sizes and strengths (See 6.2):

Strength 1 - 2 mg per ml
Strength 2 - 1 mg per ml
Size 1 - 2 ml ampul
Size 2 - 5 ml ampul
Size 3 -10 ml vial

## 2. APPLICABLE DOCUMENTS

2.1 Issues of documents. The following documents, of the issue in effect on date of invitation for bids or request for proposals, form a part of this specification to the extent specified herein.

SPECIFICATIONS

FEDERAL

PPP-C-186 - Containers, Packaging and Packing for Drugs, Chemicals and Pharmaceuticals. (See 6.3)

STANDARDS

FEDERAL

FED-STD-142 - Parenteral Preparations.

Beneficial comments (recommendations, additions, deletions and any pertinent data which may be of use in improving this document should be addressed to: Hq, Defense Personnel Support Center, ATTN: Directorate of Medical Materiel, DPSC-ATT, 2800 South 20th Street, Philadelphia, PA 19101, by using the self-addressed Standardization Document Improvement Proposal (DD Form 1426) appearing at the end of this document, or by letter.



Source: https://assist.dla.mil -- Downloaded: 202640-21T10:39Z
Check the source to verify that this is the current version before use.

MILITARY

MIL-STD-105 - Sampling Procedures and Tables for Inspection by Attributes.
MIL-STD-129 - Marking for Shipment and Storage.

(Copies of specifications and standards required by contractors in connection with specific procurement functions should be obtained from the procuring activity or as directed by the contracting officer.)

2.2 Other publications. The following documents form a part of this specification to the extent specified herein. Unless otherwise indicated, the issue in effect on date of invitation for bids or request for proposals shall apply.

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE,
FOOD AND DRUG ADMINISTRATION

Federal Food, Drug, and Cosmetic Act and Regulations Promulgated Thereunder.

(Application for copies should be addressed to the Food and Drug Administration, Washington, DC 20204.)

U. S. PHARMACOPEIAL CONVENTION, INC.

The United States Pharmacopeia.

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

National Formulary.

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

3. REQUIREMENTS

3.1 Material. Shall be Pancuronium Bromide Injection and shall be in accordance with all applicable requirements of FED-STD-142, and as specified herein.



Source: https://assist.dla.mil -- Downloaded: 202640-21T10:39Z
Check the source to verify that this is the currentversion before use.

S2. Classification. Shall be type I, class 1.

Shall be suitable for intravenous use.

S5.2.2 The following is added to this paragraph:

For strength 1, shall be a sterile, isotonic solution containing 2 mg of Pancuronium Bromide per ml within the designated assay limits for the injection.

For strength 2, shall be sterile, isotonic solution containing 1 mg of Pancuronium Bromide per ml within the designated assay limits for the injection.

Shall contain not less than 90.0 percent and not more than 110.0 percent of the required amount of Pancuronium Bromide when assayed as specified in 4.5.1.1.

The pancuronium bromide in the injection shall comply with the following identification test. The main spot of the pancuronium bromide shall conform to the main spot of the standard in Rf, size and color when tested as directed in the procedure for Related Foreign Steroids (see 4.5.1.2)

The injection shall have a total amount of related foreign steroids (decomposition products) of not more than 6 percent during the potency period when tested as specified in 4.5.1.2.

The injection shall have a pH within the range of 3.0 to 5.0 at 25°C when determined potentiometrically, using the USP method.

3.2 Ingredients.

3.2.1 Pancuronium Bromide. The Pancuronium Bromide used in the manufacture of the finished injection shall comply with the following requirements.

3.2.1.1 Melting Point. Shall melt with decomposition at about 215°C using USP method for class 1a compounds.

3.2.1.2 Residue on Ignition. Shall have not more than 0.1 percent residue using USP method.



Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:39Z
Check the source to verify that this is the current version before use.

3.2.1.3 Water Content. Shall have not more than 8.0 percent when tested using USP Karl Fischer titrimetric method.

3.2.1.4 Loss on Drying. Shall be not more than 1.0 percent greater than the water content found when determined as specified in 4.5.2.1.

3.2.1.5 Specific Rotation. Shall have a specific rotation of not less than + 38° and not more than + 44° on the anhydrous basis when determined as specified in 4.5.2.2.

3.2.1.6 Ultraviolet absorption. Shall have no absorption peaks between 220 mu-360 mu when tested as specified in 4.5.2.3.

3.2.1.7 Color of Solution. Shall have an absorbance of not greater than 0.10 between 400 mu-700 mu when tested as specified in 4.5.2.4.

3.2.1.8 Identification. Shall comply with the following tests.

(a) Thin Layer Chromatograph - the main spot of the sample conforms to the main spot of the standard in Rf, size and color when tested as directed in Related Foreign Steroid (see 4.5.2.5).

(b) Infrared absorption - the infrared absorption spectrum of a K Br dispersion of the sample previously dried for 2 hours at 60°C over $P_2O_5$ under vacuum conforms to the Pancuronium Bromide Standard spectrum similarly determined.

(c) Bromide ion-positive test for Bromide ion when tested in USP Identification Tests - General.

3.2.1.9 Related Foreign Steroids.(Decomposition Products). Shall have not more than 2 percent total foreign steroids when tested as specified in 4.5.2.5.

3.2.1.10 Assay. Shall assay to contain not less than 98.0 percent and not more than 102.0 percent of Pancuronium Bromide, on the anhydrous basis when determined as specified in 4.5.2.6.

3.3 Other ingredients. All other ingredients used in the manufacture of the injection shall comply with S5.1 of FED-STD-142.

3.4 Expiration dating period. The expiration dating period shall be not less than 24 months.



Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:39Z
Check the source to verify that this is the current version before use.

## 4. QUALITY ASSURANCE PROVISIONS

4.1 Supplier responsibility for inspection. Unless otherwise specified in the contract or purchase order, the supplier is responsible for the performance of all inspection requirements as specified herein. Except as otherwise specified in the contract or order, the supplier may use his own or any other facilities suitable for the performance of the inspection requirements specified herein, unless disapproved by the Government. The Government reserves the right to perform any of the inspections set forth in the specification where such inspections are deemed necessary to assure supplies and services conform to prescribed requirements.

4.1.1 Records of examinations and tests performed by or for the contractor shall be maintained by the contractor and made available to the Government, upon the Government's request, at any time, or from time to time, during the performance of the contract and, (i) as to any expiration dated item, for the expiration dating period specified by the contractor for such item, or such longer period as may be required by regulation of any federal agency; or (ii) as to non-dated items, for not less than 3 years after delivery of the item to the Government.

4.1.2 No company supplying any ingredient(s) to the contractor will be considered an acceptable facility for the performance of any inspection requirements specified herein.

4.2 Sampling. The sampling as shown in FED-STD-142 shall be applicable.

4.3 Packaging inspection. The inspection of the intermediate package, packing and marking for shipment and storage shall be in accordance with quality assurance provisions of the applicable container specification and the marking requirements of MIL-STD-129.

4.4 Labeling inspection. Sampling shall be in accordance with MIL-STD-105 and the unit of product shall be the unit of issue. The inspection lot shall be the number of units of issue of each batch, lot, control number, etc., offered for inspection at one time. The inspection level for labeling shall be S-2 and the acceptance number shall be zero.



Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:39Z
Check the source to verify that this is the current version before use.

4.5 Tests.

4.5.1 Finished product.

4.5.1.1 Assay for Pancuronium Bromide.

Reagents.

Weigh 100 mg finely powdered bromophenol blue into a 100 ml volumetric flask, add 50 ml of purified water and dissolve by warming, at about 50°C. Cool to room temperature and dilute to volume with purified water. Prepare this reagent fresh daily.

Standard Preparation:

Weigh accurately about 25 mg of Pancuronium Bromide Standard into a 50 ml volumetric flask. Dissolve in and dilute to volume with purified water and mix. Pipet 10 ml of this solution into a 50 ml volumetric flask and dilute to volume with purified water and mix (diluted standard).

NOTE: Correct the standard weight for as is assay.

Sample Preparation:

Dilute 5.0 ml of sample (Strength 1) or 10.0 ml of sample (Strength 2) to 100 ml with purified water in a volumetric flask.

Procedure:

Pipet 5 ml of the sample preparation and diluted standard preparation into separate 125 ml separatory funnels containing 20 ml of purified water and 40 ml methylene chloride. Add 10 ml of a 1% (W/V) solution of Anhydrous Sodium Carbonate in purified water and 3 ml of the Bromophenol blue reagent. Shake each funnel for 2 minutes and allow the layers to separate. Draw off the methylene chloride layer into 100 ml volumetric flasks, being certain not to draw off any portion of the aqueous layer. Repeat the extraction 4 times, shaking for 1 minute, using 10 ml of methylene chloride for each. To the combined methylene chloride extracts, cautiously add exactly 15 ml of absolute alcohol, mix and dilute to volume with methylene chloride. Measure the absorbance of the standard and sample, respectively, in a suitable spectrophotometer using methylene chloride as a blank at the point of maximum absorbance of about 600 mu.



Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:39Z
Check the source to verify that this is the current version before use.

Calculation:

$$\frac{au}{as} \quad \frac{mg \text{ standard}}{50} \text{ [1/]} \times \frac{10}{50} \quad \frac{100}{X} = \frac{mg \text{ Pancuronium Bromide}}{ml}$$

where au = absorbance of sample
as = absorbance of standard
X = 5 for strength 1 or 10 for strength 2

4.5.1.2 Identification and Related Foreign Steroids (Decomposition Products) of the Injection.

Plate Preparation. (For 5 plates, 20 x 20 cm). Weigh into a mortar 25 grams of Silica Gel G 2/, add 50 ml purified water and mix completely. Coat the plates with an applicator, allow them to air dry 30 minutes and dry in oven at 110°C for 1 hour. Store in a desiccator over silica gel.

Solvent system. Measure into a separatory funnel, 80 ml pyridine, 120 ml n-butanol, 24 ml glacial acetic acid and mix. Add about 50 ml of 20% W/V Ammonium Chloride in purified water and shake gently. Draw off the bottom layer and discard. Place enough of the mixture to saturate a paper lined chromatographic chamber and allow to equilibrate for 30 minutes before use.

Sample Preparation. Use sample as is.

Standard Preparations.

a) Prepare accurately a solution in purified water to contain 0.2 mg/ml (strength 2) or 0.4 mg/ml (strength 1) each of (2 Beta, 16 Beta - Dipiperidino-5 alpha-Androstane-3 alpha, 17 Beta-Diol 3 alpha-acetate Dimethobromide) plus (2 Beta, 16 Beta-Dipiperdino-5 alpha Androstane-3 alpha, 17 Beta-Diol, 17 Beta-Acetate Dimethobromide). Call this solution #a.

1/ correct for assay.
2/ Note: Use Silica Gel G, Woelm TLC, 13% Calcium Sulfate, as supplied by Waters Associates, Inc., Framingham, MA, or equivalent.



Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:39Z
Check the source to verify that this is the current version before use.

b) Into three separate 10 ml volumetric flasks, weigh accurately 10 mg of Pancuronium Bromide standard calculated on the anhydrous basis and dissolve in about 5 ml purified water. Pipet into one flask 0.5 ml of solution #a, and 1.5 ml of solution #a into the second flask and 3.0 ml of solution #a into the third flask. Dilute each flask to volume with purified water and mix. These correspond to 2% total, 6% total and 12% total of related foreign steroids respectively.

Procedure. Apply to the prepared plates, 5 microliters each of the sample and standard preparations about 2 cm apart and 2 cm from the bottom of the plate. Place the plate in the prepared chamber and allow the solvent to move about 17 cm, remove the plate from the chamber and allow it to dry in a current of air in a fume hood for at least one hour. Spray the plate evenly and completely with a fine mist of sulfuric acid in alcohol (20% V/V) and heat at 110°C for one hour. Examine the plate under ultraviolet light. The main spot of the sample conforms to the main spot of the standard in Rf, size and color (Identity Test). Any secondary spots appearing in the sample are not greater in size or intensity than the secondary spots of the standard. The total amount of related foreign steroids (decomposition) may not exceed 6% combined.

4.5.2 Tests for Active Ingredient (Pancuronium Bromide).

4.5.2.1 Loss on Drying. Dry about 200 mg of sample, accurately weighed, at 60°C under vacuum over $P_2O_5$ for 2 hours and reweigh.

4.5.2.2 Specific Rotation. Proceed according to USP except weigh 100 mg sample in 10 ml solvent prepared from 1 ml alcohol in 100 ml chloroform. Correct the sample weight for water content or loss on drying if greater than water content.

4.5.2.3 Ultraviolet Absorption. Dissolve 100 mg of sample (anhydrous basis) in 10.0 ml of purified water and measure the absorption spectrum between 220 mu and 360 mu in a 1 cm cell relative to purified water using a suitable recording spectrophotometer.



Source: https://assist.dla.mil -- Downloaded: 202640-21T10:39Z
Check the source to verify that this is the current version before use.

4.5.2.4 Color of Solution. Prepare a 5% W/V solution of sample (anhydrous basis) in purified water. Measure the absorbance of the solution in a 1 cm cell relative to purified water using a suitable recording spectrophotometer between 400 mu and 700 mu.

4.5.2.5 Identification (a) and Related Foreign Steroids (Decomposition Products) of the Pancuronium Bromide Powder.

Plate Preparation. (For 5 plates, 20 x 20 cm) Weigh into a mortar 25 grams of Silica Gel G, 1/ add 50 ml purified water, and mix completely. Coat the plates with an applicator, allow them to air dry 30 minutes and dry in oven at 110°C for 1 hour. Store in a desiccator over Silica Gel.

Solvent System. Measure into a separatory funnel, 80 ml pyridine, 120 ml n-butanol, 24 ml glacial Acetic Acid and mix. Add about 50 ml of 20% W/V Ammonium Chloride in purified water and shake gently. Draw off the bottom layer and discard. Place enough of the mixture to saturate a paper lined chromatographic chamber and allow to equilibrate for 30 minutes before use.

Sample Preparation. Prepare accurately a solution of sample powder in purified water to contain 2 mg/ml calculated on the anhydrous basis.

In addition, prepare the following solutions:

a) Prepare accurately a solution in purified water to contain 0.4 mg/ml each of (2 Beta, 16 Beta-Dipiperdino-5 alpha-Androstane-3 alpha, 17 Beta-diol 3 alpha-Acetate Dimethobromide) plus (2 Beta, 16 Beta-Dipiperdino-5 alpha-Androstane-3 alpha, 17 Beta-diol, 17 Beta-Acetate Dimethobromide). Call this solution "a".

b) 1.0, 2.0, 4.0% Standards. Into three separate 10 ml volumetric flasks, weight accurately 20 mg of Pancuronium Bromide standard calculated on the anhydrous basis and dissolve in about 5 ml purified water. Pipet into one flask 0.25 ml of solution "a", 0.5 ml of solution "a" into a second flask, and 1.0 ml of solution "a" into a third flask. Dilute each flask to volume with purified

1/ See footnote 2/ on page 7 for the type of Silica Gel to be used.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:39Z
Check the source to verify that this is the current version before use.

water and mix. These correspond to 1% total, 2% total and 4% total of related foreign steroids respectively.

Procedure. Apply to the prepared plates, 5 microliters each of the sample preparation and standard preparations about 2 cm apart and 2 cm from the bottom of the plate. Place the plate in the prepared chamber, allow the solvent to move about 17 cm, remove the plate from the chamber and allow it to dry in a current of air in a fume hood for at least one hour. Spray the plate evenly and completely with a fine mist of Sulfuric acid in alcohol (20% V/V) and heat at 110°C for one hour. Examine the plate under ultraviolet light. The main spot of the sample conforms to the main spot of the standard in Rf, size and color (Identity test a). Any secondary spots appearing in the sample are not greater in size or intensity than the secondary spots of the standard. The total amount of related foreign steroids (decomposition) may not exceed 2% combined.

4.5.2.6 Assay for Pancuronium Bromide Powder.

Reagents.

1. 0.1N Perchloric Acid. To about 500 ml acetic acid contained in a 1000 ml volumetric flask, add 9 ml Perchloric Acid, reagent grade (70%) with mixing. Add 24.5 ml of acetic anhydride, mix, and dilute to mark with acetic acid. Standardize with Potassium Biphthalate according to U.S.P.

2. Crystal Violet Solution - 0.5% W/V in acetic acid.

3. Mecuric Acetate Solution. Dissolve 1.5 gram of Mercuric Acetate, reagent grade in 50 ml acetic acid. Add 1 drop crystal violet solution and titrate to a green end point with 0.1N Perchloric acid.

Procedure.

(Note: Perform titrations rapidly to avoid difficulties in determining end-point due to precipitate formation.)

Weigh accurately about 300 mg of sample powder into a 50 ml Erlenmeyer flask. Correct the sample weight for water content. In the event the loss on drying exceeds the water content, correct the sample weight for the percent loss on drying.



Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:39Z
Check the source to verify that this is the current version before use.

Add 10 ml acetic acid and dissolve the sample by gentle warming if necessary.

Add 10 ml mercuric acetate solution, and 3 drops crystal violet solution. Mix and titrate with standardized 0.1N Perchloric Acid to a green end-point. Perform a blank titration and make any necessary corrections.

Calculation:

$$\frac{\text{ml 0.1N Perchloric Acid Consumed} \times \text{N Perchloric Acid} \times 366.3 \times 100}{\text{corrected sample weight in mg.}} = \text{percent Pancuronium Bromide}$$

5. PACKAGING.

5.1 Packaging requirements. The requirements for packaging shall be in accordance with all applicable requirements of PPP-C-186 (see 6.3) and as specified herein.

5.2 Packaging. Packaging shall be level A or C, as specified (see 6.1).

5.2.1 Unit of issue. One box containing 25 ampules of size 1, strength 1, or size 2, strength 1 or one box containing 25 vials of size 3, strength 2, as specified, constitutes one unit of issue.

5.2.2 Level A.

5.2.2.1 Immediate container.

5.2.2.1.1 Size 1, strength 1, and size 2, strength 1. The immediate container for size 1, strength 1, and size 2, strength 1, shall be a commercially available ampul considered acceptable by the Food and Drug Administration (FDA) for the product contained therein and shall be of such compatibility that neither the contents nor the container are altered in any way by each other.

5.2.2.1.2 Size 3, strength 2. The immediate container for size 3, strength 2, shall be a commercially available multiple dose vial considered acceptable by the Food and Drug Administration (FDA) for the product contained therein and shall be of such compatibility that neither the contents nor the container are altered in any way by each.

5.2.2.2 Unit packaging. Each unit shall be packaged as specified in 5.2.5 of PPP-C-186. For size 1, strength 1, and size 2, strength 1, unit box having a board thickness of .020 inch may be furnished.

Source: https://assist.dla.mil -- Downloaded: 202640-21T10:39Z
Check the source to verify that this is the current version before use.

5.2.2.3 Intermediate packaging. Twelve (12) units of issue of size 1, strength 1 shall be packaged as specified in 5.2.7.1 of PPP-C-186. Twelve (12) units of issue of size 2, strength 1, shall be packaged as specified in 5.2.8 of PPP-C-186. Twelve (12) units of issue of size 3, strength 2, shall be packaged as specified in 5.2.9 of PPP-C-186.

5.2.3 Level C. Units shall be packaged in standard commercial containers of the size and kind commonly used, which will afford the degree of protection required for shipment and use of the product for its intended purpose.

5.3 Packing. Packing shall be level A, B or C, as specified (see 6.1).

5.3.1 Level A. Seventy-two (72) units of issue of size 1, strength 1, or forty-eight (48) units of issue of size 2, strength 1, or twenty-four (24) units of issue of size 3, strength 2, shall be packed as specified in 5.4.4 of PPP-C-186.

5.3.2 Level B. Seventy-two (72) units of issue of size 1, strength 1, shall be packed as specified in 5.4.3.1.1 of PPP-C-186. Forty-eight (48) units of issue of size 2, strength 1, or twenty-four (24) units of issue of size 3, strength 2, shall be packed as specified in 5.4.3.2 of PPP-C-186.

5.3.3 Level C. The subject commodity shall be packed in substantial commercial containers of the type, size, and kind commonly used for the purpose, so constructed as to insure acceptance and safe delivery by common or other carriers, at the lowest rate, to point of delivery called for in the contract or purchase order.

5.4 Marking.

5.4.1 Labeling. Labeling shall be in accordance with the requirements of the Federal Food, Drug and Cosmetic Act and shall include the information required below:

5.4.1.1 Immediate container. Each immediate container label shall bear the following information:

(a) label information in accordance with commercial practice.
(b) the expiration date.

5.4.1.2 Unit package. Each unit package label shall bear the following information:

(a) label information in accordance with commercial practice.
(b) the National Stock Number.

NOTES: (1) The National Stock Number shall not be preceded by any prefix.
(2) Strip labeling may be used.
(3) Label adhesion test shall not apply.

Source: https://assist.dla.mil -- Downloaded: 202640-21T10:39Z
Check the source to verify that this is the current version before use.

5.4.2 Intermediate package. Each intermediate package shall be marked as specified in paragraph 5.5.3 of PPP-C-186. In addition, Type I shelf life marking as specified in MIL-STD-129 shall be shown. Marking shall include the legends: "STORE BETWEEN 2° - 8°C (35° - 46°F.)" and "DO NOT PERMIT TO FREEZE".

5.4.3 Exterior container. Exterior containers shall be marked as specified in 5.5.4 of PPP-C-186. Type I shelf life marking as specified in MIL-STD-129 shall also be shown. Marking shall include the legends: "STORE BETWEEN 2° - 8°C (35° - 46°F.)" and "DO NOT PERMIT TO FREEZE".

NOTE: The unrefrigerated shipping time for this item shall be not more than 7 days (168 hours).

6. NOTES

6.1 Ordering data. Procurement documents should specify the following:

(a) Number and date of this specification.
(b) National Stock Number.
(c) Selection of applicable levels of packaging and packing (See 5.2 and 5.3).

6.2 This specification covers the following item(s):

| National Stock No. | Item Identification | Specification Designation |
|---|---|---|
| 6505-00-149-0107 | PANCURONIUM BROMIDE INJECTION, 2 mg per ml, 2 ml, 25s | Size 1, strength 1 |
| 6505-00-149-0108 | PANCURONIUM BROMIDE INJECTION, 2 mg per ml, 5 ml, 25s | Size 2, strength 1 |
| 6505-01-012-7559 | PANCURONIUM BROMIDE INJECTION, 1 mg per ml, 10 ml, 25s | Size 3, strength 2 |

6.3 All references to PPP-C-186 refer to Interim Federal Specification PPP-C-00186B which is mandatory for use until superseded by a revision.

Custodians:
Army - MD
Navy - MS
Air Force - 03

Preparing activity:
Defense Supply Agency - DM

Project No. 6505-1548



Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:39Z
Check the source to verify that this is the current version before use.

# STANDARDIZATION DOCUMENT IMPROVEMENT PROPOSAL

OMB Approval
No. 22-R255

INSTRUCTIONS: The purpose of this form is to solicit beneficial comments which will help achieve procurement of suitable products at reasonable cost and minimum delay, or will otherwise enhance use of the document. DoD contractors, government activities, or manufacturers/vendors who are prospective suppliers of the product are invited to submit comments to the government. Fold on lines on reverse side, staple in corner, and send to preparing activity. Comments submitted on this form do not constitute or imply authorization to waive any portion of the referenced document(s) or to amend contractual requirements. Attach any pertinent data which may be of use in improving this document. If there are additional papers, attach to form and place both in an envelope addressed to preparing activity.

DOCUMENT IDENTIFIER AND TITLE
MIL-P-37415 PANCURONIUM BROMIDE INJECTION

NAME OF ORGANIZATION AND ADDRESS

CONTRACT NUMBER

MATERIAL PROCURED UNDER A

☐ DIRECT GOVERNMENT CONTRACT ☐ SUBCONTRACT

1. HAS ANY PART OF THE DOCUMENT CREATED PROBLEMS OR REQUIRED INTERPRETATION IN PROCUREMENT USE?
   A. GIVE PARAGRAPH NUMBER AND WORDING.

   B. RECOMMENDATIONS FOR CORRECTING THE DEFICIENCIES

2. COMMENTS ON ANY DOCUMENT REQUIREMENT CONSIDERED TOO RIGID

3. IS THE DOCUMENT RESTRICTIVE?

   ☐ YES ☐ NO *(If "Yes", in what way?)*

4. REMARKS

SUBMITTED BY *(Printed or typed name and address - Optional)*

TELEPHONE NO.

DATE

DD FORM 1 JAN 72 1426 REPLACES EDITION OF 1 JAN 66 WHICH MAY BE USED

Source: https://assist.dla.mil -- Downloaded: 2026-01-21 10:39Z
Check the source to verify that this is the current version before use.