MILITARY SPECIFICATION

SULFANILAMIDE, ALLANTOIN, AND AMINACRINE HYDROCHLORIDE CREAM

This specification is approved for use by all Departments and Agencies of the Department of Defense.

1. SCOPE

1.1 Scope. This specification covers Sulfanilamide, Allantoin, and Aminacrine Hydrochloride Cream (See 6.2).

2. APPLICABLE DOCUMENTS

2.1 The following documents, of the issue in effect on date of invitation for bids or request for proposals, form a part of this specification to the extent specified herein:

SPECIFICATIONS

Federal

PPP-C-186	Containers, Packaging and Packing for Drugs, Chemicals, and Pharmaceuticals. (See 6.3)

STANDARDS

Military

MIL-STD-105	Sampling Procedures and Tables for Inspection by Attributes.

FSC 6505

Source: https://assist.dla.mil -- Downloaded: 202610-21T10:16Z
Check the source to verify that this is the current version before use.

(Copies of specifications and standards required by contractors in connection with specific procurement functions should be obtained from the procuring activity or as directed by the contracting officer.)

2.2 Other publications. The following documents form a part of this specification to the extent specified herein. Unless otherwise indicated, the issue in effect on date of invitation for bids or request for proposals shall apply.

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
FOOD AND DRUG ADMINISTRATION

Federal Food, Drug, and Cosmetic Act and
Regulations Promulgated Thereunder.

(Application for copies should be addressed to the Food and Drug Administration, Washington, DC 20204.)

U. S. PHARMACOPEIAL CONVENTION, INC.

The United States Pharmacopeia.

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

National Formulary.

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

3. REQUIREMENTS

3.1 Material. Shall be a cream, uniform in texture and color containing Sulfanilamide 15%, Allantoin 2%, 9-Aminoacridine Hydrochloride 0.2% with lactose in a water miscible, non-greasy base made from propylene glycol, Stearic acid, diglycol stearate, triethanolamine, buffered with lactic acid to an acid pH. The lactic acid may be a 50 percent concentration, in lieu of the USP concentration.

3.1.1 pH. The finished preparation shall have a pH of not less than 4.0 and not more than 4.6, when determined as specified in 4.5.1.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

3.1.2 Assay. The finished preparation shall assay to contain not less than 90.0 percent and not more than 110.0 percent of the required amount of sulfanilamide and of 9-aminoacridine hydrochloride when determined as specified in 4.5.2.1 and 4.5.2.2, respectively. In addition, the cream shall assay to contain not less than 90.0 percent and not more than 110.0 percent of the required amount of allantoin when determined by a suitable, accurate, and reproducible method.

3.1.2.1 The bidder or offeror shall submit, with its bid or proposal, the proposed method of assay for the allantoin in the finished preparation. The method shall be subject to approval and acceptance by the Government. If the method is approved and accepted by the Government, such method shall be employed in the testing of the material.

3.1.3 Identity. The sulfanilamide, allantoin, and 9-aminoacridine hydrochloride in the finished cream shall be identified according to the procedure specified for each ingredient as determined in 4.5.3.1, 4.5.3.2, and 4.5.3.3, respectively.

3.1.4 Accelerated aging test. All lots of finished cream are subject to compliance with the following accelerated aging test. The contractor, as a minimum, is required to perform the aging test on the first 2 lots and on each subsequent consecutive fourth lot for each contract. At its discretion, the Defense Personnel Support Center may test any and all lots to determine compliance with the aging test:

> Select 6 tubes and subject 3 tubes to 40° C ± 1° C for 21 consecutive days and the other 3 tubes shall be stored at 70° - 80°F, as unaged control samples. The heated samples shall be returned to room temperature (70° - 80° F.) prior to test. The cream in the tubes shall be tested in accordance with the requirements for pH, assay, and color. All results shall be within the requirements of the specification, except that the color of the cream may become slightly darker after being subjected to the aging test.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

3.1.5 Bacterial and mold count. The cream shall not contain more than 20 viable microorganisms per gram. There shall be no organisms of staphylococcus aureus or the pseudomonas group and no molds. Suitable, accurate, and reproducible methods shall be employed in determining the microorganism count and presence or absence of organisms of staphylococcus aureus or the pseudomonas group and molds. Not less than 5 units, selected at random from each lot or control, shall be tested to determine compliance with the bacterial count requirements, and the presence or absence of organisms of staphylococcus aureus or the pseudomonas group and molds.

A coagulase negative response to the U.S.P. test for staphylococcus aureus and pseudomonas aeruginosa, will satisfy the requirement for no staphylococcus aureus and no organisms of the pseudomonas group.

3.2 Active ingredients.

3.2.1 Sulfanilamide. The sulfanilamide entering into the manufacture of the cream shall be in accordance with the tests, standards, and requirements of the N.F.XI, including any supplements or revision thereto, which was official from 1 October 1960 to 31 August 1965, except the assay shall not be less than 99.0% and not more than 100.5% Sulfanilamide and the melting range shall be 163.5°C - 166.5°C by USP procedure, Class I.

3.2.2 Allantoin. The allantoin entering into the manufacture of the cream shall comply with the following:

3.2.2.1 Solution and reaction. One gram dissolved in 250 ml of purified water shall produce a clear solution. Solution shall be colorless or have no more color than A.P.H.A. color standard 10, when tested in accordance with the method specified in the "Standard Methods for the Examination of Water and Wastewater," Twelfth Edition, Third Printing, October 1967, or any subsequent printing or edition published by the American Public Health Association, Inc.

3.2.2.2 Residue on ignition. Shall be not more than 0.05 percent.

3.2.2.3 Substances darkened by Sulfuric Acid, U.S.P.. Solution shall be colorless when 0.5 gram Allantoin is dissolved in 5 ml Sulfuric Acid, U.S.P.

3.2.2.4 Chloride. Dissolve 1.5 gram in 75 ml of boiling purified water. Cool well, dilute to 75 ml with purified water, mix well and filter. To 25 ml of filtrate, add 0.5 ml of nitric acid and 1 ml of silver nitrate T.S. Any opalescence produced shall correspond to not more than 0.03 mg of Chloride.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

3.2.2.5 Sulfate. Evaporate 25 ml of the filtrate obtained in the chloride test to about 10 ml. Cool, filter, if necessary, and add to the filtrate 0.5 ml N hydrochloric acid and 1 ml of barium chloride T.S. Any turbidity produced shall correspond to not more than 0.1 mg of Sulfate.

3.2.2.6 Assay. Shall assay not less than 98.5 percent and not more than 102.0 percent Allantoin, when determined as specified in 4.5.4.1.

3.2.2.7 Ninety-four (94) percent of Allantoin powder shall pass through a U.S. No. 200 sieve and 100 percent must pass through a U.S. Sieve No. 100.

3.2.3 9-Aminoacridine hydrochloride. The 9-Aminoacridine Hydrochloride entering into the manufacture of the cream shall comply with the following:

3.2.3.1 Solution. Shall be a clear, yellow solution, when 0.5 gram is dissolved in 10 ml of hot, purified water.

3.2.3.2 Residue on ignition. Sulfated ash of 0.5 gram sample shall be negligible in weight.

3.2.3.3 Arsenic. Shall contain not more than 10 p.p.m. Arsenic when determined by the procedure described in the N.F.

3.2.3.4 Loss on drying. A 1-gram sample shall be dried to constant weight at 110° C. The loss of weight shall be not more than 8.0 percent.

3.2.3.5 Assay. Shall contain not less than 82.5 percent and not more than 84.5 percent of base when determined as specified in 4.5.4.2.

3.2.3.6 Hydrochloride. The hydrochloride content shall be not less than 15.52 percent and not more than 16.00 percent when determined as specified in 4.5.4.3.

3.2.3.7 Melting range. The base from the assay, above, shall melt between 233° and 237° C when determined by the U.S.P. Melting range method Ia, with the sample inserted at approximately 228° C and the temperature of the bath increased at the rate of 1° C per minute.

Source: https://assist.dla.mil -- Downloaded: 202610-21T10:16Z
Check the source to verify that this is the currentversion before use.

3.3 Other ingredients. All other ingredients entering into the manufacture of the cream shall be of U.S.P. or N.F. quality, or, if not included in either of these compendia they shall be suitable for vaginal use. If 50 percent lactic acid is used, it shall comply with the tests, standards, and requirements of the U.S.P., except that the assay limits shall be not less than 47.5 percent and not more than 52.5 percent lactic acid.

3.4 Delivery schedule. Not more than 4 months shall have elapsed from the date of manufacture of the product to the date of delivery to the Government.

3.5 Workmanship. Workmanship shall be first class throughout. The material and its containers shall be free from defects which detract from their appearance or may impair their serviceability.

4. QUALITY ASSURANCE PROVISIONS

4.1 Supplier responsibility for inspection. Unless otherwise specified in the contract or purchase order, the supplier is responsible for the performance of all inspection requirements as specified herein. Except as otherwise specified in the contract or order, the supplier may use his own or any other facilities suitable for the performance of the inspection requirements specified herein, unless disapproved by the Government. The Government reserves the right to perform any of the inspections set forth in the specification where such inspections are deemed necessary to assure supplies and services conform to prescribed requirements.

4.1.1 Records of examinations and tests performed by or for the contractor shall be maintained by the contractor and made available to the Government, upon the Government's request, at any time, or from time to time, during the performance of the contract and, (i) as to any expiration dated item, for the expiration dating period (potency period) specified by the contractor for such item, or such longer period as may be required by regulation of any federal agency; or (ii) as to non-dated items, for not less than 3 years after delivery of the items to the Government.

4.1.2 No company supplying any ingredient(s) to the contractor will be considered an acceptable facility for the performance of any inspection requirements specified herein.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

4.2 Lot. For purposes of this specification, a lot, batch, or control is that single, uniform, and homogeneous quantity of cream produced from one formulation, subjected to the same compounding and manufacturing operations, and filled into final containers.

4.3 Sampling. Sampling shall be conducted in accordance with MIL-STD-105.

Table I. Sampling

For visual examination	Inspection level	AQL (percent defective	Unit of product
Major	S-2	1.0	Package
Minor	II	2.5	Package

4.3.1 For raw material and end item testing. Noncompliance with any of the following is cause for rejection.

Table II. For raw material and end item testing

Characteristic	Requirement paragraph	Inspection level*	Minimum sample
pH	3.1.1	S-2	5
Assay	3.1.2	S-2	5 composite
Identity	3.1.3	S-2	5
Accelerated aging test	3.1.4	S-2	5
Bacterial and mold count	3.1.5	- **	-
Ingredients :			
Allantoin	3.2.2		
9-Aminoacridine hydrochloride	3.2.3		

* Where possible, the sample shall be used for two or more tests.
** Sufficient quantities of each of the raw materials shall be selected from each lot to perform tests indicated in applicable test methods. Failure of any sample to comply with test requirements shall be cause for rejection of the lot represented.

4.3.2 Samples. If (i) the quality assurance representative submits samples to the Quality Assurance Branch, Technical & Quality Assurance Division, Directorate of Medical Materiel, Defense Personnel Support Center or (ii) the bidder (offeror)

Source: https://assist.dla.mil -- Downloaded: 202610-21T10:16Z
Check the source to verify that this is the currentversion before use.

is required by the terms of the Schedule or otherwise to submit samples to said Branch, then a quantity of each active ingredient specified in this document, sufficient to perform all of the tests stated herein, and 100 mg of 9-aminoacridine hydrochloride powder, accompanied by a certificate of analysis listing all test results, shall be submitted with the finished product to the Directorate of Medical Materiel, Technical & Quality Assurance Division, Defense Personnel Support Center, Attention: Quality Assurance Branch, 2800 South 20th Street, Philadelphia, PA 19101.

4.4 Examination. The finished cream shall be examined to determine compliance with all requirements of this specification. Nonconformance will be permitted to the extent indicated in 4.3 and 4.3.1.

4.4.1 Classification of defects. Examination will be conducted in accordance with the following classification of defects:

Categories and defects*

Major	
101	Cream not free from foreign matter.
102	Cream not uniform in color.
103	Cream not homogeneous.
Minor	
201	Exterior of immediate container is not free from contents or foreign matter.
202	Label not free from defacing marks and tears.

* Inspection is not restricted to the classified possible defects listed above.

4.5 Tests.

4.5.1 pH. Mix approximately 5 grams of cream with 20 ml of purified water in a 50 ml beaker and determine the pH potentiometrically at 25° C., using the U.S.P. method.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

4.5.2 Assay.

4.5.2.1 Assay for sulfanilamide. Accurately weigh approximately 3 grams of the cream into a 250-ml beaker, followed by approximately 50 ml of purified water and 8 ml concentrated HCl. Heat the mixture on a steam bath until the cream completely melts. Stir the mixture thoroughly and remove from the heat. Cool the mixture in an ice bath. Add about 25 grams of cracked ice and titrate with tenth-molar sodium nitrite using starch iodide test paper as the indicator. When the titration is complete, the endpoint is reproducible after the mixture has been allowed to stand for 1 minute. Each ml of 0.1M sodium nitrite is equivalent to 17.22 mg of $C_6H_8N_2O_2S$. Calculate the percent of required amount of sulfanilamide.

4.5.2.2 Assay for 9-aminoacridine hydrochloride. Accurately weigh a sample of cream, equivalent to 10 mg aminacrine hydrochloride, into a small beaker. (Note: Weigh the cream to within ± 150 mg of nominal amount required.) Add 25 ml of 1N hydrochloric acid solution and heat on a steam bath. After the cream melts, mix thoroughly. Avoid loss of sample from splashing. Stop mixture and allow the molten fatty acids to form on the surface.

Rapidly cool the mixture until the stearic acid congeals. Decant the liquid through a pledget of glass wool into a 1-liter volumetric flask. Leave as much of the congealed fatty acids, as is possible, in the beaker. Repeat this entire extraction operation with three more 25 ml portions of 1N hydrochloric acid solution. Wash the glass wool filter with purified water. Dilute to volume with purified water. Mix well. Pipet 5.0 ml of the clear solution into a 500-ml volumetric flask. Dilute with 0.1N hydrochloric acid. Mix well. Pipet 10.0 ml of this solution into a 100-ml volumetric flask, dilute with 0.1N hydrochloric acid. Mix well. Compare the fluorescence emission of the sample with that of a standard containing 0.00001 mg of aminacrine hydrochloride per ml in 0.1N hydrochloric acid, using a suitable instrument with an activating wavelength of approximately 392 nm and a fluorescent emission wavelength of approximately 455 nm.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

Calculation:

$$\frac{Fu}{Fs} \times 0.00001^{*} \text{ mg/ml} \times 100 \times \frac{500}{10} \times \frac{1000}{5} \times \frac{100}{\text{sample wt (mg)}} =$$

percent Aminacrine Hydrochloride in the cream.

*depending on the exact concentration of the final standard solution (mg/ml).
Fu = fluorescence emission of the sample solution.
Fs = fluorescence emission of the standard solution.

4.5.3 Identity.

4.5.3.1 Sulfanilamide. An orange precipitate shall be produced when finished cream is treated as follows:

To approximately 1 gram of cream in a 150-ml beaker, add 25 ml of 4 percent sodium hydroxide solution. Place the solution on the steam bath and warm gently until the cream completely melts. Cool and filter the mixture through a suitable filter paper (Whatman No. 4, or equivalent). To a test tube immersed in an ice bath, add 1 ml of filtrate, 1 ml of 10 percent hydrochloric acid solution, and mix well. Then add 4 ml of 1 percent aqueous sodium nitrite solution and mix well. To this mixture add a drop or two of a solution of 50 mg betanaphthol in 2 ml of 10 percent sodium hydroxide solution.

4.5.3.2 Allantoin. Extract 5 gram of the cream with 20 ml of hot purified water, cool, and filter. Add 2 drops of mercuric nitrate T.S. to 10 ml* of the cooled solution. A white precipitate forms which is soluble in an excess of the reagent.

*Save remainder of the solution for test in 4.5.3.3.

4.5.3.3 9-aminoacridine hydrochloride. Dilute the remainder of the above hot purified water extraction to 500 ml with purified water. A bluish fluorescence should be apparent. This fluorescence shall become immediately apparent when viewed under a U.V. light.

Source: https://assist.dla.mil -- Downloaded: 202610-21T10:16Z
Check the source to verify that this is the currentversion before use.

4.5.4 Ingredients.

4.5.4.1 Allantoin. Accurately weigh approximately 250 mg of sample, previously dried, into a glass-stoppered flask. Pipet 50 ml of ethylenediamine anhydrous into the flask and stopper. Shake until the sample is dissolved. Titrate the solution with 0.1N Sodium Methoxide Volumetric solution using 1 ml of 0.1 percent 0-nitro-aniline (in benzene) indicator until the solution changes from yellow to orange. Correct for the volume of the sodium methoxide consumed by titrating 50 ml of ethylene-diamine anhydrous plus 1 ml of indicator. Each ml of 0.1N sodium methoxide is equivalent to 15.81 mg of Allantoin.

Calculation:

$$\text{mg of 0.1N } NaOCH_3 \text{ (Corrected)} \times 15.81 \times \frac{100}{\text{Sample wt (mg)}} = \text{percent allantoin}$$

4.5.4.2 9-Aminoacridine hydrochloride. Accurately weigh approximately 500 mg of sample into a 100-ml volumetric flask. Add 10 ml of 20 percent reagent Sodium Acetate solution. Dilute to about 50 ml, stopper flask and shake until sample is dissolved. Add an excess of 10 percent Sodium Hydroxide solution to free the base. Swirl for several minutes and permit to stand for 15 minutes, occasionally agitating the solution. Dilute to volume with purified water. Mix well and filter mixture through a previously prepared tared filter crucible, catching the filtrate in a clean, dry vacuum flask. Transfer the undiluted filtrate to another flask and set aside to be used in the determination of Hydrochloride content. Rinse the 100-ml volumetric flask with purified water and transfer the rinsings to the filter crucible. Wash the precipitated base with several small portions of purified water until the washings are no longer alkaline to phenolphthalein. Dry to constant weight at 100° C. (Save for melting range.)

4.5.4.3 Hydrochloride content. Transfer exactly 75 ml of the undiluted filtrate from the aminacrine assay to a clean, dry 250-ml iodine flask. Render the solution acid to litmus with dilute Nitric Acid. Add 10 additional ml of acid, plus exactly 50.0 ml of 0.1N Silver Nitrate solution. Add 3 ml of Nitrobenzene and shake well. Rinse the stopper and walls of the flask with purified water. Add approximately 3 ml of Ferric Ammonium Sulfate T.S. and mix well. Titrate the excess Silver Nitrate with 0.1N Ammonium or Potassium Thiocyanate Volumetric Solution until the characteristic flesh color appears. Each ml of 0.1N Silver Nitrate is equivalent to 3.646 mg of Hydrochloride.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

Calculation:

$$(A-B) \times 3.646 \times \frac{100}{75} \times \frac{100}{\text{Sample (mg)}} \times \frac{100}{(100-\%\ \text{LOD})} = \text{percent hydrochloride}$$

where:

A = 50.0 ml 0.1N $AgNO_3$

B = ml 0.1N NH_4SCN or KSCN consumed

(Make appropriate corrections for exact normalities of volumetric solutions).

4.6 Examination of preparation for delivery. An examination shall be made to determine whether the packaging, packing, and marking comply with the requirements of Section 5. Defects shall be scored as specified in Table III. Sampling shall be in accordance with MIL-STD-105. The sample unit shall be one container fully prepared for delivery. The lot shall be the number of containers offered for inspection at one time. The inspection level shall be S-2 and the acceptable quality level shall be 4.0 defects per hundred units. For labeling, the acceptance number is zero.

Table III. Examination of preparation for delivery

Examination	Defect
Containers	Not as specified.
Quantity	Quantity in interior packages and shipping containers not as specified.
Packaging and Packing Materials	Any component missing or damaged.
Workmanship	Inadequate application of components such as incomplete closure of container flaps, tape improperly applied, or containers distorted.
Markings	Omitted, incorrect, illegible, improper size, location, sequence, or method of application.

NOTE: The above examination is only applicable to the unit, intermediate and exterior containers.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

5. PREPARATION FOR DELIVERY

5.1 Preparation for delivery shall be in accordance with all applicable requirements of PPP-C-186, and as specified herein.

5.2 Packaging. Packaging shall be level A or C as specified (see 6.1).

5.2.1 Unit of issue. One package containing one tube and applicator or dispenser as specified, constitutes one unit of issue.

5.2.2 Level A.

5.2.2.1 Immediate containers. Immediate container shall be a commercially available tube type container with applicator or dispenser for vaginal use considered acceptable by the Food and Drug Administration (FDA) for the product contained therein and shall be of such compatability that neither the contents nor the container are altered in any way by each other.

5.2.2.2 Unit package. Each unit shall be packaged as specified in 5.2.5 of PPP-C-186. The box may be constructed of five panels. One panel being a perforated tear-off panel.

5.2.2.3 Intermediate package. Twelve (12) units of issue shall be intermediate packaged as specified in 5.2.1.3 of PPP-C-186.

5.2.3 Level C. Units shall be packaged in standard commercial containers of the size and kind commonly used, which will afford the degree of protection required for shipment and use of the product for its intended purpose.

5.3 Packing. Packing shall be level A, B or C as specified (see 6.1).

5.3.1 Level A. Seventy-two (72) units of issue shall be packed as specified in 5.4.4 of PPP-C-186.

5.3.2 Level B. Seventy-two (72) units of issue shall be packed as specified in 5.4.3.5.1 of PPP-C-186.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

5.3.3 Level C. The subject commodity shall be packed in substantial commercial containers of the type, size, and kind commonly used for the purpose, so constructed as to insure acceptance and safe delivery by common or other carriers, to point of delivery called for in the contract or purchase order.

5.4 Marking.

5.4.1 Labeling. Labeling shall be in accordance with the requirements of the Federal Food, Drug and Cosmetic Act, and shall include the information required below.

5.4.1.1 Immediate container. Each immediate container label shall bear the following:

(a) label information in accordance with commercial practice.
(b) the date of manufacture prefixed by "MFD".

5.4.1.2 Unit package. Each unit package shall bear the following:

(a) labeling information in accordance with commercial practice.
(b) the national stock number.

Notes: (1) The national stock number shall not be preceded by any prefix.
(2) Strip labeling may be used.
(3) Label adhesion test shall not apply.

5.4.2 Intermediate package. Each intermediate package shall be marked as specified in 5.5.3 of PPP-C-186 except that the date of manufacture shall be shown in lieu of the date packed. Marking shall include the legend:

"STORE AT CONTROLLED ROOM TEMPERATURE (59° - 86°F)"

5.4.3 Exterior container. Exterior container shall be marked in accordance with 5.5.4 of PPP-C-186 except that the date of manufacture shall be shown in lieu of the date packed. Marking shall include the legend:

"STORE AT CONTROLLED ROOM TEMPERATURE (59° - 86°F)"

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

6. NOTES

6.1 Ordering data. Procurement documents should specify the following:

(a) Title, number and date of this specification.
(b) the National Stock Number.
(c) Selection of applicable levels of packaging and packing (See 5.2 and 5.3).

6.2 This specification covers the following item(s):

National Stock No.	Item Identification
6505-00-890-2217	SULFANILAMIDE, ALLANTOIN, AND AMINACRINE HYDROCHLORIDE CREAM, Vaginal, 4 oz (113.4 Grams)

6.3 All references to PPP-C-186 refer to Interim Federal Specification PPP-C-00186B which is mandatory for use until superceded by a revision.

6.4 This specification does not cover all types, classes, grades, or sizes of the commodity indicated by the title of this specification, or those which are commercially available, but is intended to cover the types which are normally procured to meet Military requirements.

Custodians:
Army - MD
Navy - MS
Air Force - 03

Preparing activity:
Defense Supply Agency - DM

Project No. 6505-1392
cs

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

STANDARDIZATION DOCUMENT IMPROVEMENT PROPOSAL

OMB Approval
No. 22-R255

INSTRUCTIONS: The purpose of this form is to solicit beneficial comments which will help achieve procurement of suitable products at reasonable cost and minimum delay, or will otherwise enhance use of the document. DoD contractors, government activities, or manufacturers/vendors who are prospective suppliers of the product are invited to submit comments to the government. Fold on lines on reverse side, staple in corner, and send to preparing activity. Comments submitted on this form do not constitute or imply authorization to waive any portion of the referenced document(s) or to amend contractual requirements. Attach any pertinent data which may be of use in improving this document. If there are additional papers, attach to form and place both in an envelope addressed to preparing activity.

DOCUMENT IDENTIFIER AND TITLE
MIL-S-37241 SULFANILAMIDE, ALLANTOIN, AND AMINACRINE HYDRO

NAME OF ORGANIZATION AND ADDRESS

CONTRACT NUMBER

MATERIAL PROCURED UNDER A
☐ DIRECT GOVERNMENT CONTRACT ☐ SUBCONTRACT

1. HAS ANY PART OF THE DOCUMENT CREATED PROBLEMS OR REQUIRED INTERPRETATION IN PROCUREMENT USE?
   A. GIVE PARAGRAPH NUMBER AND WORDING.

   B. RECOMMENDATIONS FOR CORRECTING THE DEFICIENCIES

2. COMMENTS ON ANY DOCUMENT REQUIREMENT CONSIDERED TOO RIGID

3. IS THE DOCUMENT RESTRICTIVE?
   ☐ YES ☐ NO *(If "Yes", in what way?)*

4. REMARKS

SUBMITTED BY *(Printed or typed name and address - Optional)*

TELEPHONE NO.

DATE

DD FORM 1 JAN 72 1426 REPLACES EDITION OF 1 JAN 66 WHICH MAY BE USED

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.

FOLD

DEFENSE PERSONNEL SUPPORT CENTER
2800 South 20th Street
Philadelphia, PA. 19101

POSTAGE AND FEES PAID
DEFENSE SUPPLY AGENCY
DoD - 304

U.S.MAIL

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE $300

HEADQUARTERS, DEFENSE PERSONNEL SUPPORT CENTER
ATTN: DIRECTORATE OF MEDICAL MATERIEL, CODE ATT
2800 SOUTH 20TH STREET
PHILADELPHIA, PA 19101

FOLD

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:16Z
Check the source to verify that this is the current version before use.