MILITARY SPECIFICATION

DOXEPIN HYDROCHLORIDE CAPSULES

This specification is approved for use by all Departments and Agencies of the Department of Defense.

1. SCOPE

1.1 Scope. This specification covers Doxepin Hydrochloride Capsules (See 6.2).

2. APPLICABLE DOCUMENTS

2.1 Issues of documents. The following documents, of the issue in effect on date of invitation for bids or request for proposals, form a part of this specification to the extent specified herein.

SPECIFICATIONS

FEDERAL

PPP-C-186 - Containers, Packaging and Packing for Drugs, Chemicals and Pharmaceuticals. (See 6.3)

STANDARDS

FEDERAL

FED-STD-00285 - Capsules (for Medicinal Purposes).

MILITARY

MIL-STD-129 - Marking for Shipment and Storage.

(Copies of specifications and standards required by contractors in connection with specific procurement functions should be obtained from the procuring activity or as directed by the contracting officer.)

Beneficial comments (recommendations, additions, deletions) and any pertinent data which may be of use in improving this document should be addressed to: Hq, Defense Personnel Support Center, ATTN: Directorate of Medical Materiel, DPSC-ATT, 2800 South 20th Street, Philadelphia, PA 19101, by using the self-addressed Standardization Document Improvement Proposal (DD Form 1426) appearing at the end of this document, or by letter.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

2.2 Other publications. The following documents form a part of this specification to the extent specified herein. Unless otherwise indicated, the issue in effect on date of invitation for bids or request for proposals shall apply.

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE, FOOD AND DRUG ADMINISTRATION

Federal Food, Drug, and Cosmetic Act and Regulations Promulgated Thereunder.

(Application for copies should be addressed to the Food and Drug Administration, Washington, DC 20204.)

U. S. PHARMACOPEIAL CONVENTION, INC.

The United States Pharmacopeia.

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

National Formulary.

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

3. REQUIREMENTS

3.1 Material. Shall be Doxepin Hydrochloride Capsules and shall be in accordance with all applicable requirements of FED-STD-00285, and as specified herein.

S2. Classification. Shall be type I or type II.

S5.4 The following is added to this paragraph:

Shall be capsules containing Doxepin Hydrochloride equivalent to 25 mg of Doxepin.

S6.2.6 Disintegration and solubility. Shall disintegrate in not more than 15 minutes, when tested using the U.S.P. apparatus and procedure for disintegration testing of uncoated tablets, using Simulated Gastric Fluid, T.S. as the immersion fluid. For the purpose of this test, complete disintegration is defined as that state in which any residue of the capsule remaining on the screen is a soft mass having no recognizable capsule shape.

S6.2.9 Weight variation. Shall comply with the USP Weight Variation Test for Capsules. In addition, the capsules shall comply with the USP Content Uniformity Test for Capsules when determined as specified in 4.3.1.1.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

3.1.1 Assay. The average of ten (10) individual capsule assays shall be not less than 90.0 percent and not more than 110.0 percent of the required amount of Doxepin, when determined as described in 4.3.1.1. The individual capsule assay shall meet the USP content uniformity test for the capsules.

3.1.2 Chromatographic identification. Shall show two major bands whose mobilities (R_f) are the same as the cis and trans isomers in the reference sample, when determined as specified in 4.3.1.2.

3.2 Active ingredient. The Doxepin Hydrochloride used in the manufacture of the finished capsules shall comply with the following requirements:

3.2.1 Assay (isomer content). The trans Doxepin content shall be not less than 72.0 percent and not more than 78.0 percent when determined as specified in 4.3.2.1.1. The cis Doxepin content shall be not less than 12.0 percent and not more than 16.0 percent when determined as specified in 4.3.2.1.2. The sum of the cis and trans doxepin contents shall be not less than 85.0 percent and not more than 92.0 percent.

3.2.2 Ionic chlorine content. Shall be not less than 10.9 percent and not more than 11.5 percent when determined as specified in 4.3.2.2.

3.2.3 Occluded solvents. Shall contain not more than 0.5 percent Benzene, not more than 0.5 percent Ethyl Acetate, and not more than 0.2 percent Methylene Chloride when determined as specified in 4.3.2.3.

3.2.4 Residue on ignition. Shall be not more than 2.0 percent when determined using the USP method.

3.2.5 Heavy metals. Shall be not more than 50 p.p.m. when determined using USP Method II.

3.2.6 Chromatographic examination.

3.2.6.1 Paper chromatography. Shall show two major bands whose mobilities are the same as those of the cis and trans isomers in the reference sample, when determined as specified in 4.3.2.4. The content of tertiary alcohol (11-(3-dimethylamino-1-propyl) 6H-dibenz [b, e] oxepin-11-ol) shall be not more than 0.5 percent when determined as specified in 4.3.2.5.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

3.2.6.2 Thin-layer chromatography. Shall contain not more than 0.5 percent ketone (6,11-dihydrodibenz [b,e] oxepin-11-one) when tested as specified in 4.3.2.6.

3.2.7 Total contaminants. The sum of the residue on ignition, heavy metals, benzene, ethyl acetate, methylene chloride, tertiary alcohol, and ketone shall be not more than 2.0 percent.

3.2.8 Melting range. Shall melt between 185° C. and 191° C., when determined using the USP Method, Class I.

3.2.9 Identity. Shall comply with the ultraviolet and infrared identity tests specified in 4.3.2.7 and 4.3.2.8.

3.3 Other ingredients. All other ingredients shall comply with S5.2 of FED-STD-00285.

3.4 Expiration dating period. The expiration dating period shall be not less than 36 months.

4. QUALITY ASSURANCE PROVISIONS

4.1 Supplier responsibility for inspection. Unless otherwise specified in the contract or purchase order, the supplier is responsible for the performance of all inspection requirements as specified herein. Except as otherwise specified in the contract or order, the supplier may use his own or any other facilities suitable for the performance of the inspection requirements specified herein, unless disapproved by the Government. The Government reserves the right to perform any of the inspections set forth in the specification where such inspections are deemed necessary to assure supplies and services conform to prescribed requirements.

4.1.1 Records of examinations and tests performed by or for the contractor shall be maintained by the contractor and made available to the Government, upon the Government's request, at any time, or from time to time, during the performance of the contract and, (i) as to any expiration dated item, for the expiration dating period specified by the contractor for such item, or such longer period as may be required by regulation of any federal agency; or (ii) as to non-dated items, for not less than 3 years after delivery of the items to the Government.

4.1.2 No company supplying any ingredient(s) to the contractor will be considered an acceptable facility for the performance of any inspection requirements specified herein.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

4.2 Sampling. Sampling shall be conducted in accordance with FED-STD-00285.

4.3 Tests.

4.3.1 Filled capsules.

4.3.1.1 Assay and content uniformity.

Reference Preparation.

Accurately weigh approximately 70 mg of doxepin hydrochloride reference powder and transfer to a 100-ml volumetric flask. Dissolve the reference powder in 0.01N methanolic hydrochloride acid solution, dilute to volume and mix. Pipet a 10 ml aliquot of the solution into a 200-ml volumetric flask, dilute to volume with methanolic hydrochloride acid and mix.

Procedure.

Transfer 10 capsules to two 100-ml volumetric flasks. Treat each sample as follows:

Add about 50 ml of 0.01N methanolic hydrochloric acid and shake on a mechanical shaker for 30 minutes. Dilute to volume with methanolic hydrochloric acid and mix. Transfer about 40 ml of the sample solution to a glass-stoppered centrifuge tube and centrifuge to obtain a clear supernatant liquid. Transfer a 30 ml aliquot of the clear supernatant solution to a 250-ml volumetric flask, dilute to volume with methanolic hydrochloric acid and mix well. Determine the absorbances of the sample and reference solutions relative to a methanolic hydrochloric acid blank in 1 cm cells, at approximately 297 nm, using a suitable spectrophotometer. Calculate the percent of the required amount, as follows:

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

% of required amount of Doxepin =

$$\frac{Au}{As} \times C \times \frac{250}{30} \times \frac{100}{25} \times 0.885 \times 100$$

where:

Au = absorbance of the sample
As = absorbance of the reference solution
C = final concentration of reference preparation in mg Doxepin Hydrochloride/ml
0.885 is the conversion factor for doxepin base from doxepin hydrochloride.

4.3.1.2 Chromatographic identification.

Reagents.

(1) Glycine-Sodium Chloride-Hydrochloric Acid Buffer, pH 1.6

Solution A. Dissolve 22.5 grams of glycine plus 17.5 grams of sodium chloride in 1000 ml of purified water.

Solution B. Dilute 28.4 ml of concentrated hydrochloric acid to 1000 ml with purified water.

Solution C. Mix 5.5 parts of solution A with 4.5 parts of solution B. The pH of this stock solution shall be 1.8.

Prior to use, dilute 35 ml of solution C to 100 ml with acetone.

(2) Developing solvent: Benzene-chloroform (2:1 V/V) saturated with ethylene glycol.

(3) Iodoplatinic acid reagent. Stock Solution is prepared as follows:

Platinic chloride 5% (W/V) in purified water------30 ml
Potassium Iodine 7.5% (W/V) in purified water----100 ml
Purified water---------- 70 ml

Filter this stock solution through glass wool. Dilute stock solution 1:3 with purified water prior to use.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

Chromatographic sheet.

Impregnate a sheet of Whatman No. 1 paper (approximately 7 inches by 22 inches) with Glycine- Sodium Chloride -Hydrochloric Acid Buffer. Dry at 60°C. for approximately 30 minutes, or air dry for 2 to 3 hours. Immediately prior to use, impregnate the sheet with a 7:3 solution of methanol and ethylene glycol, blot, and apply the sample.

Sample Preparation:

Transfer the contents of five capsules to a 15-ml glass-stoppered centrifuge tube. Add 5 ml of methanol. Shake the resulting suspension on a reciprocating shaker for 5 minutes and then centrifuge at 2000 r.p.m. for approximately 5 minutes. Filter through a small amount of glass wool.

Procedure:

Along the origin line of a prepared sheet, apply 0.020 ml of the sample preparation as a 1.5 inch long streak starting one inch from the edge of the sheet. Similarly, apply 0.020 ml of a solution of Doxepin Hydrochloride reference powder containing 25 mg per ml in chloroform as a 1.5 inch long streak starting one inch from the other edge of the sheet. Develop the sheet with the developing solvent using a descending technique for 20 hours ± 2 hours. After development, remove the sheet from the developing chamber, air dry for ten minutes and dry the sheet for approximately 30 minutes in an air circulating oven. Examine the dried chromatogram on a 254 nm ultraviolet scanner screen. In addition, spray the chromatogram with the iodoplatinic acid reagent and examine. The R_f values of the cis and trans isomers of the sample shall be identical to the R_f values of the cis and trans isomers of the reference powder.

4.3.2 Active ingredient.

4.3.2.1 Assay.

4.3.2.1.1 Trans isomer content.

Standard Curve.

Prepare a Standard Curve as follows:

Prepare five standard mixtures containing the following amounts of cis and trans Doxepin Hydrochloride isomers and dissolve each in 2 ml of Chloroform A.R.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

mg Trans Isomer	mg Cis Isomer	%Trans Isomer
95	5	95
90	10	90
85	15	85
80	20	80
75	25	75

Fill an 0.1 mm infrared sodium chloride cell with chloroform and insert it into the reference cell compartment of a suitable spectrophotometer. Add a chloroform solution of one of the standard cis trans isomer mixtures to another 0.1 mm cell and insert it into the sample cell compartment of the instrument. Record the infrared spectrum from 7.60 to 8.00 microns at least four times. Determine the absorbances of the maximum at 7.80 microns, using the base line technique. Similarly, for each standard cis-trans mixture, determine an average absorbance value from the replicate runs conducted. Prepare a standard reference curve by plotting the average absorbance values of the standard cis trans isomer mixtures at 7.8 against the corresponding percentages of the trans isomer in the mixtures.

Procedure:

Dissolve 100 mg of the sample of Doxepin Hydrochloride in 2 ml of chloroform. Using the procedure used for the standard mixtures, determine the average absorbance values at 7.80 microns by recording the infrared spectrum from 7.60 microns to 8.00 microns at least four times and, calculating the absorbance at 7.80 microns using the base line technique, and averaging the results. Derive the trans isomer content in the sample from the standard curve. The percentage of trans Doxepin in the sample is calculated as follows:

$$\%\text{Trans doxepin} = \%\text{trans doxepin hydrochloride} \times \frac{279.39}{315.85}$$

4.3.2.1.2 Cis isomer content.

Standard curve.

Prepare a standard curve as follows:

Prepare three standard mixtures containing the following amounts of cis and trans Doxepin Hydrochloride isomers and dissolve each in 1 ml of Chloroform, A.R.

mg Cis Isomer	mg Trans Isomer	%Cis Isomer
12.5	87.5	12.5
15.0	85.0	15.0
20.0	80.0	20.0

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

Fill an 0.1 mm infrared sodium chloride cell with a standard solucion of trans isomer (100 mg/ml in chloroform) and insert into the reference cell compartment of a suitable spectrophotometer. Add a chloroform solution of one of the standard cis-trans isomer mixtures to another 0.1 mm infrared sodium chloride cell and insert it into the sample cell compartment of the instrument. Record the infrared spectrum from 7.50 to 8.00 microns at least four times. Determine the absorbances of the maximum at 7.69 microns using the base line technique. Similarly, for each standard cis-trans isomer mixture, determine an average absorbance value from the replicate runs conducted. Prepare a standard reference curve by plotting the average absorbance values of the standard cis-trans isomer mixtures at 7.69 microns against the corresponding percentages of the cis isomers in these mixtures.

Procedure:

Dissolve 100 mg of the sample of Doxepin Hydrochloric in one ml of chloroform. Using the procedure used for the standard mixtures, determine the average absorbance value at 7.69 microns. Derive the cis isomer content in the sample from the standard curve. The percentage of cis doxepin in the sample is then calculated as follows:

$$\%\text{cis doxepin} = \%\text{cis doxepin hydrochloride} \times \frac{279.39}{315.85}$$ 1/

4.3.2.2 Ionic Chlorine content. Accurately weigh a 100 mg sample in a 250 ml beaker. Dissolve in 100 ml of purified water and 100 ml of U.S.P. Ethanol. Titrate the sample solution with standardized 0.05N Silver Nitrate solution and determine the equivalence point potentiometrically using a suitable potentiometer equipped with a calomel reference electrode, a silver electrode, and a 2N ammonium nitrate salt bridge. Calculate the percent ionic chlorine as follows:

$$\%\text{ionic chlorine} = \frac{V \times N \times 35.453 \times 100}{W}$$

Where:

V = ml of silver nitrate solution used at equivalence point.
N = normality of silver nitrate solution
W = weight of sample in mg

1/ As an alternate, an appropriate, accurate, reproducible gas chromatographic assay method may be employed.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

4.3.2.3 Occluded solvent.

Operating conditions:

Instrument: Aerograph Model 204 or its equivalent
Column: 6 feet by 1/8 inch Outside Diameter, stainless steel.
Column Packing: Porapak Q, 80-100 mesh
Detector: Flame ionization detector

Temperatures
Injector: 195°C.
Detector: 195°C.
Column Internal: 155°C.
Carrier Gas: Hydrogen-Oxygen
Flow Rate: Hydrogen 30 ml per minute
Nitrogen 35% at 60 p.s.i. (about 25 ml/minute).

Preparation of standard curves:

Transfer an accurately measured 5 µl aliquot of benzene, 5 µl aliquot of methylene chloride, and 10 µl aliquot of ethyl acetate to individual 100-ml glass-stoppered volumetric flasks containing about 90 ml of purified water. Immediately stopper the flasks and mix until the solvents are dissolved. Dilute to volume with purified water and mix again. Transfer 1, 2, 3, and 4 ml aliquots to separate No. 20 serum vials. Add sufficient purified water to bring the total volume in each vial to 10 ml and immediately cap the vials. Place the vials in a 70°C. constant temperature bath. After 30 minutes equilibrium period, withdraw 1 ml of the head space gas into a 1 ml gas-tight syringe and inject into the instrument. Record the curve. Repeat this process with each of the remaining three vials. Determine the half peak area obtained from each standard and construct a standard curve of concentration of solvent (mcg per ml) versus half-peak areas for each solvent, utilizing the density of each solvent to convert µl to mcg.

Procedure:

Transfer an accurately weighed 200 mg sample to a No. 20 serum vial containing 10 ml of purified water. Immediately cap the vial and shake until the sample is completely dissolved. Place the vial in a 70°C. water bath for 30 minutes, withdraw 1 ml of the head space gas into a 1 ml gas-tight syringe and inject into the instrument. Record the curve for the sample and determine the half peak area. Determine the amount of each solvent from the respective standard curves.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

Calculation:

$$\%\text{solvent} = \frac{\text{mcg of solvent (from standard curve)} \times 100}{\text{mcg of sample}}$$

4.3.2.4 Paper chromatography (identity).

Reagents. See 4.3.1.2

Chromatographic sheet. See 4.3.1.2

Sample Preparation.

Dissolve an accurately weighed 150 mg sample in sufficient chloroform:methanol (50:50 v/v) to make 10 ml of solution.

Reference Powder Preparation:

Prepare a reference solution as directed above for sample preparation, using Doxepin Hydrochloride reference powder.

Procedure:

Use the procedure specified in 4.3.1.2. The R_f values of the cis and trans isomers of the sample shall be identical to the R_f values of the cis and trans isomers of the reference powder.

4.3.2.5 Tertiary alcohol.

Reagents. See 4.3.1.2

Chromatographic sheet. See 4.3.1.2

Sample Preparation.

Dissolve an accurately weighed 200 mg sample in sufficient chloroform:methanol (50:50 v/v) to make 2 ml of solution.

Reference Sample Preparation:

Dissolve an accurately weighed 50 mg sample of 11-(3-dimethylamino-1-propyl) -6H-dibenz [b, e] oxepin-11-ol in sufficient chloroform: methanol (50:50 v/v) to make 100 ml of solution. The concentration of tertiary alcohol is approximately 0.5 mg per ml.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

Procedure:

Along the origin line of a prepared sheet, apply 0.050 ml of the sample preparation as a 3 inch long streak. Starting at the opposite side of the sheet, apply 0.050 ml of the reference sample solution as a 3 inch long streak so that it overlaps the sample streak for approximately one inch. Develop the sheet with the developing solvent using a descending technique for 20 +2 hours. After development, remove the sheet from the developing chamber, air dry for ten minutes, and finally dry the sheet at 60°C. for approximately 30 minutes in an air circulating oven. Spray the dried chromatogram with the idoplatinic acid reagent. Examine the sheet to see (1) if a band is present in the doxepin hydrochloride sample that forms a continuous line with the reference sample band, and (2) how the intensity of the band (if present) in the sample, compares with the intensity of the reference sample band.

Since the weight (0.05 mg) of the tertiary alcohol applied to the sheet corresponds to 0.5 percent of the weight (10 mg) of doxepin hydrochloride applied, estimate the abundance of tertiary alcohol as being (1) not detectable, (2) less than 0.5 percent, (3) approximately 0.5 percent, or (4) greater than 0.5 percent, by visually comparing the intensities of the two bands. For determining the total contaminants (3.3.7), score "less than 0.5 percent" and "approximately 0.5 percent" as "0.5 percent."

4.3.2.6 Ketone. Thin layer Examination.

Sample Preparation.

Dissolve 200 mg of doxepin hydrochloride in sufficient chloroform:methanol (50:50 v/v) to make 2 ml of solution.

Reference powder preparation.

Dissolve a 50 mg sample of 6,11-dihydrodibenz-[b, e] oxepin-11-one in sufficient chloroform methanol (50:50 v/v) to make 100 ml of solution. The concentration of this solution is 0.5 mg/ml.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

Preparation of Plate.

Mix one part of Silica Gel GF (Merck-Darmstadt) with two parts of purified water to form a slurry. Apply a uniform layer of the slurry over the surface of a six inch by eight inch glass plate. Air dry the plate for five to ten minutes and then activate at 100°C. for twenty minutes.

Procedure.

Apply 0.100 ml of the sample solution to a prepared plate as a three-inch streak to approximately one-half of the origin line. Apply 0.100 ml of the reference solution as a three inch streak to the other half of the origin line so that it overlaps approximately one inch of the sample streak. Develop the plate in a benzene-acetic acid (95:5 v/v) solvent system by an ascending technique until the solvent front has moved to within one inch of the top edge of the plate. Air dry the plate for 5 to 10 minutes and then examine the plate under a 366 nm ultraviolet light. The Doxepin Hydrochloride and 6,11-dihydrodibenz [b, e] oxepin-11-one appear as blue fluorescent bands on a dark background. Ascertain (1) whether a band is present in the doxepin hydrochloride bulk sample that forms a continuous line with the 6,11-dihydrodibenz [b, e] oxepin-11-one band and (2) how the intensity of this band (if present) compares to the intensity of the 6,11-dihydrobenz [b, e] oxepin-11-one band.

Since the weight (0.05 mg) of 6,11-dihydrodibenz [b, e] oxepin-11-one applied to the plate corresponds to 0.5 percent of the weight (10 mg) of the doxepin hydrochloride applied to the plate, estimate the abundance of 6,11-dihydrodibenz [b, e] oxepin-11-one (if present) as being (1) not detectable (2) less than 0.5 percent (3) approximately 0.5 percent, or (4) greater than 0.5 percent. For determining the total contaminants (3.2.7), score "less than 0.5 percent" and "approximately 0.5 percent" as "0.5 percent."

4.3.2.7 Ultraviolet absorption. Accurately weight a 50 mg sample of doxepin hydrochloride and transfer into a 100 ml volumetric flask. Dissolve the sample in ethanol and dilute to volume with ethanol. Transfer a 5 ml aliquot of the solution into a 50 ml volumetric flask and dilute to volume with ethanol. Record the ultraviolet absorption spectrum of the sample with a suitable spectrophotometer from 400 nm to 250 nm, using ethanol as a reference solvent. The spectrum of the sample shall exhibit a maximum and minimum absorbance at the same wavelengths as that of a similarly prepared solution of Doxepin Hydrochloride reference powder. In addition, the absorptivity of the sample at the maximum (296 nm) is not less than 96 percent and not more than 104 percent of the absorptivity of the reference sample.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

4.3.2.8 Infrared absorption. The infrared absorption spectrum of a 5 percent doxepin hydrochloride sample solution in chloroform shall compare qualitatively with that of a similar solution of doxepin hydrochloride reference powder when determined in 0.1 mm sodium chloride cells, using a suitable spectrophotometer with chloroform as a blank.

4.4 Packaging inspection. The inspection of the intermediate package, packing and marking for shipment and storage shall be in accordance with the quality assurance provisions of the applicable container specification and the marking requirements of MIL-STD-129.

4.5 Labeling inspection. Sampling shall be in accordance with MIL-STD-105 and the unit of product shall be the unit of issue. The inspection lot shall be the number of units of issue of each batch, lot, control number, etc., offered for inspection at one time. The inspection level for labeling shall be S-2 and the acceptance number shall be zero.

5. PACKAGING

5.1 Packaging requirements. The requirements of packaging shall be in accordance with all applicable requirements of PPP-C-186 (see 6.3) and as specified herein.

5.2 Packaging. Packaging shall be Level A or C, as specified (see 6.1).

5.2.1 Unit of issue. One bottle containing one hundred (100) capsules, as specified, constitutes one unit of issue.

5.2.2 Level A.

5.2.2.1 Immediate container. Immediate container shall be a commercially available immediate container considered acceptable by the Food and Drug Administration (FDA) for the product contained therein and shall be of such compatibility that neither the contents nor the container are altered in any way by each other.

5.2.2.2 Unit package. At the option of the contractor, each unit may be packaged as specified in 5.2.5 of PPP-C-186.

5.2.2.3 Intermediate package. Forty-eight (48) units of issue shall be packaged as specified in 5.2.8 of PPP-C-186.

5.2.3 Level C. Units shall be packaged in standard commercial containers of the size and kind commonly used, which will afford the degree of protection required for shipment and use of the product for its intended purpose.

5.3 Packing. Packing shall be Level A, B or C, as specified (see 6.1).

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

5.3.1 Level A. One hundred forty-four (144) units of issue shall be packed as specified in 5.4.4 of PPP-C-186.

5.3.2 Level B. One hundred forty-four (144) units of issue shall be packed as specified in 5.4.3.2 of PPP-C-186.

5.3.3 Level C. The subject commodity shall be packed in substantial commercial containers of the type, size, and kind commonly used for the purpose, so constructed as to insure acceptance and safe delivery by common or other carriers, at the lowest rate, to point of delivery called for in the contract or purchase order.

5.4 Marking.

5.4.1 Labeling. Labeling shall be in accordance with the requirements of the Federal Food, Drug and Cosmetic Act, and shall include the information required below.

5.4.1.1 Immediate container. Each immediate container label shall bear the following information:

(a) label information in accordance with commercial practice.
(b) the expiration date.

5.4.1.2 Unit package. Each unit package label shall bear the following information:

(a) label information in accordance with commercial practice.
(b) the national stock number.

NOTES: (1) The national stock number shall not be preceded by any prefix.
(2) Strip labeling may be used.
(3) Label adhesion test shall not apply.
(4) If a unit package is not supplied, the national stock number shall appear on the immediate container label.

5.4.2 Intermediate package. Each intermediate package shall be marked as specified in 5.5.3 of PPP-C-186. Type II shelf life markings shall be shown as specified in MIL-STD-129.

5.4.3 Exterior container. Exterior container shall be marked as specified in 5.5.4 of PPP-C-186. Type II shelf life marking shall be shown as specified in MIL-STD-129.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

6. NOTES

6.1 Ordering data. Procurement documents should specify the following:

(a) Title, number and date of this specification.
(b) the National Stock Number.
(c) Selection of applicable levels of packaging and packing (See 5.2 and 5.3).

6.2 This specification covers the following item(s):

National Stock No.	Item Identification
6505-00-408-8935	DOXEPIN HYDROCHLORIDE CAPSULES, Equivalent to 25 mg of Doxepin, 100s

6.3 All references to PPP-C-186 refer to Interim Federal Specification PPP-C-00186B which is mandatory for use until superseded by a revision.

Custodians:
Army - MD
Navy - MS
Air Force - 03

Preparing activity:
Defense Supply Agency - DM

Project No. 6505-1501

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

STANDARDIZATION DOCUMENT IMPROVEMENT PROPOSAL

OMB Approval
No. 22-R255

INSTRUCTIONS: The purpose of this form is to solicit beneficial comments which will help achieve procurement of suitable products at reasonable cost and minimum delay, or will otherwise enhance use of the document. DoD contractors, government activities, or manufacturers/ vendors who are prospective suppliers of the product are invited to submit comments to the government. Fold on lines on reverse side, staple in corner, and send to preparing activity. Comments submitted on this form do not constitute or imply authorization to waive any portion of the referenced document(s) or to amend contractual requirements. Attach any pertinent data which may be of use in improving this document. If there are additional papers, attach to form and place both in an envelope addressed to preparing activity.

DOCUMENT IDENTIFIER AND TITLE

MIL-D-37356 DOXEPIN HYDROCHLORIDE CAPSULES

NAME OF ORGANIZATION AND ADDRESS

CONTRACT NUMBER

MATERIAL PROCURED UNDER A

☐ DIRECT GOVERNMENT CONTRACT ☐ SUBCONTRACT

1. HAS ANY PART OF THE DOCUMENT CREATED PROBLEMS OR REQUIRED INTERPRETATION IN PROCUREMENT USE?

A. GIVE PARAGRAPH NUMBER AND WORDING.

B. RECOMMENDATIONS FOR CORRECTING THE DEFICIENCIES

2. COMMENTS ON ANY DOCUMENT REQUIREMENT CONSIDERED TOO RIGID

3. IS THE DOCUMENT RESTRICTIVE?

☐ YES ☐ NO *(If "Yes", in what way?)*

4. REMARKS

SUBMITTED BY *(Printed or typed name and address - Optional)*

TELEPHONE NO.

DATE

DD FORM 1 JAN 72 1426

REPLACES EDITION OF 1 JAN 66 WHICH MAY BE USED.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.

FOLD

DEFENSE PERSONNEL SUPPORT CENTER
2800 South 20th Street
Philadelphia, PA 19101

POSTAGE AND FEES PAID
DEFENSE SUPPLY AGENCY
DoD • 304

U.S.MAIL

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE $300

HEADQUARTERS, DEFENSE PERSONNEL SUPPORT CENTER
ATTN: DIRECTORATE OF MEDICAL MATERIEL, CODE ATT
2800 SOUTH 20TH STREET
PHILADELPHIA, PA 19101

FOLD

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T10:31Z
Check the source to verify that this is the current version before use.