MILITARY SPECIFICATION

OXYCODONE HYDROCHLORIDE, ASPIRIN, CAFFEINE, OXYCODONE TEREPHTHALATE, AND PHENACETIN TABLETS.

This specification is approved for use by the Defense Personnel Support Center, Defense Logistics Agency, and is available for use by all Departments and Agencies of the Department of Defense.

1. SCOPE

1.1 Scope. This specification covers Oxycodone Hydrochloride, Aspirin, Caffeine, Oxycodone Terephthalate, and Phenacetin Tablets.

1.2 Classification. The tablets covered by this specification shall be of the following sizes (See 6.2):

Size 1 - Individually Sealed, 250s
Size 2 - 100s

2. APPLICABLE DOCUMENTS

2.1 Issues of documents. The following documents, of the issue in effect on date of invitation for bids or request for proposals, form a part of this specification to the extent specified herein.

SPECIFICATIONS

FEDERAL

PPP-C-186 - Containers, Packaging and Packing for Drugs, Chemicals and Pharmaceuticals.

Beneficial comments (recommendations, additions, deletions) and any pertinent data which may be of use in improving this document should be addressed to: Headquarters, Defense Personnel Support Center, ATTN: Directorate of Medical Materiel, DPSC-ATT, 2800 South 20th Street, Philadelphia, Pa. 19101, by using the self-addressed Standardization Document Improvement Proposal (DD Form 1426) appearing at the end of this document or by letter.

FSC 6505

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T11:23Z
Check the source to verify that this is the current version before use.

STANDARDS

FEDERAL

FED-STD-140 - Tablets (for Medicinal Purposes).

MILITARY

MIL-STD-105 - Sampling Procedures and Tables for Inspection by Attributes.
MIL-STD-129 - Marking for Shipment and Storage.

(Copies of specifications and standards required by contractors in connection with specific procurement functions should be obtained from the procuring activity or as directed by the contracting officer.)

2.2 Other publications. The following documents form a part of this specification to the extent specified herein. Unless otherwise indicated, the issue in effect on date of invitation for bids or request for proposals shall apply.

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE,
FOOD AND DRUG ADMINISTRATION (FDA)

Federal Food, Drug, and Cosmetic Act and Regulations Promulgated Thereunder.

(Application for copies should be addressed to the Food and Drug Administration, Washington, DC 20204.)

U. S. PHARMACOPEIAL CONVENTION, INC.

The United States Pharmacopeia (USP).

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

The National Formulary (NF).

(Application for copies should be addressed to the Mack Publishing Company, Easton, PA 18042.)

Source: https://assist.dla.mil -- Downloaded: 202610-21T11:23Z
Check the source to verify that this is the current version before use.

3. REQUIREMENTS

3.1 Material. Shall be Oxycodone Hydrochloride, Aspirin, Caffeine, Oxycodone Terephthalate and Phenacetin Tablets and shall be in accordance with all applicable requirements of FED-STD-140, and as specified herein.

S2. Classification. Shall be type I.

S5.3 The following is added to this paragraph:

The tablets shall contain 4.50 mg of Oxycodone Hydrochloride, 0.38 mg Oxycodone Terephthalate, 224 mg Aspirin, 160 mg Phenacetin and 32 mg Caffeine within the designated assay limits for the tablets.

The tablets shall assay to contain not less than 90.0 percent and not more than 110.0 percent of the required amounts of Oxycodone Hydrochloride, Aspirin, Caffeine Anhydrous, Oxycodone Terephthalate and Phenacetin when assayed as specified in 4.2.2. During the expiration dating period the Salicylic acid content shall not exceed 1.0 percent of the Aspirin content when determined as specified in 4.2.3.

S6.4.9.1 Disintegration. Shall disintegrate in not more than 5 minutes when determined by the USP method for Uncoated Tablets.

3.2 Ingredients.

3.2.1 Oxycodone Hydrochloride.

3.2.1.1 Identification. Shall comply with the following:

3.2.1.1.1 Infrared absorption. The infrared absorption spectrum of a potassium bromide dispersion of the oxycodone hydrochloride, previously dried at 105° C for 2 hours, shall exhibit maxima only at the same wavelengths as that of a similar preparation of oxycodone hydrochloride reference standard. Standard Reference spectrum shall be made available to the Government upon request.

3.2.1.1.2 The ultraviolet absorption spectrum of a 1 in 6500 solution in 0.1 N HCl shall exhibit maxima and a minimum at the same wavelengths as that of a similarly prepared solution of oxycodone hydrochloride reference standard, concomitantly measured. Standard Reference spectrum shall be made available to the Government upon request.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T11:23Z
Check the source to verify that this is the current version before use.

3.2.1.1.3 Thin layer chromatograph. The sample shall show a dark blue spot at approximately the same Rf value as that of the reference standard (about 0.60) when determined as follows: Dissolve 100 mg of oxycodone hydrochloride sample in 5.0 ml of methanol. Similarly dissolve 90 mg of Oxycodone Base Reference Standard in 5.0 ml of chloroform. Plate shall be approximately 20 x 20 cm in size with a uniform 250 micron silica gel layer previously heated at 105° C for 15 minutes. Spot 5 ul of each solution. Immediately place the plate into a Gelman Rapid Development Chromatographic Chamber, or equivalent unit, lined with saturation pads and containing the solvent, ammonia-washed chloroform:ethanol (85:15). (Note: Prepare the chloroform by shaking 100 ml of chloroform with 50 ml of 3 percent aqueous ammonia in a separator.) Tape the glass cover of the chamber to prevent evaporation of the solvent, cover the chamber with a towel or other means to eliminate light, and allow the chamber to remain in a horizontal position for 15 minutes, Carefully stand the chamber up and develop the chromatograph for a distance of 10 cm. Remove the plate and dry. Spray with freshly prepared iodoplatinate reagent (consisting of equal volumes of 6 percent aqueous KI and 0.3 percent aqueous platinum chloride). Compare sample and standard spots for location, size, and intensity.

3.2.1.2 Specific rotation. Shall be between -137° and -149° when determined in an aqueous solution containing 250 mg in each 10 ml calculated on a solvent-free basis.

3.2.1.3 Acidity. Not more than 0.3 ml of 0.02 N NaOH shall be required to produce a yellow color when 300 mg of oxycodone hydrochloride are dissolved in 10 ml of purified water and 1 drop of methyl red TS added.

3.2.1.4 Water. Shall contain no more than 5 percent water when determined by the USP Karl Fischer electrometric method using 1 gram of sample.

3.2.1.5 Methanol and Ethanol. Shall contain no more than 3 percent total of methanol and ethanol when determined by an appropriate, accurate, reproducible gas chromatographic method.

3.2.1.6 Residue on ignition. Shall not exceed 0.5 mg using a sample of 200 mg when determined by the USP method.

3.2.1.7 Chloride. Shall contain not less than 9.8 percent and not more than 10.4 percent when approximately 300 mg of sample is accurately weighed into 50 ml of methanol, 5 ml of glacial acetic acid is added, plus 2 drops of Eosin Y T.S., and the solution is titrated with 0.1 N silver nitrate volumetric solution. Each ml of 0.1 N silver nitrate solution is equivalent to 3.546 mg of chloride.

Source: https://assist.dla.mil -- Downloaded: 202610-21T11:23Z
Check the source to verify that this is the current version before use.

3.2.1.8 14-Hydroxycodeinone Impurity. Shall contain a maximum of 1.0 percent when determined using an appropriate, accurate, reproducible gas chromatographic method.

3.2.1.9 Assay.

3.2.1.9.1 Non-aqueous Titration. Shall contain not less than 97.0 and not more than 103.0 percent of oxycodone hydrochloride when determined as follows: Dissolve approximately 700 mg of oxycodone hydrochloride, accurately weighed, in 50 ml of glacial acetic acid. Add 10 ml of 6 percent mercuric acetate solution in glacial acetic acid, followed by 1 drop of methyl violet, TS. Titrate with 0.1 N perchloric acid volumetric solution in glacial acetic acid to a sky blue color. Perform a blank determination and make any necessary correction. Each ml of $HClO_4$ is equivalent to 35.18 mg of oxycodone hydrochloride.

3.2.1.9.2 Gas chromatography. Shall contain not less than 97.0 and not more than 103.0 percent of oxycodone hydrochloride when determined by an appropriate, accurate, and reproducible gas chromatographic method.

3.2.2 Oxycodone Terephthalate.

3.2.2.1 Identification. Shall comply with the following:

3.2.2.1.1 The ultraviolet spectrum of the filtrate from the assay for Terephthalic acid diluted to 100.0 ml with purified water and further diluted 2:100, exhibits a maximum and minimum at the same wavelengths as the standard reference spectrum on file with the company. Standard Reference spectrum shall be made available to the Government, upon request.

3.2.2.1.2 Precipitate of dihydroxycodeinone shall melt between 217 and 221° C when prepared as follows: Transfer 50 ml of the filtrate from the terephthalic acid assay to a flask, add sufficient ammonia T.S. to render the solution alkaline and permit to stand 2 hours. Collect the ppt on a filter, wash with cold purified water, and dry at 105° C for 2 hours.

3.2.2.2 Loss on drying. Shall lose no more than 1.5 percent when dried at 105° C for 2 hours.

3.2.2.3 Residue on ignition. Shall not exceed 1.0 percent when determined by the USP method using 1 gram of sample.

3.2.2.4 Assay for Terephthalic Acid. Shall be not less than 19.8 percent and not more than 21.8 percent calculated on a dry basis when determined as follows: Accurately weigh 1 gram of sample into a beaker, add 25 ml of

Source: https://assist.dla.mil -- Downloaded: 202610-21T11:23Z
Check the source to verify that this is the currentversion before use.

0.2 N HCl and heat nearly to boiling, while stirring to break up flocks. Cover and cool to room temperature. Filter through a tared medium porosity filtering crucible, completing transfer with several small volumes of 0.2 N HCl. Collect filtrate and washings in a 100 ml volumetric flask. Wash with several small portions of 0.2 N HCl. Dilute the combined filtrate to 100.0 ml with purified water and reserve for subsequent assays of dihydrohydroxycodeinone. Dry the ppt at 105° C for 1 hour, cool to room temperature in a desiccator, and weigh:

3.2.2.5 Assay for Dihydrohydroxycodeinone. Shall contain not less than 78.5 percent and not more than 80.5 percent when assayed as follows: Transfer a 2.0 ml aliquot of the final 100 ml filtrate from the assay (see 3.2.2.4) into a second 100 ml volumetric flask and dilute to volume with purified water. Read the absorbance at 281.5 nm in a 1 cm cell with a suitable spectrophotometer using purified water as the reference.

Calculation:

$$\frac{Au \times 12.3}{\text{wht of sample (Grams)}} = \text{percent DHHC}$$

where Au = absorbance of sample

3.2.2.6 Assay for Dihydrohydroxycodeinone Terephthalate. Shall assay to contain not less than 97.0 and not more than 103.0 percent of oxycodone terephthalate calculated on a dry basis when determined as follows: Accurately weigh approximately 500 mg of oxycodone terephthalate into a 250 ml flask, add 50 ml of glacial acetic acid, 10 ml of mercuric acetate TS and 2 drops of crystal violet TS. Titrate with 0.1 N perchloric acid volumetric solution in dioxane, to a sky blue color. Perform a blank determination and make any necessary correction. Each ml of $HClO_4$ is equivalent to 39.84 mg of oxycodone terephthalate.

3.2.3 Aspirin, Phenacetin, Caffeine (Anhydrous). The Aspirin, Phenacetin (USP XVIII) and anhydrous Caffeine used in the manufacture of the tablets shall comply with the tests, standards, and requirements of the USP, including any supplements or revisions thereto.

3.3 Other ingredients. All other ingredients used in the manufacture of the tablets shall comply with S5.1 of FED-STD-140.

3.4 Expiration dating period. The expiration dating period shall be not less than 36 months.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T11:23Z
Check the source to verify that this is the current version before use.

4. QUALITY ASSURANCE PROVISIONS

4.1 Supplier responsibility for inspection. Unless otherwise specified in the contract or purchase order, the supplier is responsible for the performance of all inspection requirements as specified herein. Except as otherwise specified in the contract or order, the supplier may use his own or any other facilities suitable for the performance of the inspection requirements specified herein, unless disapproved by the Government. The Government reserves the right to perform any of the inspections set forth in the specification where such inspections are deemed necessary to assure supplies and services conform to prescribed requirements.

4.1.1 Records of examinations and tests performed by or for the contractor shall be maintained by the contractor and made available to the Government, upon the Government's request, at any time, or from time to time, during the performance of the contract and, (i) as to any expiration dated item, for the expiration dating period specified by the contractor for such item, or such longer period as may be required by regulation of any Federal agency; or (ii) as to non-dated items, for not less than 3 years after delivery of the item to the Government.

4.1.2 No company supplying any ingredient(s) to the contractor will be considered an acceptable facility for the performance of any inspection requirements specified herein.

4.2 Assay for finished tablets.

4.2.1 Oxycodone Hydrochloride Anhydrous. (Note: Oxycodone Terephethalate content is included here but reported as Oxycodone Hydrochloride equivalent.)

Standard solution.

Prepare a solution with a known final concentration of approximately 0.135 mg of Oxycodone Base Standard per ml, in chloroform.

Procedure.

Place 20 tablets in a glass-stoppered flask, add 100.0 ml of approximately 0.01 N HCl. Stopper and shake for 15 minutes. Filter through suitable filter paper rejecting the first 10 ml. Transfer 15.0 ml of the filtrate to a separatory funnel and extract with 4 x 40 ml of chloroform, discarding

Source: https://assist.dla.mil -- Downloaded: 202610-21T11:23Z
Check the source to verify that this is the current version before use.

the chloroform. Carefully adjust the aqueous solution to a pH between 9 and 10 with 10 percent ammonia, and extract with 3 x 25 ml chloroform running each chloroform extract through a cotton pledget wetted with chloroform into a 100 ml volumetric flask. Dilute to volume with chloroform.

Determine the absorbances of sample and standard in a 1 cm cell at 283.5 nm with a suitable spectrophotometer using chloroform as the blank.

Calculation:

$$\frac{Au}{As} \times \frac{\text{mg standard} \times 1.116 \times \text{purity std}}{30} = \text{mg Oxycodone Hydrochloride per tablet}$$

where Au = Absorbance sample
As = Absorbance standard

4.2.2 Aspirin, Phenacetin, and Caffeine Anhydrous.

Standard Preparation.

Working Standard Solution: Accurately weigh a mixture of USP Reference Standards in the amounts indicated below:

896 mg of aspirin
640 mg of phenacetin
128 mg of caffeine anhydrous

Add exactly 50.0 ml of chloroform and bring into solution.

Assay Preparation.

Sample Assay Solution: Weigh 20 tablets, determine the average tablet weight, and finely powder. Accurately weigh the equivalent of four tablets, and transfer quantitatively to a 125 ml glass-stoppered flask. Add exactly 50.0 ml of chloroform, stopper, and shake for 15 minutes. Filter through suitable filter paper.

Procedure.

Determine the infrared spectra of the sample assay and working standard solution, using 0.1 mm matched cells with chloroform as the reference.

Source: https://assist.dla.mil -- Downloaded: 202610-21T11:23Z
Check the source to verify that this is the current version before use.

Absorption peaks are approximately:

5.64 u - aspirin
6.60 u - phenacetin
6.02 u - caffeine

Readings should be made at wavelengths of maximum absorption provided peaks remain within experimental limits.

NOTE: Since aspirin and phenacetin have considerable absorbance at the wavelength at which caffeine is read, corrections must be made for these two components. Since it may be cumbersome to calculate the caffeine values manually, they may be computerized.

Tabulate the absorbance values for APC for all samples and standards. Using this data, plus the Average Tablet Weights, Sample Weights, Standard Weights, and their purities, calculate the Aspirin, Phenacetin and Caffeine for each sample.

Calculations:

$$\frac{\text{Abs. Sample}}{\text{abs. Std.}} \times \frac{\text{std. wt.}}{\text{sample wt.}} \times \text{avg. tab. wt.} \times \text{std. purity} = \text{mg Aspirin per tablet}$$

$$\frac{\text{Abs. sample}}{\text{abs. Std.}} \times \frac{\text{std. wt.}}{\text{sample wt.}} \times \text{avg. tab. wt.} \times \text{std. purity} = \text{mg Phenacetin per tablet}$$

Caffeine:

C' = Observed sample caffeine abs. - 0.056a - 0.110b
C" = Observed standard caffeine abs. - 0.056c - 0.110d 1/

$$\frac{C'}{C''} \times \frac{\text{std. wt.}}{\text{sample wt.}} \times \text{avg. tab. wt.} \times \text{std. purity} = \text{mg Caffeine per tablet}$$

WHERE:

abs = absorbance
C' = corrected sample caffeine absorbance
C" = corrected standard caffeine absorbance
a = Total mg aspirin in sample / 896
b = Total mg phenacetin in sample / 640
c = (mg aspirin std weighed out) (aspirin std purity) / 896
d = (mg phenacetin std weighed out) (phenacetin std purity) / 640

NOTES: I. Aspirin and phenacetin sample values must be calculated first before sample caffeine values can be calculated.
II. Correction factors are based on 50 ml of chloroform.
1/ III. Correction factors used in the caffeine calculation shall be determined for the particular instrument under specified operating conditions.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T11:23Z
Check the source to verify that this is the current version before use.

4.2.3 Free Salicylic Acid.

Reagents Preparation:

Ferric nitrate reagent: 1 gram of ferric nitrate and 1 ml of concentrated nitric acid in 100 ml of purified water.

Stock Standard Solution: Weigh accurately 100 mg of USP Salicylic Acid Reference Standard into a 100 ml volumetric flask. Dilute to volume with 3A alcohol.

Assay Preparation:

Sample Assay Solution: Transfer the equivalent of 5 tablets finely powdered to a 100 ml volumetric flask, and add approximately 50 ml of 3A alcohol. Shake the mixture for 15 minutes, and dilute to volume with 3A alcohol. Filter through a No. 1 Whatman filter paper, or equivalent.

Blank Solution: Transfer 15.0 ml of 3A alcohol and 5.0 ml of ferric nitrate reagent to a 100 ml volumetric flask, and dilute to volume with purified water.

Procedure:

Transfer a 2.0 aliquot of the stock standard solution to a 100 ml volumetric flask and add 13.0 ml of 3A alcohol and 5.0 ml of ferric nitrate reagent. Dilute to volume with purified water. Read on a suitable colorimeter or spectrophotometer at 525 nm, using the blank as reference.

Transfer a 15.0 ml aliquot of the sample assay solution to a 100 ml volumetric flask, add 5.0 ml of ferric nitrate reagent, and dilute to volume with purified water. If the solution is cloudy, filter through a millipore filter (solvinert type). Read the solution on a suitable colorimeter or spectrophotometer at 525 nm, as soon as possible after addition of ferric nitrate reagent, using the blank as reference.

Calculation:

$$\frac{A \text{ sample}}{A \text{ std.}} \times \text{conc stock std (mg/100 ml)} \times \frac{\text{avg tab wt (gram)}}{\text{sample wt (gram)}} \times 6.67 \times 0.02 =$$

mg free Salicylic acid/tablet

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T11:23Z
Check the source to verify that this is the current version before use.

5. PACKAGING

5.1 Packaging requirements. The requirements for packaging shall be in accordance with all applicable requirements of PPP-C-186 and as specified herein.

5.2 Packaging. Packaging shall be level A or C as specified (See 6.1).

5.2.1 Unit of issue. One bottle containing 100 tablets as specified; or one box containing 10 packages, each package containing 25 tablets individually sealed, as specified, constitutes one unit of issue.

5.2.2 Level A.

5.2.2.1 Immediate container.

5.2.2.1.1 Size 1. Size 1 shall be supplied, individually sealed, in a commercially available reverse-numbered blister or strip packaging considered acceptable to the Food and Drug Administration (FDA) for the product contained therein and shall be of such compatibility that neither the contents nor the container are altered in any way by each other.

5.2.2.1.2 Size 2. The immediate container for size 2 shall be a commercially available bottle considered acceptable by the Food and Drug Administration (FDA) for the product contained therein and shall be of such compatibility that neither the contents nor the container are altered in any way by each other.

5.2.3 Unit package. For size 1, each unit shall be packaged as specified in 5.2.5 of PPP-C-186. For size 2, at the option of the contractor, each unit may be packaged as specified in 5.2.5 of PPP-C-186.

5.2.4 Level C. Units shall be packaged in standard commercial containers of the size and kind commonly used, which will afford the degree of protection required for shipment and use of the product for its intended purpose.

5.3 Packing. Packing shall be level A, B or C, as specified (see 6.1).

5.3.1 Level A. For size 1, 4 units of issue shall be packed as specified in 5.4.4 of PPP-C-186. For size 2, 48 units of issue shall be packed as specified in 5.4.4 of PPP-C-186.

5.3.2 Level B. For size 1, 4 units of issue shall be packed as specified in 5.4.3.4 of PPP-C-186. For size 2, 48 units of issue shall be packed as specified in 5.4.3.2 of PPP-C-186.

Source: https://assist.dla.mil -- Downloaded: 202610-21T11:23Z
Check the source to verify that this is the currentversion before use.

5.3.3 Level C. The subject commodity shall be packed in substantial commercial containers of the type, size, and kind commonly used for the purpose, so constructed as to insure acceptance and safe delivery by common or other carriers, at the lowest rate, to point of delivery called for in the contract or purchase order.

5.4 Marking.

5.4.1 Labeling. Labeling shall be in accordance with the requirements of the Federal Food, Drug and Cosmetic Act and shall include the information required below:

5.4.1.1 Size 1.

5.4.1.1 Immediate container. Each immediate container label for size 1 shall bear the following information:

(a) label information in accordance with commercial practice.

(b) the expiration date (not required if supplied with unit package).

5.4.1.1.2 Unit package. Each unit package label for size 1 shall bear the following information:

(a) label information in accordance with commercial practice.

(b) the National Stock Number.

(c) the expiration date (not required if supplied with immediate container).

5.4.1.2 Size 2.

5.4.1.2.1 Immediate container. Each immediate container label for size 2 shall bear the following information:

(a) label information in accordance with commercial practice.

(b) the expiration date.

5.4.1.2.2 Unit package. Each unit package label for size 2 shall bear the following information:

(a) label information in accordance with commercial practice.

(b) the National Stock Number.

Source: https://assist.dla.mil -- Downloaded: 202610-21T11:23Z
Check the source to verify that this is the current version before use.

NOTES: (1) The National Stock Number shall not be preceded by any prefix.

(2) Strip labeling may be used.

(3) Label adhesion test shall not apply.

(4) If a unit package is not supplied, the National Stock Number shall appear on the immediate container label.

5.4.3 Exterior container. Exterior container shall be marked as specified in 5.5.4 of PPP-C-186. Type I shelf life marking shall be shown as specified in MIL-STD-129. Marking shall include the legend: "STORE AT CONTROLLED ROOM TEMPERATURE 15° - 30° C (59° - 86° F)".

6. NOTES

6.1 Ordering data. Procurement documents should specify the following:

(a) Title, number, and date of this specification.

(b) National Stock Number.

(c) Selection of applicable levels of packaging and packing (See 5.2 and 5.3).

6.2 This specification covers the following item(s):

National Stock Number	Item Identification	Specification Designation
6505-01-030-9492	OXYCODONE HYDROCHLORIDE, ASPIRIN, CAFFEINE, OXYCODONE TEREPHTHALATE, AND PHENACETIN TABLETS, Individually Sealed, 250s	Size 1
6505-01-030-9493	OXYCODONE HYDROCHLORIDE, ASPIRIN, CAFFEINE, OXYCODONE TEREPHTHALATE, AND PHENACETIN TABLETS, 100s	Size 2

Preparing activity:
DLA-DM

Project No. 6505-1875

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T11:23Z
Check the source to verify that this is the current version before use.

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T11:23Z
Check the source to verify that this is the current version before use.

FOLD

DEFENSE PERSONNEL SUPPORT CENTER
2800 South 20th Street
Philadelphia, PA 19101

POSTAGE AND FEES PAID
DEFENSE SUPPLY AGENCY
DoD - 304

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE $300

HEADQUARTERS, DEFENSE PERSONNEL SUPPORT CENTER
ATTN: DIRECTORATE OF MEDICAL MATERIEL, CODE ATT
2800 SOUTH 20TH STREET
PHILADELPHIA, PA 19101

FOLD

Source: https://assist.dla.mil -- Downloaded: 2026-10-21T11:23Z
Check the source to verify that this is the current version before use.

STANDARDIZATION DOCUMENT IMPROVEMENT PROPOSAL

OMB Approval
No. 22-R255

INSTRUCTIONS: The purpose of this form is to solicit beneficial comments which will help achieve procurement of suitable products at reasonable cost and minimum delay, or will otherwise enhance use of the document. DoD contractors, government activities, or manufacturers/vendors who are prospective suppliers of the product are invited to submit comments to the government. Fold on lines on reverse side, staple in corner, and send to preparing activity. Comments submitted on this form do not constitute or imply authorization to waive any portion of the referenced document(s) or to amend contractual requirements. Attach any pertinent data which may be of use in improving this document. If there are additional papers, attach to form and place both in an envelope addressed to preparing activity.

DOCUMENT IDENTIFIER AND TITLE: MIL-O-37733 (DLA-DM) OXYCODONE HYDROCHLORIDE, ASPIRIN, CAFFEINE, OXYCODONE TEREPHTHALATE, AND PHENACETIN TABLETS

NAME OF ORGANIZATION AND ADDRESS

CONTRACT NUMBER

MATERIAL PROCURED UNDER A

☐ DIRECT GOVERNMENT CONTRACT ☐ SUBCONTRACT

1. HAS ANY PART OF THE DOCUMENT CREATED PROBLEMS OR REQUIRED INTERPRETATION IN PROCUREMENT USE?
   A. GIVE PARAGRAPH NUMBER AND WORDING.

   B. RECOMMENDATIONS FOR CORRECTING THE DEFICIENCIES

2. COMMENTS ON ANY DOCUMENT REQUIREMENT CONSIDERED TOO RIGID

3. IS THE DOCUMENT RESTRICTIVE?

   ☐ YES ☐ NO *(If "Yes", in what way?)*

4. REMARKS

SUBMITTED BY *(Printed or typed name and address - Optional)*

TELEPHONE NO.

DATE

DD FORM 1 JAN 72 1426 REPLACES EDITION OF 1 JAN 66 WHICH MAY BE USED

Source: https://assist.dla.mil -- Downloaded: 202610-21T11:23Z
Check the source to verify that this is the current version before use.