INCH-POUND

A-A-197B
7 January 2020
SUPERSEDING
A-A-197A
16 September 2014

COMMERCIAL ITEM DESCRIPTION

HERBICIDE, 2,4-DICHLOROPHENOXYACETIC ACID (SALTS AND ESTERS)

The General Service Administration has authorized the use of this commercial item description for all federal agencies.

1. SCOPE. This commercial item description covers the systemic herbicide 2,4-dichlorophenoxyacetic acid (2,4-D).

2. CLASSIFICATION. The herbicide shall be of the following types and classes:

2.1 Type II. Liquid salt formulation

2.2 Type III. Liquid ester formulation

Class 1 - Volatile alkyl esters of 2,4-D
Class 2 - Low volatile esters of 2,4-D

3. SALIENT CHARACTERISTICS.

3.1 Liquid salts. The liquid salts of 2,4-D shall have the following characteristics.

3.1.1 Acid equivalent. The product shall contain a minimum of two pounds of 2,4-D equivalent per gallon of formulation at 68°F (20°C).

3.1.1.1 Acid equivalent determination. The quantity of 2,4-D contained in the salt formulation shall be determined by Official Methods of Analysis of AOAC INTERNATIONAL (OMA) method 978.05, “2,4-D in Pesticide Formulations”.

3.1.2 Solubility. The product shall be soluble in hard or soft water and shall exhibit no persistent foam.

Beneficial comments, recommendations, additions, deletions, clarifications, etc. and any data that may improve this document should be sent to DLA Aviation, VEB, 8000 Jefferson Davis Highway, Richmond, VA 23297-5616 or e-mailed to STDZNMGT@dla.mil. Since contact information can change, you may want to verify the currency of this address information using the ASSIST database at https://assist.daps.dla.mil

AMSC N/A FSC 6840

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T08:05Z
Check the source to verify that this is the current version before use.

3.1.3 Other ingredients. The product shall make a solution that contains no ingredients which will inhibit the application of the material at the concentrations normally used for weed control. The product shall remain free of solid material when held at a temperature of 25°F (-3.8°C) for a period of at least 5 consecutive 24 hour days.

3.2 Liquid esters. The liquid esters of 2,4-D shall have the following characteristics.

3.2.1 Acid equivalent. The product shall contain a minimum of two pounds of 2,4-D equivalent per gallon of formulation at 68°F (20°C).

3.2.1.1 Acid equivalent determination. The quantity of 2,4-D contained in the ester formulation shall be determined by OMA method 978.05.

3.2.2 Miscibility and emulsifibility. The formulation shall be a clear solution readily miscible with oil and emulsifiable with water. It shall contain the necessary solvents and carrying and emulsifying agents, such that the emulsion formed requires a minimum of agitation during the mixing and application period.

3.2.3 Other ingredients. The oil carrier in the formulation shall not detract from the herbicidal activity of the active ingredient and offer maximum penetration and spread of the spray solution. The product shall remain free of solid material when held at a temperature of 25°F (-3.8°C) for a period of at least 5 consecutive 24 hour days.

3.2.4 Volatility. Class 2 esters include glycol, poly-glycol, and their ether ester derivatives of 2,4-D as well as other heavy molecular weight esters of 2,4-D that are known to have low volatility. Class 2 esters shall not include esters of the lower alkyl groups such as methyl, ethyl, propyl, isopropyl, butyl, or pentyl groups. When tested according to 3.2.4.1, Class 2 liquid esters shall have an average vapor activity rating of less than 4.0.

3.2.4.1 Volatility test - relative vapor activity. Vapor activity shall be determined using gas tight cases approximately 3.5 x 3.5 x 16 inches in size. Young rapidly growing Pinto bean plants about 4 inches in height are used as test plants. A single bean plant growing in a 3 inch pot is placed in each cellophane case just prior to testing the ester. Two milligrams of acid equivalent as the ester is dissolved in 10 milliliters of 95 percent ethyl alcohol and a Whatman No. 1 filter paper (9 cm. diameter) is thoroughly moistened by dipping in the solution. (Do not reuse the container used in this impregnation.) The alcohol is then allowed to evaporate and the filter paper impregnated with the ester is inserted into the cellophane case containing the bean plant and fastened to the inside of the case 6 inches above the leave of the test plant. The open end of the case is then sealed. The case containing the test plant and treated filter paper is then placed in a dark room for a period of 24 hours. The temperature range of the room should be 80° to 90° F. Control plants are also sealed in separate cases. The experimental design is a randomized block with three replications and each test is repeated three times. The evaluations shall be made following an exposure period of 24 hours. Observation of the effect of the vapors on test plants should take into consideration whether or not the plant is slightly, moderately, or severely injured, including such symptoms as degree of stem curvature, terminal bud inhibition, and

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T08:05Z
Check the source to verify that this is the current version before use.

degree of leaf curl. The relative vapor activity of an ester can be numerically designated as follows: 0 - no visible effect; 1,2,3 - slight injury - plant usually recovered with little or no reduction in growth, slight epinasty present, stem curvature slight; 4,5,6 - moderate injury - plant usually recovered, moderate epinasty, moderate terminal bud inhibition and moderate stem curvature present; 7,8,9 – severe injury - plant usually does not recover, pronounced epinasty, together with pronounced stem curvature; 10 - plant killed. The 2,4-D acid under most conditions is rated 0 while the butyl ester has a high vapor activity with a rating of 9.0.

4. REGULATORY REQUIREMENTS.

4.1 Registration and labeling. The herbicide shall be registered and labeled in compliance with the Regulations for the Enforcement of the Federal Insecticide Fungicide and Rodenticide Act (40 CFR 162-180).

4.2 Recovered material. The offeror/contractor is encouraged to use recovered materials to the maximum extent practicable, in accordance with paragraph 23.403 of the Federal Acquisition Regulation (FAR).

4.3 Safety data sheet (SDS). Manufacturers shall prepare and submit a SDS in accordance with FED-STD-313, "Material Safety Data, Transportation Data and Disposal Data for Hazardous Materials Furnished to Government Activities", and meeting the requirements of 29 Code of Federal Regulations § 1910.1200, "Occupational Safety and Health Standards: Hazard Communication".

4.4 Marking, packaging, and labeling. The product shall be labeled, packed and marked in accordance with 49 CFR parts 100 to 185.

5. PRODUCT CONFORMANCE PROVISIONS

5.1 Product conformance. The product provided shall meet the salient characteristics of this commercial item description, conform to the producer’s own drawings, specifications, standards, and quality assurance practices, and be the same product offered for sale in the commercial market. Any certificate of analysis (COA) offered with the product to the commercial market shall be provided with each lot or batch of product. The lot or batch number and the date of inspection/analysis from the offer shall be on each COA. A full COA proving the salient characteristics of the product meets government requirements shall be provided upon request.

5.2 Market acceptability. The product offered must have been sold to the Government or commercial market.

6. PACKAGING. Preservation, packing, and marking shall be as specified in the contract or order.

7. NOTES.

7.1 Intended use. The intended use of the types and classes are:

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T08:05Z
Check the source to verify that this is the current version before use.

Type II - Herbicide is effective on easy-to-kill or moderately easy-to-kill weeds

Type III - Class 1. Herbicide is effective on deep rooted perennial weeds, old semi resistant weeds and woody species.
- Class 2. Herbicide is used the same as Class 1 however Class 2 should be used in areas where sensitive crops such as cotton are grown.

7.2 Source of documents.

7.2.1 Electronic copies of Official Methods of Analysis of AOAC INTERNATIONAL may be obtained from www.aoac.org.

7.2.2 Electronic copies of the FAR may be obtained online at www.gpo.gov.

7.2.3 Electronic copies of CFR documents can be obtained from https://gov.ecfr.io/cgi-bin/ECFR.

7.2.4 Copies of FED-STD-313 can be obtained from https://quicksearch.dla.mil.

7.3 Ordering data. The contract or order should specify the following:
a. CID document number and revision
b. Product type (and class if necessary)
c. Packaging requirements

7.4 National Stock Numbers (NSNs). The following NSNs correspond to this CID: 6840-00-664-7060 type II and 6840-00-577-4194 type III, class 2. These may not be the only NSNs associated with this document.

7.5 Key Words.

2,4-D
Weed control
Weed killer

7.6 Changes from previous issues. Marginal notations are used in this revision to identify changes with respect to the previous issue.

MILITARY INTERESTS:

CIVIL AGENCY COORDINATING ACTIVITY:
GSA - FAS

Custodians:
DLA – GS

Preparing Activity:
DLA – GS3

(Project 6840-2020-001)

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T08:05Z
Check the source to verify that this is the current version before use.

NOTE: The activities listed above were interested in this document as of the date of this document. Since organizations and responsibilities can change, you should verify the currency of the information above using the ASSIST database at https://assist.dla.mil/.

Source: https://assist.dla.mil -- Downloaded: 2026-10-26T08:05Z
Check the source to verify that this is the current version before use.