AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, IGATPURI
ZONAL AGRICULTURE RESEARCH STATION, IGATPURI.

Ph. No. 02553-244032 **E-mail: igatpuri.amfu@gmail.com**

34) Weather based Agromet Advisory committee meeting dated 31.07.2026 District: Nashik

Last Week Weather Summary (25.07.2026 to 31.07.2026)							Weather Parameters	Weather Forecast (01.08.2026 to 05.08.2026)				
25	**26**	**27**	**28**	**29**	**30**	**31**	**Date**	**01**	**02**	**03**	**04**	**05**
76.6	70.0	130.0	38.2	35.6	38.9	46.8	**Rainfall (mm)**	75	15	4	4	3
24.6	25.0	23.5	24.2	24.5	25.0	24.2	**Max. Temp. (^{0}C)**	26	27	29	29	29
21.8	22.0	22.0	22.5	22.8	23.0	23.4	**Min. Temp. (^{0}C)**	23	23	24	24	24
Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	**Cloud Cover**	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy
97	96	97	97	95	96	97	**Max. RH (%)**	96	93	93	93	93
92	96	97	95	94	96	96	**Min. RH (%)**	95	83	81	78	81
10.9	7.7	5.8	5.6	7.4	7.4	6.7	**Wind Speed (km/hr)**	20.2	28.3	21.1	19.9	18.6

Agromet Advisory Based on Weather Forecast Prediction

Crop	Stage	Advisory
Weather Summary		Considering the forecast for the next five days, there is possibility of moderate to heavy rain on 01st & 02nd Aug. 2026 & rest of the day's light rainfall in Nashik district. The sky will cloudy for next five days. Maximum Temperature staying in between 26-29 Degree Celsius & Minimum Temperature 23-24 Degree Celsius & the wind speed will remain between 18.0-28.0 kmph for the next five days.
Weather Alerts/ warning:		**Considering Impact based forecast (IBF) for Agriculture for upcoming Heavy to very heavy rainfall (Red alert) at few places with extremely heavy rainfall at isolated places strong winds reaching 50-60 kmph gusting 70 kmph very likely of Nashik district & Ghat region of Nashik district on 31st July 2026. Heavy to very rainfall (Orange alert) at isolated places of Ghat region of Nashik on 01st August 2026 (Based on the District Level Forecast and warning issued by RMC Mumbai on issued on 31.07.2026).**
General Advisory		Considering gusty wind with heavy rain Alerts/ warning forecast, protect the livestock**,** poultry and yourself from gusty wind with heavy rainfall. To protect Mango & Pomegranate orchards from gusty winds, it is advice to use the bamboo or wooden sticks for support. The Pomegranate plants should be tied to each other with the help of plastic rope. Considering weather forecast, in dry land /rainfed area prepare the structures for in situ moisture conservation like compartment bunding. Also management should be done to store falling rain water in the field. Planting of kharif season vegetable crops e.g. Bottle gourd, Bitter gourd, Ridge gourd, Sponge gourd, Pumpkin etc. Before transplanting, the roots of Tomato, Brinjal, Chilli, cauliflower and cabbage plants should be immersed in a solution of insecticide, fungicide and bacterial fertilizer. Kharif Okra and Cluster bean crops should be cultivated. Considering the forecast of rainfall**,** it is advised to spraying of insecticides and fungicides on kharif crops should be stopped or postponed for given period Six weeks old seedlings of cauliflower and cabbage should be used for planting. Rain water should not be allowed to accumulate in the planted vegetable crop.
SMS		**Considering the forecast and warning of heavy rainfall, immediate arrangements should be made to drain out excess water from areas under transplanted Kharif paddy, Finger millet and Little millet nurseries or transplanted fields, as well as from crops such as maize, soybean, pearl millet, green gram, and cotton.**
***Kharif* Paddy**	Transplanting / Growth stage	Considering the rainfall received or availability of water, the transplanting of Paddy should be completed up to 31st July 2026. When planting, 2 to 3 seedlings should be kept per hill; however, for hybrid varieties, only 1 to 2 seedlings should be kept per hill. Two to three rough weeding are essential after transplanting. Weed infestation should be controlled excellently by maintaining water level up to 5 to 6 cm. For proper growth of crops in the paddy field, there should be maintain a water level of 1 to 2 cm from the time the seedlings are transplanted until the seedlings are established. After 3-4 days of transplanting of Paddy use the Urea DAP Briquette @ 70 kg per acre & avoid nutrient losses. Maintain the water level from 1 to 2 cm until the newly transplanted paddy seedlings are stable in paddy field. Also, maintain the water level up to 2-3 cm at the plants where they are in primary stage.
***Kharif* Finger millet**	Transplanting	Considering the rainfall forecast transplanting of Finger millet seedlings should be continuing. 25 to 30 days old seedlings prepared on BBF should be used for transplanting. In light soils of sub plateau region of Maharashtra, transplanting of Finger millet /Ragi crop at row to row (twin line method) distance 20:40 cm for higher yield and economic benefits of Ragi. 5 ton of FYM per hectare + 75 percent of recommended fertilizer quantity (45: 22.5: 00 Nitrogen: Phosphorus: Potash kg/ha) in the form of briquettes should be given at the time of planting (A tablet of 2.7 gm on depth of 5 to 7 cm at 35 cm distance in 20 cm row to row method). Considering the rainfall received or availability of water, the transplanting of Finger millet should be completed up to 31st July 2026.
Niger	Sowing / Intercultural operation	Farmers who have not yet sown the crop should complete the sowing of the Niger crop by July 31st 2026. To maintain optimum plant population (3.3 lakh/ha) in the areas where the crop is sown, thinning is recommended to remove extra plants after two weeks of sowing or when the seedlings attain 8-10 cm height.
Maize	Sowing	Consideration rainfall forecast, sowing of maize during the Kharif season up to this week should be completed. During the Kharif season, maize should be sown in flat beds using the dibbling method at a depth of 4–5 cm. A seed rate of 15–20 kg per hectare should be used, maintaining a spacing of 75 × 20 cm for late and medium-duration varieties, and 60 × 20 cm for early-maturing varieties. After sowing, during a break in the rainfall, the herbicide Atratop 50% should be sprayed on the soil at a rate of 2.5 kg/hectare.
Soybean	Sowing	Considering the forecast of rains, soybean crop has been sown at 99.96 percent in the eastern drought prone and plain zone areas, but in the remaining areas, improved varieties of soybean crop such as Phule Kimaya DS-753 and Phule Kalyani DS-228 varieties recommended by Mahatma Phule Agricultural University are suitable for late sowing. hence the sowing should be completed by 31 July 2026.
Groundnut	Sowing	Considering rainfall forecast, groundnut sowing should be completed up to this week during the Kharif season, following seed treatment. Sowing should be done using the Pabhari (seed drill) or Dibbling method, maintaining a spacing of 30 × 10 cm. Seed treatment prior to sowing is essential to protect the crop against seed-borne and seedling-stage diseases.
Pearl millet	Sowing	Considering the rainfall forecast, pearl millet (bajra) sowing for the Kharif season should be completed up to this week, adhering to recommended spacing and applying the prescribed dosage of chemical fertilizers along with bacterial culture seed treatment. Sowing during the Kharif season should be done at a depth of 2–3 cm using either the ridge-and-furrow method or the flat-bed method. A seed rate of 3–4 kg/hectare should be used, maintaining a spacing of 45×15 cm in dryland areas and 30×15 cm under irrigated conditions. It is essential to subject the seeds to appropriate seed treatment before sowing.

AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, IGATPURI
ZONAL AGRICULTURE RESEARCH STATION, IGATPURI.

Ph. No. 02553-244032 **E-mail: igatpuri.amfu@gmail.com**

Grapes		Downy Mildew Control Downy mildew becomes active when there is at least 4 mm of rainfall for three consecutive days and the temperature drops to 25°C or lower. The infection typically appears first in areas where water accumulates, so keep a close watch on those spots. Spray only after the disease is detected. Since the disease spreads with the wind, prioritize spraying orchards located downwind from the infected area. Avoid using systemic fungicides at this stage; they are less necessary now and are more effective when applied 50–60 days after fruit pruning. Spraying copper-based fungicides is currently effective for controlling downy mildew, as well as fungal and bacterial blight and rust diseases. There is a low risk of copper toxicity affecting the older leaves at this stage.
Pomegranate		Ambe Bahar Irrigation should be scheduled by taking into account the weather forecast and the soil's water-holding capacity. If weather conditions become favorable for disease development, immediately spray a 0.5% Bordeaux mixture. Harvest the fruit in orchards where the crop has reached the harvesting stage.
Mango		Water Drainage Considering the possibility of moderate rains, arrangements should be made for drainage of excess water in the garden. New planting should be done when the intensity of rain is low. Care should be taken that water does not accumulate near the base of newly planted plants.
Kharif Onion	Nursery management/ Transplanting	An outbreak of wilt disease has been observed in onion nurseries due to the continuous rainfall over the past week. To control the disease, drench the soil with a mixture of 30 grams of Copper Oxychloride per 10 liters of water. Considering rainfall forecasts and the actual satisfactory rainfall or water availability farmers should adopt the following integrated crop protection method in transplanting of onion seedlings. Integrated Crop Protection Depending on the season, planting in an area should be completed in one week by maintaining a longer gap between two seasons to break the life cycle of diseases and insects. Carbendazim 1 g and carbosulfan 2 ml two hours before replanting. Per liter of water should be dipped in this solution and replanted only later. 0.6 ml of stickers used while spraying. Per liter of water should be used as follows. Pesticides should be alternated as recommended. Insect resistance increases when used as a single insecticide.
Tomato		Transplanting Rainy Season Tomatoes For transplanting, select vigorous seedlings that are 25–30 days old, 10–15 cm tall, and have 6–8 leaves; avoid using wilted, diseased, or damaged seedlings. Before planting, dip the roots for 10–15 minutes in a solution containing Imidacloprid (17.8 SL) (4 ml) and either Metalaxyl-M (31.8 ES) (6 g) or Carbendazim (10 g) per 10 liters of water. Plant at a spacing of 30 × 90 cm, ensuring that no pressure is exerted on the seedling stems.
Animal Husbandry (Cow, buffalo)		Considering the thunderstorm, moderate rain & gusty wind forecast it is advised to Keep animals away from tractors and other metal farm equipment. Keep cattle and goats in sheds & keep animals away from open water, ponds or rivers to protect them from thunder and lightning. Don't allow your animals to congregate under trees. Watch your animals closely and try to keep them under your direct control. Livestock Management Increased humidity during the monsoon leads to a higher prevalence of bacteria and parasitic worms; therefore, all animals should be administered deworming medication upon veterinary advice. Livestock should be vaccinated on time to prevent disease outbreaks. Measures must also be taken to control the infestation of ticks and fleas in the cattle shed, which tends to increase during the monsoon season.
Goat		Goat Management During July–August Control external parasites and deworm newly born kids. Vaccinate all goats against enterotoxaemia, PPR, and tetanus. Increase the feed ration for breeding does and prepare the breeding bucks. If planning for synchronized breeding, keep a 'teaser buck' in the shed to detect estrus.
Sheep		Management in sheep in July The sheep in the flock should be examined and treated for red urine disease and jaundice on the advice of a veterinarian. Sheep should be dewormed as recommended. Take special care of pregnant sheep. Body weight of lambs aged 9 to 12 months should be taken. All sheep should be vaccinated against tetanus.
Poultry		Since there is a risk of drinking water becoming contaminated during the monsoon season, chickens should be provided with clean and disinfected water. Chlorine or bleaching powder should be mixed into the water in appropriate quantities, and the water tank must be kept clean, covered, and regularly maintained.

- **In the event of a delay in sowing, the 'Contingent Crop Planning' handbook prepared by the university should be utilized. Additionally, alternative seeds suitable for late sowing should be kept available.**
- **Bio-fertilizers produced by the university are highly effective and beneficial for seed treatment; their use enhances crop germination capacity, provides essential nutrients to the crops, and reduces expenditure on chemical fertilizers.**

Source:

1) Weather Forecast : Research Section, Mumbai
2) Last week weather summary : GKMS Observatory, ZARS, Igatpuri, Dist. Nashik.

Place : ZARS, Igatpuri
Date : 31.07.2026

Sd/-
Nodal Officer, GKMS,
AMFU Igatpuri &
Associate Director of Research
ZARS, Igatpuri, Dist. Nashik