AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, IGATPURI
ZONAL AGRICULTURE RESEARCH STATION, IGATPURI.

Ph. No. 02553-244032 **E-mail: igatpuri.amfu@gmail.com**

19) Weather based Agromet Advisory committee meeting dated 09.06.2026 District: Nashik

Last Week Weather Summary (03.06.2026 to 09.06.2026)							Weather Parameters	Weather Forecast (10.06.2026 to 14.06.2026)				
03	04	05	06	07	08	09	Date	10	11	12	13	14
18.7	0.0	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)	8	4	3	2	7
35.0	33.5	35.0	35.0	35.7	35.0	34.7	Max. Temp. (^{0}C)	36	37	37	37	37
21.2	23.7	24.8	25.0	25.0	25.2	25.2	Min. Temp. (^{0}C)	26	26	26	26	26
Cloudy	P Cloud	Cloudy	Cloudy	P Cloud	Cloudy	Cloudy	Cloud Cover	P Cloud	P Cloud	P Cloud	P Cloud	P Cloud
88	84	86	85	84	82	86	Max. RH (%)	70	70	69	69	70
51	46	43	44	42	49	50	Min. RH (%)	51	50	51	52	53
6.1	6.3	6.9	6.7	7.1	7.8	8.3	Wind Speed (km/hr)	21.7	22.6	19.3	20.5	20

Agromet Advisory Based on Weather Forecast Prediction

Crop	Stage	Advisory
Weather Summary		Considering the forecast for the next five days, there is possibility of light rain between 10th to 14th June 2026 in Nashik district. The sky will partial cloudy for next five days. Maximum Temperature staying in between 36-37 Degree Celsius & Minimum Temperature 26 Degree Celsius & the wind speed will remain between 19.3-22.6 kmph for the next five days.
Weather Alerts/ warning:		Considering Impact based forecast (IBF) for Agriculture for upcoming thunderstorm accompanied with lightning, light to moderate rain & gusty wind with speed 30-40 kmph (Yellow alert) on 09th June 2026 at isolated places of Nashik & Ghat region of Nashik district..(Based on the District Level Forecast and warning issued by RMC Mumbai on issued on 09.06.2026).
General Advisory		Considering weather forecast & Alerts/ warning, protect the harvested summer crops, livestock, poultry and yourself from thunderstorm with lightning & rainfall. Considering weather forecast & Monsoon forecast farmers are advised not to rush into sowing; hence kharif crops should be sown only after 3 to 4 consecutive days of adequate monsoon rainfall (75 to 100 mm) /the availability of soil moisture/wafsa condition in the scarcity zone and plain zone (Sinnar, Yeola, Nandgaon, Malegaon, Chandwad, Nashik and Niphad) of Nashik district. Farmers should sow only after proper and sufficient rains, farmers should be careful of fake seeds and fertilizers, farmers should buy fertilizers and seeds from authorized centers only after receiving a firm bill, beware of fake varieties and stay away from fraud. Farmers should be wary of indigenous fake varieties. Unauthorized, illegal and unpaid seeds and fertilizers should not be purchased. Also, if such agricultural inputs are found to be being sold illegally, a complaint should be made to the concerned Taluka Agriculture Officer or District Superintendent of Agriculture Officer's office. Farmers are advised to get soil tested to control over-fertilization, proper planning of field crops, double economic benefits, planning of crop cultivation and the amount of nutrients in the soil. Farmers who have irrigation facilities should sow seeds to produce onion, brinjal, chilli, tomato, cauliflower and cabbage plants. Seed treatment should be done before sowing. Initially the nursery should be irrigated by water can with sprayer, then irrigation water can be used. Farmers should cultivate the land and test the soil for sowing rice and other kharif crops.
SMS		**Considering weather forecast & Alerts/ warning, farmers are advised to protect themselves by using the 'Damini' mobile app to anticipate lightning strikes.**
***Kharif* Paddy**	Preparatory Tillage	Procured good quality improved varieties of paddy like Indrayani, Phule Samridhi, Bhogavati and Phule Radha. Also for sowing of paddy crop 1 meters wide and 15 cm. height and length of BBF (broad bed furrow) as per requirement should be prep ared. 4 Are of nursery are enough for one acre of plantation**.** For 1 Are (guntha) Mix of 250 grams of farm yard manure or composted manure, 500 grams of nitrogen, 400 grams of phosphorus, 500 grams of potash fertilizer in the soil.
***Kharif* Finger millet**	Preparatory Tillage	Improved Varieties: Phule Nachani, Dapoli-1, Dapoli Safed, Dapoli-2, Phule Kasari. For nursery sowing of Finger millet crop 1 to 1.5 meters wide and 8-10 cm. height and length of BBF (broad bed furrow) as per requirement should be prepared in first week of June. per sq.m. 3 kg of farm yard manure should be applied on the area.
Maize	Pre sowing preparation	During the Kharif season, maize should be sown between June 15 and July 15 immediately after the sufficient rains (75 to 100 mm). Mixed and hybrid varieties of maize crop yield 60 to 80 percent more than local varieties. There are mixed and hybrid varieties of maize which mature in different periods and planning should be done to purchase seeds of suitable variety for planting by taking into account the fertility of the land and amount of rainfall. Name of the variety – Average grain yield (quintal/ha): Late variety (100 to 110 days): Bio-9681 60-70; HQPM-1 60-65; HQPM-1 55-60; Sangam 75-80; Kuber 75-80 Early variety (80 to 90 days) and very early variety (70 to 80 days): Pusa Hybrid Maize-1 40-50; Vivek Hybrid Maize-21 45-50; Vivek Hybrid Maize-27 50-55; Maharaja 60-65 Moderate variety (90 to 100 days): Rajashri 70-75 Kharif 95-100 Rabi; Bio-9637 70-75; Phule Maharshi 75-80 For Fodder - Late composite varieties (100 to 110 days) African tall 60-70 tons of green fodder 40-50 km. grain For Sweet corn Phule Madhu 125-130 km/ha. Corn
Soybean	Pre sowing preparation	Procured good quality seeds of improved varieties of soybean crop such as Phule Durva (KDS 992), Phule Kimaya (KDS 753), Phule Sangam (KDS 726), Phule Agrani (KDS 344) , Phule Kalyani (KDS 228), K. S. 103, J. S. 335, J. S. 9305, J. S. 9560, Samriddhi (MAUS 71), MAUS 162, MAUS 158, MAUS 612, MAUS 1188, Plan to buy MAUS 1281. Soybean crop should be sown only after 3 to 4 consecutive days of adequate monsoon rainfall (65 to 100 mm) /the availability of soil moisture/wafsa condition till the first week of July.
Groundnut	Pre sowing preparation	During the kharif season, Groundnut should be sown between 15th June and 15th July immediately only after the sufficient rains (75 to 100 mm). Procured good quality seeds of Groundnut improved varieties like SB-11, JL-24 (Phule Pragati), TAG-24, TG-26, JL-501, Phule Unnati, JL - 1085 (Phule Dhani).

AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, IGATPURI
ZONAL AGRICULTURE RESEARCH STATION, IGATPURI.

Ph. No. 02553-244032 **E-mail: igatpuri.amfu@gmail.com**

Grapes		Management of Stromacium barbatum stem bore From the beginning of June, 1 light trap per acre should be installed outside the vineyard. Spraying should be planned after these stemborer are found in the trap. During the period from 1st to 20th June, 5 percent neem extract or Neem leaf extract or Azadirachtin (1000 ppm) 2.3 ml. Per liter of water should be sprayed five to six times on the main trunk and across at intervals of three to four days. (**Undertake the agricultural practices only when during clear weather).**
Pomegranate		Coating the Stem Before taking bahar in the two years pomegranate orchard, after harvesting of fruits apply the following coating on the stem/branch up to two feet from the ground. Red clay 4 kg + Imamectin benzoate (5% SG) 20 gm + Copper oxychloride (50% WP) 25 gm per 10 liters of water above coating should be applied in rotation with 10 percent Bordeaux mixture. (**Undertake the agricultural practices only when during clear weather).**
Mango		For fully grown mangoes in monsoons every year 50 kg of farm yard manure + 1500 grams of nitrogen + 500 grams of phosphorus + 500 grams of potash should be given. Nitrogen dose should be divided into two equal installments in July and September. So phosphorus and Potash should be given in one installment in July.
Tomato		Field should be prepared for nursery of tomato crop. Seedlings should be prepared on broad bed furrows. Seeds should be sown between May to June to produce tomato plant. Seed treatment should be done before sowing. Initially the nursery should be irrigated by water can with sprayer, then irrigation water can be used. Nursery period Season — Seed sowing period — Transplanting period Kharif — May to June — June to July Rabi — September to October — October to November Summer — January to February — February to March
Animal Husbandry (Cow, buffalo)		Considering the thunderstorm & moderate rain forecast it is advised to keep milch animals indoors. Keep livestock in sheds and keep animals away from open water, lakes, or rivers to protect them from hail. Considering the moderate rain forecast farmers should advice to keep the stored fodder of livestock in a safe place or covered with plastic/tarpaulin. Care should be taken to ensure that contaminated or silty water does not enter the drinking water of the animals. Deworming the animals. Sulfa tablets should be used for the disease of Bovine cholera caused by young grass. Do not tie the animals in open spaces. Keep them in a barn, shed or strong building. If there is a rainy weather and lightning, stop the work in the field immediately and move the animals to a closed and safe place. The roof and walls of the barn should not be made of metal, because metal attracts electricity. If the animals are in water, take them out immediately.
Goat		Management of goats in June Goats with low production and reproductive disorders should be removed from the herd. (20 percent of the total number). New breeds and good goats should be selected in their place. New breeding buck should be selected. Keep an eye on various physical diseases of goats. Prepare for reproduction and breeding of goats.

Source:
1) Weather Forecast : Research Section, Mumbai
2) Last week weather summary : GKMS Observatory, ZARS, Igatpuri, Dist. Nashik.

Place : ZARS, Igatpuri
Date : 09.06.2026

Sd/-
Nodal Officer, GKMS,
AMFU Igatpuri &
Associate Director of Research
ZARS, Igatpuri, Dist. Nashik