AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR
ZONAL AGRICULTURAL RESEARCH STATION, SHENDA PARK, KOLHAPUR

Ph. No. 0231 2692416 **E-mail: adrkolhapur@rediffmail.com**

Weather based Agromet Advisory committee meeting dated 30.04.2024

District: SANGLI

Last Week Weather Summary (24.0.2024 to 30.04.2024)							Weather Parameters	Weather Forecast (01.05.2024 to 08.05.2024)				
24	25	26	27	28	29	30	Date	01	02	03	04	08
-	-	-	-	-	-	-	Rainfall (mm)	0	0	0	0	0
-	-	-	-	-	-	-	Max. Temp. (^{0}C)	42	41	40	41	41
-	-	-	-	-	-	-	Min. Temp. (^{0}C)	26	25	24	24	25
-	-	-	-	-	-	-	Cloud Cover	1	0	1	0	1
-	-	-	-	-	-	-	Max. RH (%)	61	41	42	60	64
-	-	-	-	-	-	-	Min. RH (%)	30	29	25	25	28
-	-	-	-	-	-	-	Wind Speed(km/hr)	17	18	15	16	17
-	-	-	-	-	-	-	Wind Direction	WNW	WNW	WNW	WNW	WNW

Agromet Advisory Based on Weather Forecast

Crop	Stage	Advisory
Weather Summary		Weather forecast given by Regional Meteorological Centre, Mumbai has indicated that there is possibility of **manly clear sky from 1st to 5th May in** Sangli district. During next 5 days **Maximum temperatures will remain nearby 40.0 to 42.0 and minimum nearby 24.0 to 26.0** ^{o}C respectively. Morning relative humidity will remain between 41 to 64% and Afternoon relative humidity will remain 25 to 30%. Wind speed will remain between 15 to 18 km/hr. According to Extended Range Forecast (ERF) in Madhya Maharashtra Sub Division rainfall will remain below normal, maximum temperature will remain normal and minimum temperature will remain normal valid from 5th to 11th May, 2024
General Advisory		❖ Farmers should carry out farm operations during morning or evening hours. Working in field during afternoon hours should avoided as there is forecast of increase in Tmax upto 42.0^{o}C ❖ Possibility of increase in maximum temperature upto 42°C thus farmers should take care of themselves and their cattles.
SMS Advisory		➢ Carry out mulching in fruits and vegetable crop. Provide irrigation to Orchards and vegetable crop as per the availability and as per the requirement of crop.
Sugarcane	**Establishment**	➢ During May month the rate of evaporation increases up to 9.24 mm per day thus sugarcane plant requires 9.33 liters water per day to fulfill that requirement it is need to run Drip irrigation unit (4 liters dripper) for 2 hour and 20 minutes per day. ➢ At the time of earthing up of Suru Sugarcane give fertilizer dose of 100 kg Nitrogen (217 kg Urea), 55 Kg Phosphorus (344 kg Single Super Phosphate) and 55 Kg Potash (Murate of Potash). ➢ Remove lower dried and matured leaves and use these leaves as mulching in furrows. It will minimise evaporation from soil and it will helpful for maintaining soil moisture.
Sunflower	**Maturity**	➢ Carry out harvesting of matured Sunflower crop and dry under bright sunlight and then carry out threshing.
Tomato	**Fruit development**	➢ There is forecast of increase in Tmax > 40.0 thus increased Tmax will increase flower and fruit drop in Tomato thus to minimise flower and fruit drop provide sufficient irrigation to Tomato crop
Grapes	**Prunning**	➢ After foundation pruning, during shoot growth stage, apply 12,750 – 15,300 L/acre per day of irrigation water.
Ginger	**Seed treatment**	➢ Purchase Ginger rhizomes for cultivation and keep the rhizomes under shade after proper Chemical and Biological rhizome treatment. ➢ Carry out rhizome treatments with Azospirillum @ 25 gram and P.S.B @ 25 gram per liter of water as biological treatment of Rhizomes. ➢ Select Rhizomes having 25 to 55 gram weight, 2.5 to 5 cm length for cultivation. The

AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR
ZONAL AGRICULTURAL RESEARCH STATION, SHENDA PARK, KOLHAPUR

Ph. No. 0231 2692416 **E-mail: adrkolhapur@rediffmail.com**

		dormancy of rhizomes should be over and bulging of 2 to 3 buds. Cultivation of 1 hector of Ginger 25 Quintals of rhizomes seed require.
Mango	**Fruit development**	➢ Carry out harvesting of mango fruit at 80 to 85% maturity stage. Harvesting should be done with help of nutan mango harvester keep the harvested fruits at safe place as rainfall is forecasted. ➢ Keep the harvested fruits in shade to prevent development of spongy tissue and from heat. Transport of matured fruit should be done during night. Do not spray the plant with any type of fungicide or insecticide at least 8 days before harvesting of fruit.
Coconut	**Growth**	➢ There is possibility of increase in maximum temperature thus Make arrangement of shade with help of shade net to newly planted Coconut plants.
Animals	**Growth**	➢ Keep the cattle's under shade. If possible sprinkle water 4 to 5 times through out day. ➢ Provide sufficient, pure and cool water for drinking to animals. ➢ Do not allow cattle's for grazing during afternoon hours as the intensity of Sunshine and maximum temperature will be high during that period. ➢ Carry out proper ventilation in cattle shed, install fans and foggers for maintaining temperature in cattle shed. ➢ Allow Buffalo for wallowing in water as they have fewer sweat glands compared to cattle to maintain their body's thermo equilibrium. Water soaked gunny bag or other cloth should kept on a body to avoid occurrence of heat stroke.
Poultry	**Birds**	➢ To protect Poultry birds from heat cover the sides of Poultry shed with gunny bags and spray water on these gunny bags to minimize the temperature inside the poultry house. Cover the roof of poultry house with grass or with gunny bags which will help to keep the reduce temperature of poultry house.

Source:
1) Weather Forecast : Research Section, Mumbai

Place : ZARS, KOLHAPUR
Date :30.04.2024

Sd/-
Nodal Officer, GKMS, &
Associate Director of Research,
ZARS, Kolhapur