

Development and Validation of a french kidney donor marginality score

Corentin Choisy¹ Magali Giral^{2,3} Etienne Dantan¹


¹SPHERE UMR 1246, Nantes Université, Univ Tours, CHU Nantes, Inserm, Nantes, France

²CR2TI UMR 1064, Nantes Université, ITUN, CHU Nantes, RTRS Centaure, Nantes, France

³Centre d'Investigation Clinique en Biothérapie, Nantes, France

Background

- **Kidney transplantation (KT)**: recognised as the best treatment for **end-stage chronic renal disease**
 - **Graft shortage** in countries with aging population
 - ⇒ **Necessity for expansion of graft pool**
- Wide use of **marginal grafts** with suboptimal properties for patient-graft survival
- Decision making tools to assist clinicians in evaluating graft proposals
 - **ECD [1]** Older than 60 years old or between 50 and 59 with at least 2 comorbidities among: high serum creatinine, history of hypertension and death by CVA
 - ⇒ **Binary criterion, no gradient between less and more marginal donors**
 - **KDRI/KDPI [2]** Continuous/percentile scale defined by 10 donor features
 - ⇒ **Not adapted to the french population [3] and prone to increased graft refusal rate [4]**




Objectives

Develop and validate a **kidney donor marginality score**, adapted to the french population and suited for the current practices in KT in France

- **Donor/recipient interactions** will be studied in order to express donor marginality in relation to recipient characteristics
- **Recipient loss-of-chance** related to receiving a marginal graft as defined by the proposed score will be studied

Materials and Methods

- **7622 patients** from the DIVAT national kidney transplant cohort
- First-time deceased donor graft recipients without donor-recipient ABO incompatibility
- **Multivariate Cox regression** model built via blockwise variable selection approach (Donor/Graft/Recipient features integrated/removed from the model in 11 steps)
- Assessment of **period/center effect** via adjusted and frailty models
- New variable selection steps after integration of period/center effects



Funding

This work was supported by the French Biomedicine Agency (reference: AOR Greffe 2022)

Results




Figure 1. Patient/allograft survival analysis

	β	95% CI
Donor age	-0.020	[-0.037– -0.003]
Donor after cardiac death	0.691	[0.416–0.967]
Donor death by CVA	-0.616	[-1.224– -0.008]
Donor serum creatinine	0.001	[0.0001–0.002]
Donor height	-0.011	[-0.018– -0.004]
Donor weight	0.005	[0.001–0.010]
Positive donor CMV serology	0.124	[0.010–0.239]
HLA incompatibilities ≥ 4	0.151	[0.009–0.293]
Time on dialysis	0.0001	[0.0001–0.0001]
Transplanted before 2012	0.190	[0.040–0.340]
Recipient age	-0.021	[-0.036– -0.005]
Recipient BMI	0.011	[-0.003–0.025]
Recipient history of diabetes	0.376	[0.234–0.517]
Recipient history of cardiovascular disease	0.408	[0.239–0.527]
Hemodialysis	0.250	[0.048–0.452]
Donor age*Recipient age	0.001	[0.0001–0.001]
Donor death by CVA*Recipient age	0.014	[0.003–0.024]

Table 1. Proportional hazards Cox model analysis of retained factors and interactions after multivariate selection

- Donor age, beating heart status ($HR = 2.00$), cause of death, creatininemia, height, weight and CMV serology ($HR = 1.13$) will be integrated into the score. For a given donor, the score will differ depending on recipient age and donor/recipient HLA ABDR incompatibilities.

What's next

- **Internal validation** of the model, $\frac{1}{3}$ of the database allocated to a random validation sample
- **Validity assessed through:**
 - **Calibration:** calibration plot, predicted survival by score quantile, Brier score
 - **Discrimination:** time-dependent ROC curve (AUC, PPV/NPV) and time-dependent ROC curve per recipient strata
- **Propensity score**-based analysis to estimate population-averaged effects and propose a recipient loss-of-chance-based approach
- Development of an **online score computation tool** for clinicians

References

- [1] Robert A Metzger, Francis L Delmonico, Sandy Feng, Friedrich K Port, James J Wynn, and Robert M Merion. Expanded criteria donors for kidney transplantation. *American Journal of Transplantation*, 3:114–125, 2003.
- [2] Panduranga S Rao, Douglas E Schaubel, Mary K Guidinger, Kenneth A Andreoni, Robert A Wolfe, Robert M Merion, Friedrich K Port, and Randall S Sung. A comprehensive risk quantification score for deceased donor kidneys: the kidney donor risk index. *Transplantation*, 88(2):231–236, 2009.
- [3] Etienne Dantan, Florent Le Borgne, Magali Giral, Angelina Dion, Anne-Hélène Querard, and Yohann Foucher. Covariates adjustment questioned conclusions of predictive analyses: an illustration with the kidney donor risk index. *Journal of Clinical Epidemiology*, 135:103–114, 2021.
- [4] Olivier Aubert, Peter P Reese, Benoit Audry, Yassine Bouatou, Marc Raynaud, Denis Viglietti, Christophe Legendre, Denis Glotz, Jean-Philippe Empana, Xavier Jouven, et al. Disparities in acceptance of deceased donor kidneys between the united states and france and estimated effects of increased us acceptance. *JAMA internal medicine*, 179(10):1365–1374, 2019.
- [5] Rapport annuel 2015. *Agence de la Biomédecine*, 2015.
- [6] V Bongard, J Dallongeville, D Arveiler, J-B Ruidavets, D Cotel, A Wagner, and J Ferrières. Estimation et caractérisation de l'insuffisance rénale chronique en france. In *Annales de Cardiologie et d'Angéiologie*, volume 61, pages 239–244. Elsevier, 2012.
- [7] Transplantation rénale - accès à la liste d'attente nationale - recommandation de bonne pratique. *Haute Autorité de Santé*, 2015.

