

Table 1. Main sample characteristics

	Children/Early adolescents		Late adolescents	
	ASD (N=339)	no ASD (N=80)	ASD (N=360)	no ASD (N=31)
Age (years)				
Mean (SD)	11.0 (2.64)	10.9 (2.41)	19.8 (3.10)	19.5 (3.12)
Median [Min, Max]	11.0 [6.00, 24.0]	11.0 [6.00, 16.0]	20.0 [15.0, 25.0]	19.0 [15.0, 25.0]
Sex				
Female	66 (19.5%)	24 (30.0%)	140 (38.9%)	9 (29.0%)
Male	273 (80.5%)	56 (70.0%)	220 (61.1%)	22 (71.0%)
FSIQ				
Mean (SD)	105 (15.9)	102 (14.2)	110 (16.8)	98.7 (13.1)
Median [Min, Max]	104 [70.5, 149]	102 [75.0, 144]	110 [73.0, 158]	99.0 [75.0, 123]
Missing	32 (9.4%)	7 (8.8%)	40 (11.1%)	8 (25.8%)
Vineland (socialization)				
Mean (SD)	53.2 (26.5)	69.1 (29.0)	50.7 (25.8)	51.2 (19.7)
Median [Min, Max]	50.0 [14.0, 114]	70.0 [20.0, 134]	50.0 [12.0, 118]	52.0 [20.0, 78.0]
Missing	76 (22.4%)	16 (20.0%)	197 (54.7%)	16 (51.6%)
Vineland (total)				
Mean (SD)	60.3 (24.3)	75.5 (24.6)	55.8 (25.3)	49.5 (19.8)
Median [Min, Max]	60.0 [20.0, 124]	77.0 [21.0, 132]	56.0 [20.0, 141]	50.0 [20.0, 95.0]
Missing	102 (30.1%)	25 (31.3%)	200 (55.6%)	16 (51.6%)
ADOS total score				
Mean (SD)	12.3 (4.84)	7.78 (4.55)	10.9 (4.27)	6.33 (4.69)
Median [Min, Max]	12.0 [0, 26.0]	7.00 [0, 21.0]	11.0 [1.00, 24.0]	5.00 [1.00, 20.0]
Missing	2 (0.6%)	0 (0%)	2 (0.6%)	1 (3.2%)
Other psychiatric condition				
Any	242 (71.4%)	59 (73.8%)	274 (76.1%)	17 (54.8%)
None	61 (18.0%)	12 (15.0%)	52 (14.4%)	11 (35.5%)
Missing	36 (10.6%)	9 (11.3%)	34 (9.4%)	3 (9.7%)
Ongoing psychiatric treatment				

	Children/Early adolescents		Late adolescents	
	ASD (N=339)	no ASD (N=80)	ASD (N=360)	no ASD (N=31)
At least one treatment	82 (24.2%)	19 (23.8%)	78 (21.7%)	2 (6.5%)
No treatment	205 (60.5%)	50 (62.5%)	220 (61.1%)	24 (77.4%)
Missing	52 (15.3%)	11 (13.8%)	62 (17.2%)	5 (16.1%)