

Methods/Results

Methods

Population

We present a spatial explicit stochastic agent based model that recreates the day to day dynamics in a typical nursery home during the COVID pandemic in the United States. We use an hourly time step resolution and we ran the model for 150 days.

Population structure

We used the floor plans and satellite imagery to recreate the spatial structure of a typical nursery home in the US (figure 1). The nursery home consist on 58 bedrooms designated for the residents, recreation areas (such as dining room, and activities rooms), and rooms for staff use. In the initial conditions there are 3 residents per room (total 174) and 170 staff divided into 3 different turns. The decision on the population distribution was based on information obtained from an interview with a nursery home in California.

Population dynamics

In our simulation, an agent can interact with other agents based on its location. Given the current guidelines of recommendations for long term care facilities, there are no visitations and the residents spend most of the day in their rooms, so they can only interact with their roommates and the staff. In our model each resident will have at least one interaction with the staff per day which is based on different contact rates depending on the staff type (CN, RN, LPN), The staff will have different contact rates that were parametrized according on the average number of resident contacts in a normal day (REFERENCE: Table shared via email??). The contact rates are presented in table from supplementary materials.

Type of staff	Distribution of resident contacts per turn
CN	$x \sim Mult(n, \pi)$
RN	$x \sim Mult(n, \pi)$
LPN	$x \sim Mult(n, \pi)$

The staff agents are assigned to one of 3 different work schedules (morning, afternoon or night) and they spend 8 hours inside the nursery home and the rest of the time outside in the community. We only follow the agents inside the nursery home and when the agents are outside we assume that they all have the same probability of contacting other people.

Disease dynamics:

The transmission between agents inside the facility will depend on two parts, which are the probability that a person will shed the virus and the probability that another person will get the virus. We decided on model the transmission this way to represent scenarios where the infected and susceptible could have

different combination of interventions (i.e. only infected received the intervention, only susceptible received the intervention, both received the intervention, etc..). The parametrization of the transmission parameters was based on observed outbreaks in nursery homes in California.

The introduction from the community to the facility depends on a parameter *Introduction_p* that represents how likely is that a staff agent will be infected in the community.

All the agents start as susceptible and after 1 day there is a resident introduced with the disease. Then we follow up for 150 days or until the disease has been absent for more than 14 simulation days. Once the transmission between a infectious agent to a susceptible agent has been successful, the susceptible agent becomes exposed and based on a distribution for the latent period λ , the agent becomes infectious after λ number of days, which can be either symptomatic and asymptomatic. The agent can infect other agents only when its in the Infectious state, then they remain infectious during 15 days and they transition to recovered. The agents can transition to infectious to hospitalized at any moment based on the hospitalization rate. When the agents has been recovered they acquire infection immunity, which lasts for ____ days.

Disease parameters:

Name	Value	Reference
Latent period (λ)	<i>Lognormal</i> (7,3) ^b	(He et al. 2020) ^c
Shedding probability	0.4 ^a	
Infection probability	0.4 ^a	
Introduction probability	0.01 ^a	
Asymptomatic probability	0.25 ^a	
Infection duration	15 <i>days</i>	
Hospitalization rate	0.11	

^aExplored for sensitivity analysis and scenario modeling, ^bTruncated distribution between a boundary of reasonable values, ^cFitted to a distribution

Interventions

We explore 3 different

- Testing, we explore different testing strategies based on the frequency and proportion of individuals tested
 - Passive, individuals are tested once that they present symptoms
 - Active, a proportion of individuals are tested with a given frequency. The baseline scenario test
- Isolation
- Use of PPE
- Vaccination

Interventions parameters:

Name	Value	Reference
Test detection probability	75%	
Proportion of Staff tested		
Proportion of Residents tested		
Frequency of testing		
PPE Effect		
Vaccine effect first and second dose	60%,90%	
Vaccine immunity duration	120 <i>days</i> ^b	

Scenario modeling

Results

Sensitivity analysis

Scenario modeling

References

He, Xi, Eric H. Y. Lau, Peng Wu, Xilong Deng, Jian Wang, Xinxin Hao, Yiu Chung Lau, et al. 2020. “Temporal dynamics in viral shedding and transmissibility of COVID-19.” *Nature Medicine* 26 (5): 672–75. <https://doi.org/10.1038/s41591-020-0869-5>.