

# Methods/Results


## Methods

### Population

We present a spatial explicit stochastic agent based model that recreates the day to day dynamics in a typical nursery home during the COVID pandemic in the United States. We use an hourly time step resolution and we ran the model for 150 days.

#### Population structure

We used the floor plans and satellite imagery to recreate the spatial structure of a typical nursery home in the US (figure 1). The nursery home consist on 58 bedrooms designated for the residents, recreation areas (such as dining room, and activities rooms), and rooms for staff use. In the initial conditions there are 3 residents per room (total 174) and 170 staff divided into 3 different turns. The decision on the population distribution was based on information obtained from an interview with a nursery home in California.



## Population dynamics

In our simulation, an agent can interact with other agents based on its location. Given the current guidelines of recommendations for long term care facilities, there are no visitations and the residents spend most of the day in their rooms, so they can only interact with their roommates and the staff. In our model each resident will have at least one interaction with the staff per day which is based on different contact rates depending on the staff type (CN, RN, LPN), The staff will have different contact rates that were parametrized according on the average number of resident contacts in a normal day (REFERENCE: Table shared via email??). The contact rates are presented in table from supplementary materials.

The staff agents are assigned to one of 3 different work schedules (morning, afternoon or night) and they spend 8 hours inside the nursery home and the rest of the time outside in the community. We only follow the agents inside the nursery home and when the agents are outside we assume that they all have the same probability of contacting other people.


## Disease dynamics:

The transmission between agents inside the facility will depend on two parts, which are the probability that a person will shed the virus and the probability that another person will get the virus. We decided

on model the transmission this way to represent scenarios where the infected and susceptible could have different combination of interventions (i.e. only infected received the intervention, only susceptible received the intervention, both received the intervention, etc..). The parametrization of the transmission parameters was based on observed outbreaks in nursery homes in California.

The introduction from the community to the facility depends on a parameter *Introduction\_p* that represents how likely is that a staff agent will be infected in the community.

All the agents start as susceptible and after 1 day there is a resident introduced with the disease. Then we follow up for 150 days or until the disease has been absent for more than 14 simulation days. Once the transmission between a infectious agent to a susceptible agent has been successful, the susceptible agent becomes exposed and based on a distribution for the latent period  $\lambda$ , the agent becomes infectious after  $\lambda$  number of days, which can be either symptomatic and asymptomatic. The agent can infect other agents only when its in the Infectious state, then they remain infectious during 15 days and they transition to recovered. The agents can transition to infectious to hospitalized at any moment based on the hospitalization rate. When the agents has been recovered they acquire infection immunity, which lasts for 120 days.



Transmission parameters:

Name	Value	Reference
Latent period ( $\lambda$ )	<i>Lognormal</i> (7, 3)	(He et al. 2020) <sup>b,c</sup>
Shedding probability	0.5	<sup>a</sup>
Infection probability	0.5	<sup>a</sup>
Introduction probability	0.01	<sup>a</sup>
Asymptomatic probability	0.25	<sup>a</sup>
Infection duration	15 <i>days</i>	
Hospitalization rate	0.11	

<sup>a</sup>Explored for sensitivity analysis and scenario modeling, <sup>b</sup>truncated distribution between a boundary of reasonable values, <sup>c</sup>fitted to a distribution

## Interventions

We explore 3 different COVID-19 control strategies and the combination of them. Each of the interventions have an impact in the transmission of the disease, interventions such as the use of PPE and vaccination reduces the probability of transmission affecting directly the *Shedding* and *Infection probability*, while the isolation affects the transmission indirectly stopping the agent to interact with other agents. The equation 1 shows the effect of *PPE effect* and *Vaccine effect* on the transmission probability, where  $odds_{\omega}$  represent the global transmission probability for all agents,  $OR_{\gamma}$  represent the odds ratio for the *PPE effect*, and  $OR_{\pi}$  is the *Vaccine effect*. This probability is computed for all agents at each step so we can have different probabilities of transmission based on the interventions each individual received.

$$p_T = \frac{e^{\ln(odds_{\omega}) + \ln(OR_{\pi}) + \ln(OR_v)}}{1 + e^{\ln(odds_{\omega}) + \ln(OR_{\pi}) + \ln(OR_v)}}$$

For the implementation of the vaccination, we specified by the proportion of residents and staff vaccinated, and a fixed interval between the first and second dose of 21 days. After the first dose, the agents will only obtain a 60% of the total immunity protection assumed to be conferred by the vaccine, then on the second dose the agents will have 100% of the assumed effect. Then the vaccination immunity will have a decay of 120 days and the individual will no longer have the vaccination immunity protective effect.

Since there is still some uncertainty in the effect of the use of PPE and the vaccine for older population, we started with values that are within the range of reported values and then varied these values for the sensitivity analysis and scenario modeling.

## Testing and isolation

Our model represents the testing of the population with 2 different approaches:

- Passive, individuals are tested once that they present symptoms, this approach is focused on the early detection of symptomatic individuals.
- Active, a proportion of individuals are tested with a given frequency. In baseline scenario, 1 resident per room and all the staff are tested weekly. If 1 of the residents in a room is detected positive, the rest of the residents in that room are also tested.

Once a individual has been detected positive is isolated. There are special isolation rooms for the residents and in the case of the staff they are sent home. Once the individual is tested negative it return to the facility.

Interventions parameters:

Name	Value	Reference
Test detection probability	85%	<sup>a</sup>
Proportion of Staff tested	90%	<sup>a</sup>
Proportion of Residents tested	33.3	<sup>a</sup>
Frequency of testing	Weekly	<sup>a</sup>
PPE Effect ( $OR_{\pi}$ )	0.34089	(Chu et al. 2020) <sup>a</sup>
Vaccine effect first and second dose ( $OR_v$ )	60%,90%	(Pfizer-BioNTech 2020) <sup>a</sup>
Vaccine immunity duration	120 <i>days</i>	<sup>a</sup>

<sup>a</sup>Explored for sensitivity analysis and scenario modeling, <sup>b</sup>truncated distribution between a boundary of reasonable values, <sup>c</sup>fitted to a distribution

## Scenario modeling

**Vaccine Effect:** To explore the vaccination implementation we considered 3 scenarios according to the current vaccine effect reported by the trials from the Moderna and Pfizer vaccine (Baden et al. 2020; Polack et al. 2020). Since there is still some uncertainty about the effect of the vaccine in population  $>65$  years, we defined 3 scenarios that explore the possible outcomes under 3 different assumptions:

- Scenario 00: The vaccine has the same effect both populations ( $>65$  years and  $>65$  years).
- Scenario 01: The vaccine is less effective on populations  $>65$  years parametrized according to the reported by (Baden et al. 2020).
- Scenario 02: The vaccine is less effective on populations  $>65$  years parametrized according to the reported by (Polack et al. 2020).


The parameters used for the vaccine effect in the 3 scenarios are the following:

Scenario	$OR_{V,S}$	$OR_{V,R}$	$V_S$	$V_R$
S00	0.0493	0.0493	80%	20%
S01	0.0434	0.0619	80%	20%
S02	0.0441	0.1357	80%	20%

## Vaccine Distribution:

Scenario	$OR_{V,S}$	$OR_{V,R}$	$V_S$	$V_R$
S00	0.0441	0.1357	50%	50%
S01	0.0441	0.1357	80%	20%
S02	0.0441	0.1357	20%	80%

$T_f$  Frequency of testing,  $T_r$  Residents tested,  $T_s$  Staff tested,  $V_e$  Vaccination efficacy,  $V_r$  Residents vaccinated,  $V_s$  Staff vaccinated



## Results



### Sensitivity analysis

We performed sensitivity analysis on some parameters to explore the influence on the outbreak size and duration.

- Disease Parameters:
  - Transmission probability (for both shedding and infection)
  - Introduction probability (low risk, medium risk, high risk)
- Interventions
  - Frequency of testing (1 week, 5 days, 3 days)
  - Detection probability
  - PPE effect
  - Vaccine effect
  - Vaccine immunity decay



## Transmission rate

Scenario ■ Low transmission ■ Medium transmission ■ High transmission








## Probability of Introduction

Scenario ■ 1% Introduction Probability ■ 5% Introduction Probability ■ 10% Introduction Probability





## Probability of Detection

Scenario  75% Detection Probability  85% Detection Probability  95% Detection Probability




## Effect of PPE

Scenario  Highly effective  Medium effective  Low effective







## Scenario modeling



With 5% probability of introduction:



with 1% probability of introduction:



## References

- Baden, Lindsey R., Hana M. El Sahly, Brandon Essink, Karen Kotloff, Sharon Frey, Rick Novak, David Diemert, et al. 2020. "Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine." *New England Journal of Medicine*, December, NEJMoa2035389. <https://doi.org/10.1056/NEJMoa2035389>.
- Chu, Derek K., Elie A. Akl, Stephanie Duda, Karla Solo, Sally Yaacoub, Holger J. Schünemann, Amena El-harakeh, et al. 2020. "Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis." *The Lancet* 395 (10242): 1973–87. [https://doi.org/10.1016/S0140-6736\(20\)31142-9](https://doi.org/10.1016/S0140-6736(20)31142-9).
- He, Xi, Eric H. Y. Lau, Peng Wu, Xilong Deng, Jian Wang, Xinxin Hao, Yiu Chung Lau, et al. 2020. "Temporal dynamics in viral shedding and transmissibility of COVID-19." *Nature Medicine* 26 (5): 672–75. <https://doi.org/10.1038/s41591-020-0869-5>.
- Pfizer-BioNTech. 2020. "Vaccines and Related Biological Products Advisory Committee Meeting December 10, 2020." Pfizer-BioNTech.
- Polack, Fernando P, Stephen J Thomas, Nicholas Kitchin, Judith Absalon, Alejandra Gurtman, Stephen Lockhart, John L Perez, et al. 2020. "Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine." *The New England Journal of Medicine*, 2603–15. <https://doi.org/10.1056/NEJMoa2034577>.