

Foreign Domestic Workers' Perspectives on an LLM-Based Emotional Support Tool for Caregiving Burden

Shin Shoon Nicholas Teng
Singapore University of Technology and Design
Singapore, Singapore
nicholastengshin@hotmail.com

Kenny Tsu Wei Choo
Singapore University of Technology and Design
Singapore, Singapore
kenny_choo@sutd.edu.sg

Abstract

Foreign Domestic Workers (FDWs) play a central role in home-based eldercare yet often experience substantial emotional caregiving burden shaped by linguistic barriers, social isolation, and limited access to support. While caregiving burden has been extensively studied among familial caregivers, little is known about how FDWs engage with emotional support technologies. We present an exploratory qualitative study of how FDWs in Singapore interact with a Large Language Model (LLM)-driven chatbot as an everyday, non-clinical form of emotional support. Through interviews and guided chatbot interactions, we conducted an inductive thematic analysis of participants' experiences. We identify three design-relevant themes: chatbots were experienced as psychologically safe and emotionally validating; they supported linguistic accessibility by accommodating imperfect and fragmented language; and they were appropriated as multifunctional resources for reassurance, guidance, and companionship. We discuss implications for designing LLM-driven emotional support tools that foreground psychological safety, accessibility, and flexible appropriation.

CCS Concepts

• **Human-centered computing** → **Empirical studies in HCI**; *Caregiving and emotional support applications*.

Keywords

Human-AI interaction, Caregiving support, Foreign Domestic Workers, Natural language processing, Conversational agents, Emotional well-being

ACM Reference Format:

Shin Shoon Nicholas Teng and Kenny Tsu Wei Choo. 2026. Foreign Domestic Workers' Perspectives on an LLM-Based Emotional Support Tool for Caregiving Burden. In *Extended Abstracts of the 2026 CHI Conference on Human Factors in Computing Systems (CHI EA '26)*, April 13–17, 2026, Barcelona, Spain. ACM, New York, NY, USA, 5 pages. <https://doi.org/10.1145/3772363.3798761>

1 Introduction

Foreign Domestic Workers (FDWs) play a central role in supporting home-based eldercare in ageing societies such as Singapore [6, 24]. In many households, FDWs provide day-to-day assistance for frail older adults, managing both the practical and emotional demands

of caregiving [23]. Despite this central role, FDWs often experience substantial emotional caregiving burden shaped by linguistic barriers, social isolation, and structural constraints associated with migrant domestic work [10, 24]. These conditions can restrict opportunities for emotional expression, limit access to social support, and intensify feelings of stress, uncertainty, and loss of control in everyday caregiving situations.

While caregiving burden has traditionally been examined through the lens of familial caregivers such as spouses or adult children [15, 18, 25], this body of work implicitly marginalizes non-familial caregivers like FDWs, whose caregiving labor is situated within employment relationships rather than kinship ties. As a result, little is known about how FDWs emotionally appraise caregiving stress or seek support in everyday caregiving contexts where formal resources are limited or inaccessible.

Recent advances in Large Language Model (LLM)-driven chatbots present a potential form of everyday, non-clinical emotional support that may align with FDWs' constraints. Chatbots offer private, low-cost, readily accessible interaction and have demonstrated promise in providing emotional support and fostering a sense of being heard through empathetic and reflective dialogue [8, 14]. For FDWs, who may face stigma, social isolation, and linguistic barriers when seeking help [12, 20], LLM-driven chatbots may offer a distinct form of emotional support that is difficult to access through existing social or institutional channels.

While prior research has examined the use of chatbots for mental health and caregiving support more broadly, there is limited understanding of how FDWs themselves perceive and engage with LLM-driven chatbots, and how these experiences can inform the design of emotional support technologies. In particular, it remains unclear how FDWs experience these systems as conversational partners, how linguistic and cultural factors shape interaction, and what expectations FDWs form regarding the role such tools might play in their caregiving lives.

To address this gap, we conducted an exploratory qualitative study with seven FDWs working in Singapore to examine how they interact with an LLM-driven chatbot when reflecting on emotionally challenging caregiving experiences. Specifically, this study seeks to understand how FDWs interact with LLM-driven chatbots in relation to managing their emotional caregiving burden. Therefore, we investigated the following research question:

"How might FDWs in Singapore interact with LLM-driven chatbots in relation to managing their caregiving burden?"

By foregrounding FDWs' perspectives, this study contributes qualitative insights into how FDWs experience LLM-driven emotional



This work is licensed under a Creative Commons Attribution 4.0 International License. *CHI EA '26, Barcelona, Spain*

© 2026 Copyright held by the owner/author(s).

ACM ISBN 979-8-4007-2281-3/26/04

<https://doi.org/10.1145/3772363.3798761>

support in everyday contexts, and surfaces design implications for such systems that aim to support an FDW's emotional well-being.

2 Related Work

2.1 Caregiving Burden and Emotional Appraisal

Caregiving burden has been widely conceptualized as a subjective and multidimensional experience shaped not only by the objective demands of care, but by caregivers' emotional appraisal of their caregiving role. Pearlin's stress process model [18] frames caregiving burden as arising from the interaction between caregiving-related stressors and the availability of social and emotional support. In this view, caregiving burden intensifies not only as stressors accumulate, but also when caregivers lack access to emotional, social, or communicative forms of support that enable them to cope with and make sense of these demands.

For FDWs, these dynamics are often amplified by structural and communicative constraints associated with migrant domestic work. Long working hours, limited rest days [17], and caregiving in a non-native language [10] restrict opportunities for emotional expression and access to social support. Caregiving occurs within employment relationships rather than kinship ties, where expectations of professionalism and power asymmetries can further limit safe emotional disclosure [23]. As a result, FDWs may experience heightened emotional caregiving burden even when the objective demands of care are comparable to those faced by familial caregivers. Despite these insights, existing caregiving frameworks offer limited guidance on how FDWs access emotionally validating or expressive support in their everyday caregiving practices.

2.2 Technologies for Caregiving and Emotional Support

Prior HCI research has examined technologies that support caregivers' everyday practices, including role management, experiential knowledge sharing, and the emotional labor of care [13, 19, 21]. Beyond familial caregiving contexts, CSCW scholarship has explored how domestic workers engage with digital technologies under conditions of labor precarity, where adoption is shaped by trust, autonomy, and perceived risk [7]. Together, this body of work shows that caregiving technologies mediate not only coordination and emotional labor, but also the broader socio-structural constraints shaping how FDWs engage with conversational systems.

In parallel, advances in Large Language Model (LLM)-driven conversational agents have explored chatbots as tools for providing empathetic and emotionally responsive interaction [4]. Prior studies suggest that chatbots can encourage emotional disclosure and foster a sense of being heard, particularly in contexts where users fear judgment or lack access to human support [5, 11]. However, this work has largely focused on clinical mental health settings or system effectiveness, with limited attention to how caregivers experience and appropriate LLM-driven emotional support in everyday, non-clinical contexts

For FDWs in particular, it remains unclear how LLM-driven chatbots are experienced as conversational partners under conditions of linguistic constraint, how tolerance for imperfect or fragmented language shapes access to support, and how such systems may be flexibly appropriated to address caregiving-related uncertainty,

isolation, and emotional strain. Addressing these gaps requires attending to FDWs' lived experiences of interaction with LLM-driven systems, rather than evaluating chatbot performance or outcomes alone.

3 Methodology

3.1 Study Design and Qualitative Data Collection

We conducted a single-session exploratory qualitative study to examine how FDWs engage with an LLM-driven chatbot, *Sophia*, when reflecting on emotionally challenging caregiving experiences. This format was adopted to minimize demands on participants' limited free time and to capture initial, situated impressions of the chatbot within an ethically appropriate research setting. Participants were FDWs working in Singapore with experience in eldercare and basic English proficiency. Seven FDWs (all female), aged 23 to 50, were recruited through in-person intercept recruitment in public spaces frequented by FDWs and via snowball sampling. These participants were selected as they completed the 12-item short-form Zarit Burden Interview (ZBI-12) [1], with a median score of 14, indicating overall mild caregiver burden.

Each participant took part in one in-person session lasting approximately 1 to 1.5 hours. Sessions included a semi-structured interview guided by the Critical Incident Technique [3, 9]. Based on the recalled caregiving incident, participants then engaged in a guided interaction with *Sophia*, which was configured to provide empathetic, non-clinical emotional support. *Sophia* operated under a single, fixed system prompt that framed it as a non-judgmental conversational partner, encouraged emotional articulation through open-ended questions, and explicitly declined to provide medical, legal, or financial advice. The restriction was implemented to mitigate ethical risks and to ensure the system remained within a non-clinical, emotionally supportive scope. During the interaction, participants were encouraged to verbalize their thoughts using the think-aloud method [22]. Sessions concluded with a brief reflective interview about *Sophia*'s perceived usefulness and limitations. All sessions were audio-recorded and transcribed, and informed consent was obtained in accordance with IRB-approved procedures. The research team consisted of university-based researchers trained in HCI and qualitative methods. As we do not share the lived experience of FDWs, we remained attentive to power dynamics during recruitment, interviews, and analysis.

3.2 Data Analysis

We analyzed the data using inductive thematic analysis [2]. Audio recordings were transcribed verbatim and iteratively coded to identify patterns related to participants' caregiving experiences and interactions with *Sophia*. Initial codes were compared across participants and refined into higher-level themes through repeated reference to the transcripts to ensure grounding in participants' accounts. All participants were assigned unique identifiers to preserve anonymity.

4 Results

We identified three themes characterizing how FDWs experienced and engaged with *Sophia* in relation to caregiving burden: (1) psychologically safe and emotionally validating interaction, (2) linguistic accessibility through tolerance of imperfect language, and (3) multifunctional appropriation beyond emotional support.

4.1 Psychologically Safe and Emotionally Validating Interaction

Participants commonly experienced *Sophia* as a psychologically safe and emotionally validating space for reflecting on caregiving challenges. Several participants (n=2) described disclosing thoughts and emotions to *Sophia* that they would not share with others, emphasizing the absence of social judgment or evaluation. As one participant explained, “Yes, I can talk to it about anything... I don’t want to say to other people but I want to say what is on my mind. So talk with AI is okay, outside other people no one can know better.” (P1).

Nearly all participants (n=5) described *Sophia*’s responses as reassuring and stress-reducing, helping them continue caregiving tasks. Some participants noted that these interactions helped reduce stress and gave them emotional strength to continue caregiving tasks. Notably, participants emphasized that emotional acknowledgment mattered more than cultural specificity or conversational naturalness, suggesting that perceived validation played a central role in shaping positive experiences.

4.2 LLM-driven Chatbots Support Linguistic Accessibility by Accepting Imperfect Language

Linguistic accessibility emerged as a key factor shaping participants’ engagement with *Sophia*. While some participants (n=3) initially found it challenging to formulate detailed prompts in English, they reported greater comfort interacting in their native languages, which allowed them to express emotions more naturally and with less effort.

At the same time, more than half of the participants (n=4) described being able to communicate effectively in English despite limited proficiency. Participants frequently used short, fragmented, or grammatically non-standard input, yet perceived *Sophia*’s responses as emotionally supportive and relevant. Several expressed surprise that *Sophia* could infer meaning from minimal or imperfect input. As one participant noted, “Because we’re just talking a little bit, but it can reply a long long thing that is correct.” (P3). This tolerance for imperfect language reduced pressure to perform linguistic competence and lowered the effort required to seek support. However, linguistic accessibility was not uniform. One participant reported difficulty understanding specific English vocabulary in *Sophia*’s responses, indicating that accessibility depended not only on accommodating user input but also on generating comprehensible responses.

4.3 LLM Chatbots as Multifunctional Support Resources in Caregiving

Beyond emotional disclosure, participants appropriated *Sophia* as a multifunctional support resource, using it to seek reassurance, cope with loneliness, practice English, and navigate caregiving-related uncertainties (n=4). These uses were not framed as separate from caregiving but as indirectly supporting caregiving by reducing uncertainty, isolation, and emotional strain. One participant described relaying on *Sophia* when other supports were unavailable: “Yeah I’m happy la, for me it’s helping me so much much la...so better to have the app to ask. If I ask the employer, she also don’t know what to do” (P5). Participants described turning to *Sophia* when social, workplace, or informational supports were insufficient, highlighting its role as a flexible and accessible resource within constrained caregiving contexts.

5 Discussion

5.1 Interpreting FDW’s Experiences with LLM-Driven Emotional Support

Our findings suggest that FDWs’ interactions with *Sophia* supported caregiving burden primarily through expressive emotional support. Participants described feeling heard and validated, aligning with Pearlin’s stress process model [18], which positions emotional acknowledgment as a key resource when stressors cannot be reduced. For FDWs operating within employment-based care relationships, such support is often constrained by professional norms and power asymmetries. In this context, *Sophia* functioned as a psychologically safe space for disclosure without social or occupational risk.

Participants valued non-judgmental validation over cultural specificity or conversational naturalness, suggesting that emotional support technologies may be most effective when prioritizing psychological safety over surface-level personalization. However, this safety is structurally distinct from reciprocal human support. Unlike personal networks, *Sophia* entails no mutual accountability or relational negotiation. While this asymmetry may lower barriers to disclosure, it reframes emotional support as a one-directional rather than relational exchange.

5.2 Linguistic Accessibility as a Condition for Emotional Support

Our findings highlight linguistic accessibility as a key condition enabling emotional support. Many participants interacted using fragmented or non-standard language, yet still experienced *Sophia*’s responses as emotionally supportive and meaningful. This tolerance for imperfect input reduced the effort required to articulate distress and lowered barriers to seeking support across language differences.

These findings suggest that linguistic accessibility is not merely a usability concern but is closely tied to emotional access. For migrant caregivers operating in non-native languages, the ability to communicate without performing linguistic competence can shape whether emotional support feels approachable or burdensome. At the same time, difficulties understanding certain chatbot responses

indicate that accessibility depends on both input tolerance and output clarity. Designing for linguistic accessibility therefore requires attention to how conversational systems accommodate limited proficiency while generating responses that remain emotionally and cognitively accessible.

5.3 LLM Chatbots as Flexible Support Resources in Caregiving Contexts

Participants' appropriated *Sophia* as a multifunctional resource extending beyond emotional disclosure, using it to seek reassurance, cope with loneliness, and navigate caregiving-related uncertainty when other supports were unavailable. These uses indirectly supported caregiving by addressing isolation and informational gaps.

This flexible appropriation reflects prior work showing that caregiving burden arises from a combination of emotional, relational, and communicative stressors [10, 18]. Rather than replacing existing support networks, *Sophia* appeared to fill gaps created by linguistic barriers, social hierarchies, and limited access to caregiving knowledge. These findings suggest that emotional support technologies for caregivers may be most valuable when designed as adaptable, non-judgmental resources that caregivers can appropriate according to their lived constraints and evolving needs.

5.4 Design Implications for Emotional Support Technologies for FDWs

Taken together, our findings point to several implications for the design of LLM-driven emotional support tools for FDWs. First, designers should foreground psychological safety and emotional validation as core interactional properties, enabling caregivers to express distress without fear of judgment or professional failure. Second, linguistic accessibility should be prioritized over linguistic correctness, accommodating imperfect input while generating clear and emotionally accessible responses. Finally, emotional support technologies should support flexible, multifunctional appropriation, recognizing that caregiving burden is shaped by broader everyday stressors beyond emotional disclosure alone. Designing conversational systems as adaptable support resources may better align with FDWs' lived realities and caregiving contexts.

5.5 Ethical and Temporal Considerations

Although participants experienced *Sophia* as psychologically safe within the session, emotionally responsive systems introduce a form of bounded emotional engagement. Emotional disclosure may extend beyond the temporal limits of a study interaction, raising questions about containment and continuity. Designing such systems therefore requires attention not only to validation and accessibility, but also to how emotional support is introduced, framed, and withdrawn. While LLM-driven emotional support may provide expressive relief, it does not address the structural labour conditions that give rise to caregiving burden. There is a risk that positioning such systems as solutions may inadvertently individualize responsibility for coping, shifting attention away from broader socio-economic inequities shaping FDWs' work. Emotional support technologies should therefore be understood as supplementary resources rather than substitutes for structural change.

5.6 Limitations and Future Work

This study has several limitations. First, participants' interactions with *Sophia* occurred within a single, time-limited session, capturing initial impressions rather than longer-term patterns of use, trust, or reliance that may emerge over time. Second, the small sample size and focus on FDWs working in Singapore limit the generalizability of our findings to other caregiving contexts or migrant worker populations. Finally, participants engaged with a single chatbot configuration, and different prompt designs or system behaviors may shape caregiving experiences in other ways.

Future work could adopt longitudinal field studies to examine how FDWs integrate LLM-driven emotional support tools into everyday caregiving routines over time, and how such systems influence emotional well-being, coping practices, and caregiving burden. Exploring alternative designs grounded in caregiving and self-care frameworks (e.g., Orem's Self-Care Deficit Nursing Theory [16]) may further inform the development of emotionally supportive technologies for FDWs.

6 Conclusion

This paper examined how FDWs in Singapore engage with an LLM-driven chatbot as a form of everyday emotional support in caregiving contexts. Our findings show that FDWs experienced *Sophia* as a psychologically safe and emotionally validating space, valued its tolerance for imperfect and non-standard language, and appropriated it as a flexible support resource addressing caregiving-related uncertainty and isolation. Together, these insights highlight the potential of LLM-driven conversational systems to support caregivers' emotional well-being under conditions of social, linguistic, and structural constraint. We conclude by emphasizing the importance of designing emotional support technologies that foreground psychological safety, linguistic accessibility, and flexible appropriation to better align with the lived realities of FDWs and other marginalized caregivers.

References

- [1] Martin Bédard, David W. Molloy, Larry Squire, Sylvie Dubois, Jeffrey A. Lever, and Mark O'Donnell. 2001. The Zarit Burden Interview: A New Short Version and Screening Version. *The Gerontologist* 41, 5 (Oct. 2001), 652–657. doi:10.1093/geront/41.5.652
- [2] Virginia Braun and Victoria Clarke. 2006. Using thematic analysis in psychology. *Qualitative Research in Psychology* 3, 2 (2006), 77–101. arXiv:https://doi.org/10.1191/1478088706qp0630a doi:10.1191/1478088706qp0630a
- [3] Lee D. Butterfield, William A. Borgen, Norman E. Amundson, and Asa-Sophia T. Maglio. 2005. Fifty years of the critical incident technique: 1954-2004 and beyond. *Qualitative Research* 5, 4 (2005), 475–497. arXiv:https://doi.org/10.1177/1468794105056924 doi:10.1177/1468794105056924
- [4] Mirko Casu, Sergio Triscari, Sebastiano Battiato, Luca Guarnera, and Pasquale Caponnetto. 2024. AI Chatbots for Mental Health: A Scoping Review of Effectiveness, Feasibility, and Applications. *Applied Sciences* 14, 13 (2024). doi:10.3390/app14135889
- [5] Ana Paula Chaves and Marco Aurélio Gerosa. 2019. How should my chatbot interact? A survey on human-chatbot interaction design. *CoRR* abs/1904.02743 (2019). arXiv:1904.02743 http://arxiv.org/abs/1904.02743
- [6] Theresa W. Devasahayam. 2010. Placement and/or protection? Singapore's labour policies and practices for temporary women migrant workers. *Journal of the Asia Pacific Economy* 15, 1 (2010), 45–58. arXiv:https://doi.org/10.1080/13547860903488229 doi:10.1080/13547860903488229
- [7] Mariana Fernandez-Espinosa, Mariana Gonzalez-Bejar, Jacobo Wiesner, and Diego Gómez-Zarzá. 2025. When Technologies Are Not Enough: Understanding How Domestic Workers Employ (and Avoid) Online Technologies in Their Work Practices. *Proc. ACM Hum.-Comput. Interact.* 9, 7, Article CSCW342 (Oct. 2025), 34 pages. doi:10.1145/3757523

- [8] Kathleen Kara Fitzpatrick, Alison Darcy, and Molly Vierhile. 2017. Delivering Cognitive Behavior Therapy to Young Adults With Symptoms of Depression and Anxiety Using a Fully Automated Conversational Agent (Woebot): A Randomized Controlled Trial. *JMIR Ment Health* 4, 2 (06 Jun 2017), e19. doi:10.2196/mental.7785
- [9] John C. Flanagan. 1954. The Critical Incident Technique. *Psychological Bulletin* 51, 4 (1954), 327–358. doi:10.1037/h0061470
- [10] Ngoc Huong Lien Ha, Mei Sian Chong, Robin Wai Munn Choo, Wai Jia Tam, and Philip Lin Kiat Yap. 2018. Caregiving burden in foreign domestic workers caring for frail older adults in Singapore. *International Psychogeriatrics* 30, 8 (August 2018), 1139–1147. doi:10.1017/S1041610218000200
- [11] Annabell Ho, Jeff Hancock, and Adam S Miner. 2018. Psychological, Relational, and Emotional Effects of Self-Disclosure After Conversations With a Chatbot. *Journal of Communication* 68, 4 (05 2018), 712–733. arXiv:https://academic.oup.com/joc/article-pdf/68/4/712/25410448/jqy026.pdf doi:10.1093/joc/jqy026
- [12] Ken Hok Man Ho, Chen Yang, Alex Kwun Yat Leung, Daniel Bressington, Wai Tong Chien, Qijin Cheng, and Daphne Sze Ki Cheung. 2022. Peer Support and Mental Health of Migrant Domestic Workers: A Scoping Review. *International Journal of Environmental Research and Public Health* 19, 13 (June 2022), 7617. doi:10.3390/ijerph19137617
- [13] Long-Jing Hsu and Chia-Fang Chung. 2024. Dancing with the Roles: Towards Designing Technology that Supports the Multifaceted Roles of Caregivers for Older Adults. In *Proceedings of the 2024 CHI Conference on Human Factors in Computing Systems* (Honolulu, HI, USA) (CHI '24). Association for Computing Machinery, New York, NY, USA, Article 1010, 12 pages. doi:10.1145/3613904.3642728
- [14] Becky Inkster, Shubhankar Sarda, and Vinod Subramanian. 2018. An Empathy-Driven, Conversational Artificial Intelligence Agent (Wysa) for Digital Mental Well-Being: Real-World Data Evaluation Mixed-Methods Study. *JMIR Mhealth Uhealth* 6, 11 (23 Nov 2018), e12106. doi:10.2196/12106
- [15] M. Powell Lawton, Michael H. Kleban, Miriam Moss, Michael Rovine, and Allen Glicksman. 1989. Measuring caregiving appraisal. *Journal of Gerontology* 44, 3 (May 1989), P61–P71. doi:10.1093/geronj/44.3.p61
- [16] Dorothea E. Orem. 2001. *Nursing: Concepts of Practice* (6 ed.). Mosby, St. Louis, MO.
- [17] Rhacel Salazar Parreñas, Krittiya Kantachote, and Rachel Silvey. 2021. Soft violence: migrant domestic worker precarity and the management of unfree labour in Singapore. *Journal of Ethnic and Migration Studies* 47, 20 (2021), 4671–4687. arXiv:https://doi.org/10.1080/1369183X.2020.1732614 doi:10.1080/1369183X.2020.1732614
- [18] Leonard I. Pearlin, Joseph T. Mullan, Shirley J. Semple, and Marilyn M. Skaff. 1990. Caregiving and the stress process: an overview of concepts and their measures. *The Gerontologist* 30, 5 (October 1990), 583–594. doi:10.1093/geront/30.5.583
- [19] Ji Youn Shin, Dima Chaar, Catherine Davis, Sung Won Choi, and Hee Rin Lee. 2021. Every Cloud Has a Silver Lining: Exploring Experiential Knowledge and Assets of Family Caregivers. *Proc. ACM Hum.-Comput. Interact.* 5, CSCW2, Article 416 (Oct. 2021), 25 pages. doi:10.1145/3479560
- [20] Slesh A. Shrestha and Dean Yang. 2019. Facilitating Worker Mobility: A Randomized Information Intervention among Migrant Workers in Singapore. *Economic Development and Cultural Change* 68, 1 (October 2019), 63–91. doi:10.1086/700620
- [21] Diva Smriti, Lu Wang, and Jina Huh-Yoo. 2024. Emotion Work in Caregiving: The Role of Technology to Support Informal Caregivers of Persons Living With Dementia. *Proc. ACM Hum.-Comput. Interact.* 8, CSCW1, Article 48 (April 2024), 34 pages. doi:10.1145/3637325
- [22] Maarten Someren, Yvonne Barnard, and Jacobijn Sandberg. 1994. *The Think Aloud Method - A Practical Guide to Modelling Cognitive Processes*.
- [23] Wai Jia Tam, Gerald Choon-Huat Koh, Helena Legido-Quigley, Ngoc Huong Lien Ha, and Philip Lin Kiat Yap. 2018. “I Can’t Do This Alone”: a study on foreign domestic workers providing long-term care for frail seniors at home. *International Psychogeriatrics* 30, 9 (2018), 1269–1277. doi:10.1017/S1041610217002459 Issue Theme: Quality of Care for Frail Older Adults.
- [24] Brenda S. A. Yeoh and Shirlena Huang. 2010. Transnational Domestic Workers and the Negotiation of Mobility and Work Practices in Singapore’s Home-Spaces. *Mobilities* 5, 2 (2010), 219–236. arXiv:https://doi.org/10.1080/17450101003665036 doi:10.1080/17450101003665036
- [25] Steven H. Zarit, Kathleen E. Reeve, and Joan Bach-Peterson. 1980. Relatives of the impaired elderly: correlates of feelings of burden. *The Gerontologist* 20, 6 (December 1980), 649–655. doi:10.1093/geront/20.6.649