

SoulSkipper: A Voice-Controlled Emotional Adaptive Game to Complement Therapy for Social Anxiety Disorder

Xiaohan Li
Department of Computer Science and
Technology, Tsinghua University
Beijing, China
lixiaoha21@mails.tsinghua.edu.cn

Qixin Wang
Department of Computer Science and
Technology, Tsinghua University
Beijing, China
wqx22@mails.tsinghua.edu.cn

Zishan Wang
School of Software, Tsinghua
University
Beijing, China
wang-zs21@mails.tsinghua.edu.cn

Zeyu Jin
Department of Computer Science and
Technology, Tsinghua University
Beijing, China
jinzeyu23@mails.tsinghua.edu.cn

Jia Jia*
Department of Computer Science and
Technology, Tsinghua University,
Beijing National Research Center for
Information Science and Technology
Beijing, China
jjia@mail.tsinghua.edu.cn

ABSTRACT

In this paper, we introduce SoulSkipper, a voice-controlled emotional adaptive game based on the use of Cognitive Behavioral Therapy (CBT) to complement therapy for social anxiety disorder (SAD). The game allows players to interact with non-player characters (NPCs) through voice input, which helps players identify negative thoughts and practice speech performance through 1) specific social scenarios, 2) emotional adaptation, and 3) real-time voice feedback. Through our initial evaluation, we find that the combination of game elements and voice interactions helps players improve their awareness of therapeutic skills associated with SAD and face social interactions in real-life scenarios more positively.

CCS CONCEPTS

• **Human-centered computing** → **Interaction design; HCI design and evaluation methods.**

KEYWORDS

social anxiety disorder; voice-controlled game; emotional adaptation; CBT

ACM Reference Format:

Xiaohan Li, Qixin Wang, Zishan Wang, Zeyu Jin, and Jia Jia. 2024. SoulSkipper: A Voice-Controlled Emotional Adaptive Game to Complement Therapy for Social Anxiety Disorder. In *Extended Abstracts of the CHI Conference on Human Factors in Computing Systems (CHI EA '24)*, May 11–16, 2024, Honolulu, HI, USA. ACM, New York, NY, USA, 7 pages. <https://doi.org/10.1145/3613905.3650822>

*Corresponding author

Permission to make digital or hard copies of part or all of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for third-party components of this work must be honored. For all other uses, contact the owner/author(s).
CHI EA '24, May 11–16, 2024, Honolulu, HI, USA
© 2024 Copyright held by the owner/author(s).
ACM ISBN 979-8-4007-0331-7/24/05
<https://doi.org/10.1145/3613905.3650822>

1 INTRODUCTION

Social Anxiety Disorder (SAD) is a common and widespread anxiety disorder. Patients typically experience fear and anxiety in one or more specific social scenarios and exhibit physical symptoms such as poor speech and increased heart rate [62]. This state of fear in social interactions can have multiple adverse consequences for patients, leading to disruption of social relationships, loss of confidence in life, and increased risk of suicide [14].

Cognitive Behavioral Therapy (CBT) is the main evidence-based treatment for SAD [18, 40, 50], in which psychotherapists face-to-face contact with patients to help them identify negative thoughts and face their fears by exposing them to specific social scenarios [45]. However, psychotherapy for SAD faces multiple barriers, including a lack of medical resources and privacy concerns, which make psychotherapy less accessible in clinical settings. In addition, the fear of social stigmatization among people with SAD makes it more difficult for them to seek professional therapeutic help [2, 64].

In this context, researchers have explored several digital interventions to improve SAD, thus providing anonymity and accessibility to therapy [6, 41]. Findings suggest that a range of digital interventions was not different from traditional face-to-face psychotherapy in terms of effectiveness [11, 21], however, they have shown their poor engagement and high attrition rates overtime [25]. Research has shown that engagement is associated with improved outcomes for digital interventions [59]. Therefore, there is a need for further research into the user experience and engagement of digital interventions to implement them more effectively among people with SAD, thus filling a gap that is not present in traditional therapies [25].

Serious games and gamification are considered as an attractive solution to increase user engagement [39]. Games and especially fun interactions may be more engaging than non-game digital interventions [25]. However, many current serious games for mental health are not evidence-based and have limitations in their design approach [31]. Moreover, research suggests that existing digital mental health interventions generally neglect human-computer interaction which negatively affects user engagement [5].

To further address these challenges, in this paper, we present a role-playing game based on voice-controlled and emotional adaptation to complement therapy for SAD. We adopt an evidence-based design approach using CBT to help SAD patients challenge negative thoughts and improve behaviors. Considering the speech symptoms of SAD, the game allows players to interact with non-player characters (NPCs) through voice input. It enhances user engagement by increasing the level of personalization of the game through 1) specific social scenarios, 2) emotional adaptation, and 3) real-time voice feedback.

2 GAME GOALS

Our game goals are as follows: **1)** Propose an innovative digital game to increase engagement with SAD patients to complement therapy. **2)** Support self-help CBT training through specific social scenarios that create a game place for players to engage in close to real-life situations, helping them to identify negative thoughts and learn strategies for coming up with positive thoughts. **3)** Combine emotional adaptation with timely feedback to help players reflect and adjust, and face social interactions in real-life scenarios more positively.

3 GAME DESIGN

We refer to the player/game/therapy model (P/G/T model) [39] for game design, this framework describes the game design as a two-step process. We began by independently analyzing all aspects of the game, including understanding the SAD players we were targeting, selecting the therapeutic elements to be embedded in the gameplay process, and defining the underlying genre of the game. We then combined the understanding of the player/game/therapy to generate our game.

3.1 Player Research

Our target players are adults with mild social anxiety disorder. First, we want to answer a research question: What are the anxiety conditions experienced by these SAD patients? To answer this question, we first collected and analyzed posts in online forums discussing social anxiety. Secondly, we organized a semi-structured interview with SAD patients (N=10).

We collected data from the Chinese online forum Douban¹, which allows members to create rich online groups for discussion. We obtained 15 online groups based on keywords such as "social" and "social phobia", and further filtered them by keywords such as "social anxiety", "SAD", and "fear" within the groups. To ensure timeliness, we screened to get a total of 4015 posts and comments with keywords between 2022-2023. 232 valid posts and comments were obtained.

To further identify user needs, we recruited 10 participants (3 males, 7 females, aged 23-29) from offline and conducted semi-structured interviews with them. After obtaining consent from the participants, we assessed the participants using the Liebowitz Social Anxiety Scale (LSAS) [42], and all of them demonstrated mild (>30) and higher social anxiety scores (M=63.2, scored 35-93). The interviews focused on their experiences of social anxiety, its

resolution, and their expectations for assisted relief. Each interview lasted an average of 45-60 minutes.

The qualitative data were analyzed using an open coding approach [35], and two co-authors worked together to identify the initial themes. After coding and analyzing the data, we first derived 8 different social anxiety themes, such as "talking to strangers", "public speaking", and "talking in a group". We used keywords in different posts to identify similar themes. For example, multiple posts mentioning "meeting strangers", "meeting people for the first time", "when I'm in a crowd of strangers", etc., which were categorized under the "talking to strangers" theme. Secondly, we analyzed the anxiety reduction methods that SAD patients considered effective by repeatedly studying the supportive comments in online forums and the data from semi-structured interviews, we summarised them as "cognitive restructuring", "exposure therapy", "social skills training" etc. Finally, in the semi-structured interviews, participants expressed their expectations for assistive relief methods, including gamification ("The game tasks are less stressful than talking to a counselor") and timely feedback ("I need some objective feedback that tells me how I'm doing", "I'd like to know how effective I am at practicing my expression"). At the same time, we asked participants to rate their game-playing level using a Likert scale of 1-5 (1=low-high=5), and most of them expressed confidence (M=3.8, SD=0.75). These results helped us to define our game design.

3.2 Therapy Approaches

CBT uses cognitive techniques, as well as behavioral techniques, such as exposure therapy [53], to help people with SAD identify and challenge their anxious thoughts and change problematic behaviors. Research has shown that combining cognitive restructuring and exposure therapy is effective for SAD [7].

In our interviews, we asked participants to assess their knowledge of CBT (1=Low-High=5), most of them did not know about CBT (M=2, SD=0.89), and a small number of participants (N=2) would use methods similar to "thought records" [3, 56] to help them cognitively reorganize themselves ("I would reflect by keeping a journal to check my negative emotions"), but did not identify the concept of cognitive distortions [9]. Meanwhile, most users (N=8) emphasized the importance of exposure therapy ("To overcome social anxiety in addition to psychological counseling, it is more important to deliberately practice speaking, exercise social skills"). These findings are consistent with the research, therefore, we embedded these two CBT techniques in our game:

1) Cognitive Restructuring: CBT believes that many negative emotions arise from irrational thoughts [4]. In Cognitive Restructuring, the therapist helps patients: 1) understand his or her emotional state; 2) identify negative thoughts associated with the emotional state [16]; and 3) challenge these negative thoughts [27]. In our design, players will express emotions, identify negative thoughts, and make suggestions to non-player characters (NPCs) through an interactive form of dialog with NPCs. Players' verbal emotions will affect the NPCs' emotional state, thus helping players understand that using more positive thoughts to replace negative thoughts will reduce negative emotions. At the same time, players will also be guided by knowledge of concepts such as cognitive distortion in recognizing negative thoughts.

¹Douban: <https://www.douban.com/>

2) Exposure Therapy: Exposure therapy is a behavioral technique in which patients face their fears under the guidance of a therapist [8]. Exposure therapy for social anxiety requires: 1) selecting specific exposure scenarios tailored to social interactions [55] and 2) effectively modeling the social scenarios [24, 60]. In our design, players will enter the specific social scenarios we set after completing the cognitive reconstruction task, engage in social tasks with NPCs in the form of a first-person perspective, and receive timely feedback, which will help players reflect and adjust their speech expression.

3.3 Game Overview

Our game, SoulSkipper, is a CBT-based role-playing game that combines voice control and emotion recognition. In this game, players will take on the role of "Soul Skipper" to help travelers who are lost in the "Sea of Emotions" to solve their emotional distress. Players will take a third-person perspective to understand the traveler's social anxiety situation, judge and identify the traveler's negative thoughts, and advise on voice input. Then, players will help the traveler reverse their emotional state through verbal emotion and will gain the "Rudder of Time" through the traveler's positive emotional changes, helping them return to the time before the emotional distress occurred. Finally, players will help the traveler perform social tasks from a first-person perspective to see the impact of positive behavior.

3.3.1 Game Genre. We chose "role-playing games" as our game genre. Role-playing is a commonly used tool in the treatment of SAD and can be used as an imaginative exposure to certain social situations [28]. At the same time, role-playing can also be better embedded with player-responsive narrative and emotional elements, increasing player empathy and engagement [1]. In terms of interaction, our game allows players to use voice input to talk to NPCs. This approach mimics the talk-back scenarios of actual therapy [61].

3.3.2 Game Play. Following the guidance of CBT, our game tasks will progress gradually according to the sequence of emotions, thoughts, and behaviors (Fig.1). Moreover, we embedded a fantastical narrative in the game to keep players continuously immersed in the gameplay [47]:

1) Accept Mission: Players navigate the "Sea of Emotions" and are guided by the "Nautical Map" to the location where the traveler is lost and accepts a new mission.

2) Enter Situation: Players will learn about the traveler's social difficulties and emotional distress through the traveler's narration and the display of images. Players will be guided by the "Navigation Light" to make judgments about irrational thoughts. If players' choices reflect any cognitive distortions, the "Navigation Light" will show players the concept of cognitive distortions and reasonable alternative thoughts.

3) Give Advice: Players will give advice to the traveler after identifying the traveler's negative thoughts. Players' verbal emotions will be detected. When players express positive emotions that shift the traveler's emotional state to positive, the traveler's emotions will provide the energy to help them return to the point before the emotional distress occurred.

4) Take action: Players use the "Rudder of Time" to go back in time, enter the traveler's perspective, and complete social tasks. Players will receive objective feedback on the quality of their speech expression.




Figure 1: The main interface of the game

3.4 Design Process

Drawing on our research into game players, therapy approaches, and gameplay, we further integrated three aspects: storytelling, interaction design, and visual design.

3.4.1 Storylines. 1) Scenario Development: The locus of SAD is often personalized [55]. Therefore, in our design, we focused on developing personalized social scenarios. Our story scenarios were all derived from the user research we completed in Phase 1. We ultimately identified 8 social scenarios including "talking to strangers" [36], "public speaking" [8], "talking in a group" [36], "talking in a job interview", "talking to people in authority" [32], "ordering a meal", "shopping at the mall", and "speaking at a conference". These scenarios were based on the themes we identified in the first phase and were further reviewed to generate stories.

2) Story Generation: After identifying the social scenario in which the stories took place, we further coded the themes in each scenario. For example, the scenario "talking to strangers" contains several sub-themes such as "I was afraid to speak", "I was afraid to look at each other", "My facial expressions and movements started to be unnatural", etc. These rich narrative details provide materials for our story generation. Based on these materials, we adapted our stories in conjunction with the Freytag's Pyramid [33] (a theatre structure in a five-act framework) and made sure they did not point to any particular original story.

3) Narrative Structure: Each story we generate will be treated as an independent chapter, and each chapter follows a consistent structure, using the ABCDEF model of CBT [48] as a framework. Referring to this framework, we divided the episodes into the following six sections: social anxiety situations (Activating Event), negative thoughts (Beliefs), behavioral and emotional consequences of negative thoughts (Consequences), counterarguments to negative thoughts (Disputation), shifts in positive thinking patterns (Effective New Beliefs), and behaviors (Future Action). Fig.2 shows the storylines for the chapter "Talking to Strangers".




Figure 2: Mapping the storylines to the ABCDEF-model

3.4.2 Interaction Design. 1) Emotional Adaptation: The emotions that players experience during gameplay can stimulate their immersion and engagement [13, 65]. In our design, we chose to embed an element of emotional adaptation to create a personalized gaming experience for the player [12], as well as to help players understand the principles of CBT and how emotions affect our thoughts and behaviors.

In previous studies, various modalities have been used to detect the player's emotional state, including physiological signals [26], voice intonation [19], and facial expressions [57]. In our design, we encourage players to communicate with NPCs through voice input, we use acoustic and textual features to recognize players' emotional state, which will be detected as positive or negative.

We adapt to the player's emotional state in two ways (Fig.3): 1) Adaptation of NPCs' emotional appearance: NPCs will display corresponding positive or negative emotional states according to the player's emotion; 2) Adaptive automation of task content: the game will provide automated assistance and prompts according to the player's emotion to encourage the player to shift to a positive mindset. The automatic feedback is selected from a predetermined list. When the player's emotion is detected as negative, the game guide will provide positive suggestions. Through this design, we enhance the connection between the player and the game character, helping the player visualize how they should support and encourage their shy friend, thus helping them recognize their irrational mindset. At the same time, the adaptive nature of the content encourages players to advance the game's tasks in a more dynamic way [43].

2) Objective Feedback: SAD patients often exhibit behaviors such as poor speech and avoidance of eye contact [4, 10], these




Figure 3: Emotional Adaptation and Objective Feedback

nonverbal behaviors can interfere with the delivery of information, and exacerbate the patient's negative emotions. For example, speaking too fast can limit the listener's understanding, and too low is not conducive to clarifying a message [29]. Exposure therapy helps patients gradually overcome fears and improve their behavior through planned exercises. In our design, we help players understand and adjust their vocal performance by extracting their volume and speech rate values and providing real-time personalized feedback.

In the game story, players should help the protagonist (the traveler) go back to the past, complete social tasks, and face the game's conversational objects from a first-person perspective (we set up one or more NPCs according to different social scenarios). We designed specific dialog quests for each chapter of the game and provided players with prompted texts that players could read or speak in his or her own way. When players speak, the game interface displays real-time objective feedback metrics, including volume and speech rate. When a speech task is completed, players will receive nonverbal feedback (a warm smile) from the NPCs, which encourages players to complete the task.

For the visualization of objective feedback, we considered two forms of display (Fig.3): 1) real-time quantitative display: dynamic volume bar and speech rate value; 2) qualitative indicator prompts: we provided alerts to players concerning the appropriate volume (50-60 dB) and speech rate value (150-180 words/min) [37] for the Chinese language (which is the native language of all our players). When players' voice performance was not within these thresholds, we provided them with textual reminders of "volume is a bit low/high" and "speech speed is a bit slow/fast". Considering the amount of information players can receive during a conversation [44], we set an interval of 20s [58] between these text alerts. During the design process, we will use an iterative design methodology to invite players to experience each of the above two forms of objective feedback and improve our design through players' evaluation results.

3.4.3 Visual Design. We refer to previous research and use positive visualization strategies to help SAD players reduce the negative experiences caused by stress. We embedded natural elements such

as plants and water in the game scene, research has shown that these visual elements have a positive impact on the player's emotional state [22]. For the overall visual background, we also chose a visual scheme with bright colors and high saturation to reduce the player's associations related to negative emotions [34].

4 TECHNICAL SUPPORT

In this game, we encourage players to interact with the game through voice input. To achieve this goal, we record their speaking voice during gameplay through a microphone and extract audio and text features to recognize their emotional changes. At the same time, we provide players with objective feedback on speech quality by measuring the speed and volume of the input audio.

4.1 Speech Emotion Recognition

4.1.1 Methodology. Initially, we extract text content through the pre-trained model Whisper [51]. Then, multiple pre-trained embeddings are employed to abstract features for multi-model SER-Fuse [49], including BERT [23] (Bidirectional Encoder Representations from Transformers) for text embedding feature vector generation and x-vector [54], ECAPA [52], d-vector [63] for audio embedding feature vector generation. After obtaining multiple feature vectors, a linear combination of all features into a single feature vector is achieved through concatenation. Subsequently, this fused feature vector is fed into an FC (Fully Connected) layer to generate the final prediction, providing results for negative, neutral, and positive sentiment in three-class emotion prediction (Fig.4).




Figure 4: The workflow of our speech emotion recognition

4.1.2 Implementation. The model is trained for 200 epochs on the CH-SIMS v2.0 [38] dataset, after which we input the collected 226 real player data into the model to obtain prediction results. It can be observed that due to the generally calm tone of users in everyday speech, the model sometimes exhibits errors in distinguishing between negative and neutral emotions. However, the total accuracy is still 71.68%, demonstrating a reasonable level of emotion recognition.

4.2 Real-time Feedback on Speed and Volume.

The calculation of the speech speed is implemented using the Tencent Automatic Speech Recognition [30, 66]. Based on real data testing, we found that when the duration of the input audio reaches three seconds, stable and accurate speed results can be obtained. To provide more immediate feedback, we record the speech and calculate the speed at three-second intervals. Then, we report it

to the player as a numerical value (words/minute). The player's volume is obtained through Audioworklet [20], which gets the microphone volume in real-time and maps the changes in the volume value through a bar graph.

5 PLAYER EVALUATION

We invited 10 participants who were recruited in user interviews to participate in the initial evaluation, each player was asked to complete 3 different game scenarios. In conjunction with our game goals, we initially evaluated our game prototype through the following three components: 1) assess whether our game was engaging for players through the System Usability Scale (SUS) [15], as well as the Engagement Scale adapted from the User Engagement Scale (UES) [46] and the Narrative Engagement Scale [17]; 2) assess whether our game was effective for player's CBT skills learning through the CBT Knowledge Test before and after the game; 3) assess whether the interactive features we designed were appropriate by inviting users to rate them (1=Low-High=5). After completing the questionnaire, we invited players to participate in a semi-structured interview to discuss their perceptions of our game.

Overall, our system had an average usability score of 80.75% and an average user engagement score of 3.73 (SD=0.43), which suggests that our system is attractive to players. Some players (N=3) believe that "the game gives some close-to-life scenarios, and through the experience of the game I can have better control of my emotions in actual social scenarios". The mean scores on our pre-game and post-game CBT knowledge tests (10 questions) were 4.0 (SD=1.53) and 7.17 (SD=1.5) respectively, which suggests that players improved their understanding of CBT skills through our game.

We designed two voice interactions: emotional adaptation and objective feedback. Initial evaluation results showed that players favored the emotional adaptation (M=4.08, SD=0.76), believing that "after I said what I thought, I felt that the character of the story got my help and it was very fulfilling". For both visual forms of objective feedback, users preferred real-time quantitative display (M=3.25, SD=1.64), and some players (N=5) felt that "the visual bars gave me a sense of participating in a positive challenge", but others (N=2) felt that this approach would "stress them out". Therefore, continued discussion of the balance between volume, speed, and participant comfort is necessary. Players also suggested that they needed a reminder of the qualitative indicators (M=2.83, SD=1.07). However, the current textual reminder was "not very attention-grabbing"; therefore, we need to consider more engaging and gentle approaches in our future work.

6 FUTURE WORK

Our future work will focus on two main areas:

1) Randomized Controlled Trial: First, we plan to conduct an iterative design to optimize the user experience and consider cultural adaptation. Second, due to the limitations and potential biases associated with the small sample size of the current experiment, we plan to invite more users to conduct a randomized controlled trial, set up a control group, and randomize participants. In addition, CBT requires consistency, and we plan to study the long-term effects of using this game; therefore, we will ask users to complete the full range of game scenarios we have developed and conduct repeated

measures in future experiments. These improvements aim to make the game a more effective digital mental health tool for SAD treatment, offering a creative solution to current treatment challenges and potentially broadening its applicability and effectiveness in the mental health domain.

2) Improvement of Technical Methods: To further improve the accuracy of speech emotion recognition, we are considering additional pre-trained speech emotion models as auxiliary resources. By combining the results of these models with the SER-Fuse model in a weighted manner, our goal is to obtain more accurate comprehensive outcomes, focusing on distinguishing between negative and neutral emotions. Additionally, we recognize that the dataset's quality significantly influences the model's training quality. Therefore, we plan to augment the dataset with high-quality data to enhance the accuracy of emotion classification.

ACKNOWLEDGMENTS

We thank all participants for their participation.

REFERENCES

- [1] Matthew S Abbott, Kimberly A Stauss, and Allen F Burnett. 2022. Table-top role-playing games as a therapeutic intervention with adults to increase social connectedness. *Social Work with Groups* 45, 1 (2022), 16–31.
- [2] Kristin N Anderson, Andrew B Jeon, Jordan A Blenner, Richard L Wiener, and Debra A Hope. 2015. How people evaluate others with social anxiety disorder: A comparison to depression and general mental illness stigma. *American Journal of Orthopsychiatry* 85, 2 (2015), 131.
- [3] Martin M Antony and Richard P Swinson. 2000. *The shyness & social anxiety workbook: Proven techniques for overcoming your fears*. New Harbinger Publications.
- [4] Joe Ayres. 1988. Coping with speech anxiety: The power of positive thinking. *Communication Education* 37, 4 (1988), 289–296.
- [5] Luke Balcombe and Diego De Leo. 2022. Human-computer interaction in digital mental health. In *Informatics*, Vol. 9. MDPI, 14.
- [6] Imre Dániel Báldy, Nikolaj Hansen, and Thomas Bjørner. 2020. How to design and evaluate a serious game aiming at awareness of therapy skills associated with social anxiety disorder. In *Proceedings of the 6th EAI International Conference on Smart Objects and Technologies for Social Good*. 156–162.
- [7] David H Barlow. 2021. *Clinical handbook of psychological disorders: A step-by-step treatment manual*. Guilford publications.
- [8] Terri L Barrera, Derek D Szafranski, Chelsea G Ratcliff, Sarah L Garnaat, and Peter J Norton. 2016. An experimental comparison of techniques: Cognitive defusion, cognitive restructuring, and in-vivo exposure for social anxiety. *Behavioural and cognitive psychotherapy* 44, 2 (2016), 249–254.
- [9] Aaron T Beck. 1963. Thinking and depression: I. Idiosyncratic content and cognitive distortions. *Archives of general psychiatry* 9, 4 (1963), 324–333.
- [10] Graham D Bodie. 2010. A racing heart, rattling knees, and ruminative thoughts: Defining, explaining, and treating public speaking anxiety. *Communication education* 59, 1 (2010), 70–105.
- [11] Johanna Boettcher, Per Carlbring, Babette Renneberg, and Thomas Berger. 2013. Internet-based interventions for social anxiety disorder-an overview. *Verhaltens-therapie* 23, 3 (2013), 160–168.
- [12] Boyan Bontchev. 2016. Adaptation in affective video games: A literature review. *Cybernetics and Information Technologies* 16, 3 (2016), 3–34.
- [13] Boyan Bontchev and Dessislava Vassileva. 2016. Assessing engagement in an emotionally-adaptive applied game. In *Proceedings of the fourth international conference on technological ecosystems for enhancing multicultural*. 747–754.
- [14] Stephane Bouchard, Stephanie Dumoulin, Genevieve Robillard, Tanya Guitard, Evelyn Klingler, Helene Forget, Claudie Loranger, and Francois Xavier Roucaut. 2017. Virtual reality compared with in vivo exposure in the treatment of social anxiety disorder: a three-arm randomised controlled trial. *The British Journal of Psychiatry* 210, 4 (2017), 276–283.
- [15] John Brooke et al. 1996. SUS-A quick and dirty usability scale. *Usability evaluation in industry* 189, 194 (1996), 4–7.
- [16] David D Burns. 1989. *The feeling good handbook: Using the new mood therapy in everyday life*. William Morrow & Co.
- [17] Rick Busselle and Helena Bilandzic. 2009. Measuring narrative engagement. *Media psychology* 12, 4 (2009), 321–347.
- [18] Gillian Butler. 2016. *Overcoming social anxiety and shyness: A self-help guide using cognitive behavioural techniques*. Hachette UK.
- [19] Ricardo A Calix, Leili Javadpour, and Gerald M Knapp. 2012. Detection of affective states from text and speech for real-time human–computer interaction. *Human factors* 54, 4 (2012), 530–545.
- [20] Hongchan Choi. 2018. AudioWorklet: The future of web audio. In *ICMC*.
- [21] Pim Cuijpers, Tara Donker, Annemieke van Straten, J Li, and Gerhard Andersson. 2010. Is guided self-help as effective as face-to-face psychotherapy for depression and anxiety disorders? A systematic review and meta-analysis of comparative outcome studies. *Psychological medicine* 40, 12 (2010), 1943–1957.
- [22] Li Deng, Xi Li, Hao Luo, Er-Kang Fu, Jun Ma, Ling-Xia Sun, Zhuo Huang, Shi-Zhen Cai, and Yin Jia. 2020. Empirical study of landscape types, landscape elements and landscape components of the urban park promoting physiological and psychological restoration. *Urban Forestry & Urban Greening* 48 (2020), 126488.
- [23] Jacob Devlin, Ming-Wei Chang, Kenton Lee, and Kristina Toutanova. 2018. Bert: Pre-training of deep bidirectional transformers for language understanding. *arXiv preprint arXiv:1810.04805* (2018).
- [24] Meriem El-Yamri, Alejandro Romero-Hernandez, Manuel Gonzalez-Riojo, and Borja Manero. 2019. Emotions-responsive audiences for VR public speaking simulators based on the speakers' voice. In *2019 IEEE 19th international conference on advanced learning technologies (ICALT)*, Vol. 2161. IEEE, 349–353.
- [25] Sandra Garrido, Chris Millington, and Emery Schubert. 2019. What works and what doesn't work? A systematic review of digital mental health interventions for depression and anxiety in young people. *Frontiers in psychiatry* 10 (2019), 441740.
- [26] Kunal Gupta, Yuewei Zhang, Yun Suen Pai, and Mark Billingham. 2021. Wizard-ofvr: An emotion-adaptive virtual wizard experience. In *SIGGRAPH Asia 2021 XR*. 1–2.
- [27] Richard G Heimberg. 2002. Cognitive-behavioral therapy for social anxiety disorder: current status and future directions. *Biological psychiatry* 51, 1 (2002), 101–108.
- [28] Richard G Heimberg, Debra G Salzman, Craig S Holt, and Karl A Blendell. 1993. Cognitive-behavioral group treatment for social phobia: Effectiveness at five-year followup. *Cognitive therapy and Research* 17, 4 (1993), 325–339.
- [29] Rebecca Hincks. 2005. Measures and perceptions of liveliness in student oral presentation speech: A proposal for an automatic feedback mechanism. *System* 33, 4 (2005), 575–591.
- [30] Geoffrey Hinton, Li Deng, Dong Yu, George E Dahl, Abdel-rahman Mohamed, Navdeep Jaitly, Andrew Senior, Vincent Vanhoucke, Patrick Nguyen, Tara N Sainath, et al. 2012. Deep neural networks for acoustic modeling in speech recognition: The shared views of four research groups. *IEEE Signal processing magazine* 29, 6 (2012), 82–97.
- [31] Chris Hollis, Caroline J Falconer, Jennifer L Martin, Craig Whittington, Sarah Stockton, Cris Glazebrook, and E Bethan Davies. 2017. Annual Research Review: Digital health interventions for children and young people with mental health problems—a systematic and meta-review. *Journal of Child Psychology and Psychiatry* 58, 4 (2017), 474–503.
- [32] Jon-Chao Hong, Ming-Yueh Hwang, Chin-Hao Hsu, Kai-Hsin Tai, and Yen-Chun Kuo. 2015. Belief in dangerous virtual communities as a predictor of continuance intention mediated by general and online social anxiety: The Facebook perspective. *Computers in Human Behavior* 48 (2015), 663–670.
- [33] Carol Jago. 2004. Literature: Elements of Style: Stop Pretending and Think about Plot. *Voices from the Middle* 11, Teaching English Language Learners (2004), 50–51.
- [34] Naz Kaya and Helen H Epps. 2004. Relationship between color and emotion: A study of college students. *College student journal* 38, 3 (2004), 396–405.
- [35] Shahedul Huq Khandkar. 2009. Open coding. *University of Calgary* 23, 2009 (2009).
- [36] Eleanor Leigh and David M Clark. 2018. Understanding social anxiety disorder in adolescents and improving treatment outcomes: Applying the cognitive model of Clark and Wells (1995). *Clinical child and family psychology review* 21, 3 (2018), 388–414.
- [37] Changshuan Li. 2010. Coping strategies for fast delivery in simultaneous interpretation. *The Journal of Specialised Translation* 13 (2010), 19–25.
- [38] Yihe Liu, Ziqi Yuan, Huisheng Mao, Zhiyun Liang, Wanqiu Yue Yang, Yuanzhe Qiu, Tie Cheng, Xiaoteng Li, Hua Xu, and Kai Gao. 2022. Make acoustic and visual cues matter: CH-SIMS v2. 0 dataset and AV-Mixup consistent module. In *Proceedings of the 2022 International Conference on Multimodal Interaction*. 247–258.
- [39] Stéphanie Mader, Guillaume Levieux, and Stéphanie Natkin. 2016. A game design method for therapeutic games. In *2016 8th International Conference on Games and Virtual Worlds for Serious Applications (VS-GAMES)*. IEEE, 1–8.
- [40] Evan Mayo-Wilson, Sofia Dias, Ifigeneia Mavranzouli, Kayleigh Kew, David M Clark, AE Ades, and Stephen Pilling. 2014. Psychological and pharmacological interventions for social anxiety disorder in adults: a systematic review and network meta-analysis. *The Lancet Psychiatry* 1, 5 (2014), 368–376.
- [41] Hugh Cameron McCall, Chris G Richardson, Fjola Dogg Helgadóttir, and Frances S Chen. 2018. Evaluating a web-based social anxiety intervention among university students: randomized controlled trial. *Journal of Medical Internet Research* 20, 3 (2018), e91.

- [42] Douglas S Mennin, David M Fresco, Richard G Heimberg, Franklin R Schneier, Sharon O Davies, and Michael R Liebowitz. 2002. Screening for social anxiety disorder in the clinical setting: using the Liebowitz Social Anxiety Scale. *Journal of anxiety disorders* 16, 6 (2002), 661–673.
- [43] Domen Novak, Matjaž Mihelj, and Marko Munič. 2012. A survey of methods for data fusion and system adaptation using autonomic nervous system responses in physiological computing. *Interacting with computers* 24, 3 (2012), 154–172.
- [44] Eyal Ofek, Shamsi T Iqbal, and Karin Strauss. 2013. Reducing disruption from subtle information delivery during a conversation: mode and bandwidth investigation. In *Proceedings of the SIGCHI Conference on Human Factors in Computing Systems*. 3111–3120.
- [45] James C Overholser. 2002. Cognitive-behavioral treatment of social phobia. *Journal of Contemporary Psychotherapy* 32 (2002), 125–144.
- [46] Heather L O'Brien, Paul Cairns, and Mark Hall. 2018. A practical approach to measuring user engagement with the refined user engagement scale (UES) and new UES short form. *International Journal of Human-Computer Studies* 112 (2018), 28–39.
- [47] Louise E Parker and Mark R Lepper. 1992. Effects of fantasy contexts on children's learning and motivation: Making learning more fun. *Journal of personality and social psychology* 62, 4 (1992), 625.
- [48] Jonathan Passmore, Tracy Sinclair, Jonathan Passmore, and Tracy Sinclair. 2020. Cognitive Behavioral Approach and ABCDEF Model. *Becoming a Coach: The Essential ICF Guide* (2020), 127–132.
- [49] Nhat Truong Pham, Le Thi Phan, Duc Ngoc Minh Dang, and Balachandran Manavalan. 2023. SER-Fuse: An Emotion Recognition Application Utilizing Multi-Modal, Multi-Lingual, and Multi-Feature Fusion. In *Proceedings of the 12th International Symposium on Information and Communication Technology*. 870–877.
- [50] KATHRYN Ponniah and SD Hollon. 2008. Empirically supported psychological interventions for social phobia in adults: a qualitative review of randomized controlled trials. *Psychological medicine* 38, 1 (2008), 3–14.
- [51] Alec Radford, Jong Wook Kim, Tao Xu, Greg Brockman, Christine McLeavey, and Ilya Sutskever. 2023. Robust speech recognition via large-scale weak supervision. In *International Conference on Machine Learning*. PMLR, 28492–28518.
- [52] Mirco Ravanelli, Titouan Parcollet, Peter Plantinga, Aku Rouhe, Samuele Cornell, Loren Lugosch, Cem Subakan, Nauman Dawalatabad, Abdelwahab Heba, Jianyuan Zhong, et al. 2021. SpeechBrain: A general-purpose speech toolkit. *arXiv preprint arXiv:2106.04624* (2021).
- [53] Thomas L Rodebaugh, Robert M Holaway, and Richard G Heimberg. 2004. The treatment of social anxiety disorder. *Clinical Psychology Review* 24, 7 (2004), 883–908.
- [54] David Snyder, Daniel Garcia-Romero, Gregory Sell, Daniel Povey, and Sanjeev Khudanpur. 2018. X-vectors: Robust dnn embeddings for speaker recognition. In *2018 IEEE international conference on acoustics, speech and signal processing (ICASSP)*. IEEE, 5329–5333.
- [55] Susan H Spence and Ronald M Rapee. 2016. The etiology of social anxiety disorder: An evidence-based model. *Behaviour research and therapy* 86 (2016), 50–67.
- [56] Andrew A Sweet and Andrew L Loizeaux. 1991. Behavioral and cognitive treatment methods: A critical comparative review. *Journal of Behavior Therapy and Experimental Psychiatry* 22, 3 (1991), 159–185.
- [57] Chek Tien Tan, Sander Bakkes, and Yusuf Pisan. 2014. Inferring player experiences using facial expressions analysis. In *Proceedings of the 2014 Conference on Interactive Entertainment*. 1–8.
- [58] M Iftikhar Tanveer, Emy Lin, and Mohammed Hoque. 2015. Rhema: A real-time in-situ intelligent interface to help people with public speaking. In *Proceedings of the 20th international conference on intelligent user interfaces*. 286–295.
- [59] Hailey Tremain, Carla McEnery, Kathryn Fletcher, Greg Murray, et al. 2020. The therapeutic alliance in digital mental health interventions for serious mental illnesses: narrative review. *JMIR mental health* 7, 8 (2020), e17204.
- [60] Bao Truong, Trung-Nghia Le, Khanh-Duy Le, Minh-Triet Tran, and Tam V Nguyen. 2022. Public Speaking Simulator with Speech and Audience Feedback. In *2022 IEEE International Symposium on Mixed and Augmented Reality Adjunct (ISMAR-Adjunct)*. IEEE, 855–858.
- [61] Samuel M Turner, Deborah C Beidel, and Michele R Cooley-Quille. 1995. Two-year follow-up of social phobics treated with Social Effectiveness Therapy. *Behaviour Research and Therapy* 33, 5 (1995), 553–555.
- [62] Vihang N Vahia. 2013. Diagnostic and statistical manual of mental disorders 5: A quick glance. *Indian journal of psychiatry* 55, 3 (2013), 220–223.
- [63] Ehsan Variani, Xin Lei, Erik McDermott, Ignacio Lopez Moreno, and Javier Gonzalez-Dominguez. 2014. Deep neural networks for small footprint text-dependent speaker verification. In *2014 IEEE international conference on acoustics, speech and signal processing (ICASSP)*. IEEE, 4052–4056.
- [64] Philip S Wang, Patricia Berglund, Mark Olfson, Harold A Pincus, Kenneth B Wells, and Ronald C Kessler. 2005. Failure and delay in initial treatment contact after first onset of mental disorders in the National Comorbidity Survey Replication. *Archives of general psychiatry* 62, 6 (2005), 603–613.
- [65] Georgios N Yannakakis and John Hallam. 2008. Entertainment modeling through physiology in physical play. *International Journal of Human-Computer Studies* 66, 10 (2008), 741–755.
- [66] Yongqiang Zou, Xing Jin, Yi Li, Zhimao Guo, Eryu Wang, and Bin Xiao. 2014. Mariana: Tencent deep learning platform and its applications. *Proceedings of the VLDB Endowment* 7, 13 (2014), 1772–1777.