Supplementary Material

PRISM: an auditable, phase-aware pipeline for post-treatment glioma surveillance — rigorous evaluation of imaging, clinical and treatment-course information

Draft, 2026-10-15. Every number below is copied from the result documents in `docs/results/` *and* `docs/manuscript/` *or from* `outputs/results/`*; the file is named in each section. "Grade 4" (G4) is the primary population; "all" = all gliomas. AUC = IPCW AUC(120 d), mean over 5 repeats of the per-repeat pooled out-of-fold value, with 95% patient-cluster bootstrap CI (1,000 draws), unless stated otherwise. PRISM v1 is the first version of PRISM, described in an unpublished internal report; "PRISM's initial radiomics model" and "our initial evaluation split" are its radiomics model and its seed-62 test split. "Baseline" = PRISM's core model (clinical + timing + as-of-scan treatment, ridge), the deployed configuration and the reference for every imaging comparison.*

Contents

S1 Pre-specification register **3**

S2 Cohort flow **4**

S3 Outcome, censoring rules and label formulations **5**

S4 Predictor definitions and the as-of-scan treatment rule **6**

S5 Radiomics: extraction, configurations and outlier clipping **8**

S6 Frozen deep features **11**

S7 PRISM v1: the initial radiomics model, the initial evaluation split and the fusion components **12**

S8 Split variance: full tables and verifier checks **13**

S9 Estimator and evaluation details **15**

S10 Feature-group search with permutation control **15**

S11 Phase-of-care ladder: full results **16**

S12 Secondary analyses: baselines, sensitivity, imbalance, schedule, feature conflicts, SHAP, calibration and utility **18**

S12.1 Literature baselines and tuning 18

S12.2 Sensitivity of the primary comparison 18

S12.3 Class imbalance 19

S12.4 Schedule coupling 19

S12.5 Feature conflicts 20

S12.6 SHAP attributions 20

S12.7 Calibration, decision curves, added value and sample size 20
S13 External evaluation on LUMIERE: full results **21**
S14 Imaging-defined label, imaging diagnostics and re-segmentation **23**
S15 The auditable layer: full results and an example audit record **25**
S16 Reproducibility **29**
S17 Reporting checklists (TRIPOD+AI, CLEAR) cross-referenced to the manuscript **29**
S18 CFB-GBM diffusion analysis **31**
S19 PRISM report generator: design and faithfulness evaluation **32**
Supplementary figures **32**

S1 Pre-specification register

Analysis	Status	Plan record	Decision rule / primary quantity	Outcome
Radiomics vs baseline, nested over 70 configurations, grade 4	Pre-specified primary (development)	Publication plan §3.3	ΔAUC with 95% CI	−0.032 [−0.084, +0.017]; no gain
LUMIERE locked clinical + timing + volumes, grade-4-trained	Pre-specified external model	`EXTERNAL_LUMIERE.md`	Pooled IPCW AUC	0.608 / 0.620 (ridge / boosting)
Phase-of-care ladder, post-RT R2 + I2 vs R2	Pre-specified primary; plan in module docstring, not separately committed; a smoke run printed some I0–I2 numbers before the full run; plan unchanged	`PHASE_LADDER.md` §1	ΔAUC with CI and 100-permutation p	+0.061 [−0.038, +0.156], p = 0.21; not supported
Imaging-defined (clean) label	Pre-specified; plan snapshot saved before the run	`outputs/results/analyses/clean_label/PLAN_snapshot.md`	ΔAUC ≥ 0.03 with CI excluding 0	+0.027 [−0.033, +0.082]; not met
Re-segmentation	Rule written before any result was seen	`RESEGMENTATION.md`	Detection Δ ≥ 0.05 with CI > 0, or ET larger at progression in > 65% of pairs	−0.016 [−0.057, +0.025]; 45%; not met
CFB-GBM ADC	Pre-specified; plan snapshot saved before the run	`outputs/results/analyses/cfb_adc/PLAN_snapshot.md`	Δ(d − c) ≥ 0.03 with CI excluding 0	+0.007 [−0.019, +0.035], p = 0.151; not met
Audit layer P1 (coverage) and P2 (abstention beyond case mix)	Pre-specified; plan snapshot saved before the run	`outputs/results/analyses/audit_system/PLAN_snapshot.md`	P1 coverage ≥ 0.78; P2 CI above 0	P1 0.841 [0.790, 0.885], met; P2 +0.006 [−0.023, +0.036], not met
Split variance (500 splits)	Descriptive; designed after the single-split result was known	`SPLIT_VARIANCE.md`	—	—
Positive controls, SHAP, feature conflicts, imbalance, schedule coupling, baselines, sensitivity, utility	Secondary or exploratory	respective documents	—	—

Sequence: the clean-label, re-segmentation, CFB-GBM and audit analyses were designed after the main imaging result (no added value) was known. Each had its own rule fixed before its own run. Contrasts beyond the primary ones are not corrected for multiplicity.

S2 Cohort flow

Source: `docs/manuscript/FLOW.md` (`outputs/data/mu_flow.csv`, `outputs/results/analyses/tables/lumiere_flow.csv`). Figure: `docs/manuscript/cohort_flow.png` (Figure S1).

Supplementary Table S1a. MU-Glioma-Post.

Step	Scans	Patients	Excluded
All timepoints	596	203	–
All four sequences and mask present	594	203	2
Tumour (labels 1–3) present	592	203	2
MRI day known	589	202	3
MRI day plausible (≥ -365 d, not before an earlier timepoint)	588	202	1
Progression date known for progressors	588	202	0
MRI not after recorded death	580	202	8
Before late treatment (2nd-line, immunotherapy, new treatment)	501	199	79
Before first progression (index scans)	**308**	**155**	193
Grade 4 subset	249	126	–

Supplementary Table S1b. LUMIERE.

Step	Studies	Patients	Excluded
All rated studies	616	91	–
Post-operative (week ≥ 2 after surgery)	456	81	160
Rated SD / PR / CR / Post-Op	199	70	257
Before first progression	132	61	67
Not after recorded death (index scans)	**132**	**61**	0

Censored scans (52 MU, 39 grade 4, 9 LUMIERE at 120 d) stay in the cohort and are handled by the censoring-aware labels and IPCW.

Table S2. The full cohort table with definitions is `docs/manuscript/TABLE1_COHORT.md` (`table1.csv`). Additional rows not in main Table 1: last documented follow-up after scan, median [IQR] days 275 [107–471] (all), 285 [112–478] (G4), 462 [238–721] (LUMIERE); time to progression or censoring 125 [50–237], 111 [48–208], 168 [91–287]; progression documented after the scan at any time 220 (71%), 190 (76%), 122 (92%); LUMIERE T1c field strength 1.5 T / 3 T / other or missing 81 (61%) / 44 (33%) / 7; MU field strength not distributed. WHO grade in all gliomas: grade 1, 3 (2%); grade 2, 23 (15%); grade 3, 2 (1%); "3 vs 4", 1 (1%); grade 4, 126 (81%).

S3 Outcome, censoring rules and label formulations

Source: `docs/results/FIX_LABELS.md`, `docs/results/C1_endpoint.md`, `docs/results/C1_full_tables.md`.

Estimand. Documented first progression within H = 120 d of the index scan (90 / 180 d as sensitivities). Death without documented progression censors at the death day. Label 0 requires progression after H or progression-free follow-up $\geq$ H. Composite (progression or death) is a sensitivity analysis only.

Supplementary Table S3. Rules adopted in the label audit.

#	Problem found	Rule
1	Death without documented progression treated as a verified negative at every horizon	Death censors at the death day (primary); composite event as sensitivity
1b	MRIs dated after the recorded death (15/96 MU deaths)	If more than one dated record follows the death day, the death date is ignored (7 patients, 3 with cohort scans); if exactly one, that scan is excluded
2	Progression flag 0 with a progression date (3 patients); two date fields disagree (29 patients)	Any patient with a progression date is a progressor; primary date `date_of_first_progression`; reverse order as sensitivity
3	One MRI day of −965; progressor who died before progression	MRI day ≥ -365 and not before an earlier timepoint; death-before-progression flagged
4	Hazard formulation counted scans censored mid-interval as surviving the interval	Censored scans contribute only fully observed intervals
5	Follow-up extended by non-imaging dates	Kept as primary; imaging-only follow-up as sensitivity
6	Unknown RT end coded as "not early post-RT"	`early_post_rt` is 1 / 0 / missing (94 scans missing, 65 grade 4)

Effect: six scans left the cohort (314 → 308; grade 4 253 → 249). At 120 d, 17 negatives became censored (16 deaths within 120 d without progression, 1 ignored death date) and 1 negative became positive; no positive changed.

Supplementary Table S4. Label counts (positive / verified negative / censored).

Population	Label	H = 90	H = 120	H = 180
All (308 / 155)	primary	65 / 195 / 48	100 / 156 / 52	144 / 105 / 59
All	composite	79 / 195 / 34	116 / 156 / 36	161 / 105 / 42
All	alternative progression date	63 / 191 / 54	99 / 151 / 58	141 / 102 / 65
All	imaging-only follow-up	65 / 187 / 56	100 / 149 / 59	144 / 99 / 65
Grade 4 (249 / 126)	primary	62 / 152 / 35	93 / 117 / 39	131 / 75 / 43
Grade 4	composite	74 / 152 / 23	107 / 117 / 25	146 / 75 / 28
LUMIERE (132 / 61)	primary	24 / 101 / 7	47 / 76 / 9	62 / 61 / 9
LUMIERE	composite	26 / 101 / 5	51 / 76 / 5	66 / 61 / 5

LUMIERE gives a survival time without a death indicator; a recorded survival time is read as a death and used only as a censoring time in the primary label.

IPCW AUC. Cumulative/dynamic AUC at 120 d with Kaplan–Meier censoring weights (Uno et al., 2007; Blanche et al., 2013). Cases are scans with an event by 120 d, weighted by the inverse of the censoring survival at their event time; controls are scans event-free beyond 120 d. The vectorised implementation matches scikit-survival to 2×10^{-16} (`FIX_EVALUATION.md` §1). In the bootstrap, censoring weights are re-estimated in each resample; if a resample has no censoring, G = 1.

Supplementary Table S5. Label formulations, grade 4, H = 120 d (`C1_endpoint.md`; ΔAUC vs naive).

Features	Formulation	AUC [95% CI]	ΔAUC vs naive [95% CI]	O/E	Slope
volumes + clinical	naive	0.651 [0.567, 0.727]	—	1.14	0.95
	verified	0.657 [0.577, 0.732]	+0.006 [−0.020, +0.030]	0.95	0.91
	IPCW	0.656 [0.578, 0.731]	+0.005 [−0.021, +0.029]	0.97	0.92
	hazard	0.653 [0.574, 0.726]	+0.002 [−0.018, +0.022]	0.87	0.50
legacy radiomics + clinical	naive	0.648 [0.553, 0.736]	—	1.13	0.52
	verified	0.655 [0.575, 0.730]	+0.007 [−0.021, +0.038]	0.96	0.51
	IPCW	0.654 [0.575, 0.730]	+0.006 [−0.022, +0.036]	0.98	0.51
	hazard	0.669 [0.583, 0.746]	+0.021 [−0.027, +0.068]	0.90	0.33

All 36 ΔAUC CIs (2 populations × 2 feature sets × 3 horizons × 3 formulations) include 0. The naive label under-predicts risk in all 12 settings (O/E 1.12–1.19 against 0.95–0.99 for verified / IPCW). The legacy radiomics pipeline in C1 predates the clipping fix (S5).

S4 Predictor definitions and the as-of-scan treatment rule

Source: `prism/features/tabular.py`; `docs/results/FIX_TREATMENT_ASOF.md`; `docs/results/FIX_FEATURES.md`; audit record input tables (`outputs/results/analyses/audit_system/records.md`).

Supplementary Table S6. Baseline predictors (23 raw; 31 design columns after missingness indicators).

Group	Predictor	Definition (as of scan day)	Source column(s)
Clinical	age	age at diagnosis, years	`clinical_age_at_diagnosis`
	male	sex at birth	`clinical_sex_at_birth`
	idh_mutant, idh_unknown	IDH1/IDH2 mutant; neither documented	`clinical_idh1_mutation`, `clinical_idh2_mutation`
	mgmt_methylated, mgmt_unknown	MGMT promoter methylated; unknown code	`clinical_mgmt_methylation`
Timing	days_from_diagnosis	scan day	`days_from_diagnosis_to_mri`
	days_since_rt_end	days since documented RT end; missing until RT completed	RT end day vs scan day
	rt_date_unknown	no completed RT with known end date at the scan	RT end day
	early_post_rt	0–84 d after RT end (RANO 12-week window); missing if RT end unknown	RT end day
	early_post_rt_x_mgmt_methylated	interaction	RT end day × MGMT
	visit_index	rank of the scan among the patient's MRIs	MRI days
Treatment	rt_started	RT start day ≤ scan day	`clinical_radiation_therapy` + RT start day
	rt_completed	RT end day ≤ scan day	RT end day
	rt_dose_gy, rt_fractions, rt_hypofractionated	prescription, visible once RT started; hypofractionation ≤ 15 fractions	`clinical_dose`, `clinical_number_of_fractions`
	rt_start_day	days from diagnosis to RT start, visible once started	RT start day
	rt_duration_days	RT end − start, visible once completed	RT start and end days
	chemo_started	initial TMZ / lomustine start ≤ scan day	initial chemotherapy fields
	surgery_day, biopsy_before_resection	visible once the first surgery has happened	first surgery fields
	previous_brain_tumor	history, known at any scan	`clinical_previous_brain_tumor`

Unknown values are imputed in-fold with the training median plus a missingness indicator; nothing is zero-filled silently. Therapy marked given without a start date (2 RT, 2 chemotherapy patients) is assumed to have started before the scan.

Supplementary Table S7. What leaked before the as-of-scan fix (scans affected, grade 4 / all gliomas).

Feature	Problem	Affected
`rt_given`, `chemo_given`	"will receive RT / chemotherapy", true for scans before the course started	13 / 23 (RT), 13 / 21 (chemo)
dose, fractions, hypofractionation, `rt_start_day`	prescription and start date of a course that had not started	13 / 19–23
`rt_duration_days`	duration of a course not yet finished (also reveals early stopping)	17 / 29
`days_since_rt_end`, `rt_date_unknown`	negative values from the future RT end date	17 / 29
`surgery_day`, `biopsy_before_resection`	one scan precedes the first surgery	1 / 1

Progression, death, last follow-up and late-treatment dates are used only for labels, censoring and eligibility. Change features use earlier scans only.

Effect on repeated CV (intermediate inputs; earlier repeat-averaged estimator; `FIX_TREATMENT_ASOF.md` §3). Baseline ridge grade 4 0.694 [0.613, 0.775] → 0.729 [0.648, 0.811]; all gliomas 0.719 [0.648, 0.788] → 0.749 [0.675, 0.820]. Boosting grade 4 0.723 [0.633, 0.806] → 0.731 [0.641, 0.813]; all gliomas 0.743 [0.666, 0.816] → 0.740 [0.663, 0.815]. Timing-only boosting fell (grade 4 0.718 → 0.696; all 0.735 → 0.705): part of its earlier signal came from the future RT end date. The MGMT × window flag changed repeated-CV AUC by ≤ 0.008. These intermediate values are superseded by the final run (main Table 2).

Molecular look-ahead found by the audit (exploratory). IDH and MGMT are used at scans taken before the diagnosis date: 8 grade 4 scans (7 patients; 1 event, 4 non-events, 3 censored) and 11 all-glioma scans, 1–58 days before diagnosis (`AUDIT_SYSTEM.md` §2). Its effect on the baseline AUC was not quantified.

Imaging measurements. Volumes (ET, NETC, SNFH, RC, tumour = ET + NETC, lesion), morphology (ET fraction, multifocality, largest axial ET area, ET-to-cavity distance and proximity, T1c / FLAIR intensities relative to brain > 10 mm (Euclidean) from lesion and cavity), and past-only change (from previous scan and nadir, growth rate, `vol_pd_flag` requiring a measurable nadir ≥ 0.1 cc, `et_new_from_zero_nadir`, missingness flags for undefined relative change; 25 change columns). For LUMIERE transport, necrosis and cavity were merged (`log_necrosis_cavity_cc`) because DeepBraTumIA labels the cavity as necrosis (`FIX_FEATURES.md` §1).

S5 Radiomics: extraction, configurations and outlier clipping

Source: `docs/results/C3_radiomics.md`, `docs/results/FIX_RADIOMICS_CLIP.md`, `docs/results/FIX_FEATURES.md` §6, `prism/features/radiomics.py`.

Extraction. PyRadiomics (van Griethuysen et al., 2017). *Legacy* features are those of PRISM's initial radiomics model: T1c and FLAIR, original, LoG and wavelet images, 2,446 columns. Under the legacy within-lesion z-score and fixed bin width the texture matrices are effectively two-level (GLCM joint entropy ≤ 2 bits), so legacy texture behaves as a binary habitat descriptor (above vs below lesion mean). *Corrected* (v2) features use a whole-brain z-score, a fixed bin count of 32 (joint entropy 4.7–6.3 bits), no second N4 correction, and are extracted per compartment (ET, NETC, SNFH, core, lesion, 5 mm peri-cavity rim, shape) and for all four sequences (2,316 columns). Known limitations: a fixed bin count rather than the IBSI-recommended fixed bin width (Zwanenburg et al., 2020); the ICC stability pass covers T1c and FLAIR texture and first-order features only (1,116 of 2,316 columns); the 5 mm rim uses a city-block dilation.

Configurations (70). Variant (legacy; corrected whole lesion; corrected per compartment; corrected all four sequences) $\times$ stability filter (none; ICC ≥ 0.75 under 1-voxel erosion, in-fold) $\times$ selection (top-10 or top-30 univariate F; L1 with non-zero coefficients only, at most 30; PCA-10; decorrelation + top-20) $\times$ integration (concatenation with the baseline; or a stacked radiomics score from inner out-of-fold predictions). Nested choice: inner 3-fold patient-grouped AUC in each outer training fold. Missing compartments are imputed with the median and one indicator per missingness pattern.

Outlier clipping. Three legacy columns (FLAIR LoG $\sigma = 1$, 2, 3 mm NGTDM Coarseness) held the PyRadiomics sentinel 10^6 in four scans (two grade 4); other values in these columns have medians of 10^{-4} to 8×10^{-4}. One such value inflates a training SD by orders of magnitude and makes PCA and the stacked score depend on these scans. Before the fix, the stacked score had inner-OOF training values of -10^4 to -4.6×10^6 in 20/25 grade 4 fits. The fix, `QuantileClip(q = 0.01)`, is fitted in each training (and inner) fold, clips to the fold's 1st/99th order statistics, and applies the same bounds to test scans. It does not look at labels and applies to all configurations and to deep features.

Supplementary Table S8. Effect of the clipping fix (`FIX_RADIOMICS_CLIP.md`).

Population	Nested AUC before → after	ΔAUC before → after	Cal. slope
Grade 4	0.672 [0.597, 0.743] → 0.666 [0.586, 0.739]	−0.026 [−0.073, +0.017] → −0.032 [−0.084, +0.017]	0.50 → 0.60
All gliomas	0.732 [0.664, 0.794] → 0.747 [0.678, 0.811]	+0.012 [−0.029, +0.050] → +0.028 [−0.017, +0.071]	0.77 → 0.85

Largest change in any single configuration: 0.020. Single configurations with CI above 0: grade 4 0/70 before and after (below 0: 12/70 → 8/70); all gliomas 0/70 → 1/70 (`legacy|none|pca10|stack|ridge`, 0.754, +0.035 [+0.002, +0.068], $p(\Delta \leq 0) = 0.021$). Not covered by the fix: the radiomics group of the feature-group search (S10), the C1 radiomics feature set (S3), and the feature-conflict B + radiomics row (S12).

Supplementary Table S9. Radiomics results by population (`C3_radiomics.md`; `outputs/results/c3/{grade4,all_glioma}/results.csv`).

Model	Grade 4 AUC	Δ [95% CI]	Slope	All gliomas AUC	Δ [95% CI]	Slope
Baseline ridge	0.698 [0.615, 0.774]	—	0.89	0.720 [0.638, 0.792]	—	0.88
Baseline boosting	0.687 [0.601, 0.768]	−0.012 [−0.072, +0.048]	0.82	0.718 [0.642, 0.792]	−0.002 [−0.052, +0.047]	0.87
Best post hoc (legacy PCA-10 stack)	0.721 [0.638, 0.793]	+0.023 [−0.013, +0.057]	0.92	0.754 [0.685, 0.823]	+0.035 [+0.002, +0.068]	0.96
Nested over 70 (honest)	**0.666 [0.586, 0.739]**	**−0.032 [−0.084, +0.017]**	0.60	**0.747 [0.678, 0.811]**	**+0.028 [−0.017, +0.071]**	0.85

Median ΔAUC by variant, grade 4: legacy +0.005, v2 all four sequences −0.016, v2 compartments −0.017, v2 lesion −0.001; all gliomas +0.024, +0.000, −0.005, +0.005. Nested choices: legacy in 18/25 (grade 4) and 25/25 (all) outer folds. Stacking beat concatenation in grade 4 (median Δ −0.003 vs −0.027).

S6 Frozen deep features

Source: `docs/results/DEEP_FEATURES.md`; `outputs/results/analyses/deep/`.

Encoders. MONAI SwinUNETR (four channels FLAIR / T1c / T1 / T2, feature size 48), encoder only: BraTS 2021 weights; BrainSegFounder (UK Biobank self-supervised pretraining, BraTS fine-tuning; Cox et al., 2024 [CHECK]); and random weights (seed 0) as the control. **Input.** Skull-stripped 1 mm volumes, z-scored within the brain, reoriented to LPS, zero-padded to a multiple of 32; eight flip test-time augmentations. **Features.** Mask-weighted mean pooling at stages 0–3 (48 + 96 + 192 + 384 = 720 per region); MU regions lesion, ET, NETC, SNFH, 5 mm rim and ET ∪ SNFH; LUMIERE regions ET, SNFH, ET ∪ SNFH and lesion. The 768-d bottleneck was not used (between-scan SD 3×10^{-4} for BraTS 2021 weights). **Families.** D1 pretrained, 32 configurations (2 encoders × {lesion, compartments} × 4 selections × 2 integrations); D0 random, 16; D1b deep + legacy radiomics stacked, 16; DS shared regions, 16. **Licences.** BraTS 2021 SwinUNETR weights Apache-2.0; BrainSegFounder weights under the UK Biobank material transfer agreement (not redistributed).

Supplementary Table S10. Deep features on MU.

Arm	Grade 4 AUC	Grade 4 Δ [95% CI]	All AUC	All Δ [95% CI]
Baseline	0.698 [0.615, 0.774]	—	0.720 [0.638, 0.792]	—
D1 pretrained, nested over 32	0.694	−0.004 [−0.069, +0.063]	0.731	+0.011 [−0.032, +0.056]
D0 random weights, nested over 16	0.649	−0.049 [−0.085, −0.014]	0.711	−0.009 [−0.032, +0.009]
D1b deep + legacy radiomics, nested over 16	0.709	+0.011 [−0.032, +0.054]	0.751	+0.031 [−0.003, +0.066]
DS shared regions, nested over 16	0.699	+0.001 [−0.057, +0.060]	0.725	+0.005 [−0.035, +0.046]
Baseline + volumes + D1 vs baseline + volumes	0.694 vs 0.702	−0.008 [−0.045, +0.033]	0.739 vs 0.733	+0.007 [−0.021, +0.036]

Single configurations with CI above 0: 0/80 (grade 4); 3/80 (all gliomas; +0.035 to +0.040, lower bounds +0.001 to +0.004); about 2 of 80 expected by chance. Image-only stacked score, grade 4: pretrained 0.652 (modal configuration), random 0.566; correlation with log tumour volume r = 0.58 and with the baseline score r = 0.28. Calibration slope of nested D1: 0.55 (grade 4),

0.79 (all).

S7 PRISM v1: the initial radiomics model, the initial evaluation split and the fusion components

Source: `docs/results/SHAP_PRISM_SPLIT.md`; `docs/manuscript/FIGURES.md` (Figure 1); `outputs/results/analyses/ shap_prism_split/`; `outputs/results/analyses/paper_figures/single_split{,_pairs}.csv`.

Reconstruction. The input matrix was rebuilt from the legacy feature files plus the clinical codes using PRISM v1's case table; the shipped bundle reproduced the stored test predictions on all 84 rows (maximum absolute difference 1.1×10^{-15} calibrated, 1.4×10^{-15} raw). The model is an L2 logistic regression (liblinear, C = 0.0087, balanced class weights) on 46 standardised columns (two of 48 selected columns are dropped by a correlation filter), with a monotone Platt calibrator. Exact linear SHAP reproduced the shipped importance table (Spearman 0.997, identical top 10). Modality shares: T1c 52%, FLAIR 26%, clinical codes 22% (share of mean absolute per-scan group sum).

Supplementary Table S11. Composition of the test set of our initial evaluation split (percentile among 500 design splits).

	Full cohort	Initial split test set	500 splits: mean (2.5–97.5%)	Percentile
Scans / patients	308 / 155	82 / 29	84.5 / 29.8	
Event rate, naive	0.325	0.268	0.254 (0.165–0.348)	62
Grade 4 / GBM	0.81 / 0.70	0.80 / 0.77	0.82 / 0.63	44 / 97
IDH-mutant scans	0.195	0.146	0.271 (0.131–0.403)	5
Pre-RT scans	0.373	0.439	0.353 (0.220–0.474)	89
Event rate in pre-RT scans	0.087	0.000 (0 / 36)	0.044 (0–0.138)	15
$\leq$ 12 weeks post-RT	0.338	0.244	0.283	16
Later post-RT	0.263	0.268	0.345 (0.243–0.450)	6
AUC of the rule "scan is post-RT"	0.677	0.740	0.675 (0.575–0.772)	92

Supplementary Table S12. Pair decomposition (naive AUC) (`single_split_pairs.csv`; 22 positives, 36 pre-RT negatives, 24 other negatives; 60% of pairs involve a pre-RT negative).

Model	All pairs	Positives vs pre-RT negatives	Positives vs other negatives
PRISM initial radiomics model	0.802	0.823	0.771
Baseline ridge	0.871	0.962	0.735
Baseline GBM	0.844	0.933	0.710
Baseline + legacy radiomics	0.862	0.934	0.754
Clinical only (ridge)	0.791	0.789	0.793

On the 36 pre-RT scans the initial radiomics model's T1c features lowered the log-odds by −0.50 on average; the baseline ridge did the same through treatment (−0.38) and timing (−0.15). Our models' SHAP profiles on this split match their whole-cohort out-of-fold profiles at group level (baseline ridge: treatment 55%, clinical 29%, timing 16% on the split; 52 / 29 / 19% out of fold); the split does not lean on unusual features.

Post-RT test scans (42 scans, 19 patients, 19 positives; IPCW AUC [95% CI], naive in brackets): initial radiomics model 0.770 [0.604, 0.907] (0.778); baseline ridge 0.718 [0.481, 0.880] (0.730); baseline GBM 0.699 [0.463, 0.846] (0.705); + legacy radiomics 0.758 [0.605, 0.886] (0.769); clinical only 0.818 [0.664, 0.924] (0.826), Δ vs baseline +0.100 [−0.015, +0.293].

Before and after the as-of-scan fix (30 test patients / 84 scans, naive / IPCW; `head_to_head_{before,after}.csv`): initial radiomics model 0.802 / 0.811 in both; baseline ridge 0.870 / 0.844 → 0.880 / 0.860; baseline boosting 0.901 / 0.881 → 0.869 / 0.855; baseline + legacy radiomics 0.869 / 0.848 → 0.871 / 0.852; baseline + corrected compartments 0.869 / 0.843 → 0.886 / 0.870. With the final labels (29 / 82): ridge 0.871 / 0.851, boosting 0.844 / 0.824.

PRISM v1 encoder and fusion components (source: `legacy/reproduction/{README,FINDINGS,AUDIT}.md`; v1's own case table, labels and scoring on the 30-patient / 84-scan test set). All v1 numbers reproduced from the code. The v1 image encoder was a small SwinUNETR-style transformer with random, untrained weights; its embeddings alone gave AUC about 0.50 in fusion (linear probe 0.55 in training CV, 0.62 on the test set). The v1 late-fusion MLP scored AUC 0.8006, below its radiomics input alone (0.8043); its checkpoint had been selected on test loss and its 0.48 threshold hand-tuned, so its reported recall and F1 gains over the radiomics model were an operating-point shift. A BraTS-pretrained SwinUNETR fine-tuned on MU (lesion-masked pooling) reached image-only test AUC 0.64–0.65, and late fusion with the radiomics score gave 0.790–0.809, with a patient-bootstrap CI of the difference spanning 0. These results motivated the frozen-encoder design with a random-weight control (S6) and the single regularised model with ablatable branches (main §2.1). The v1 report generator's RANO section was computed from the radiomics score, not from mask measurements; the rebuilt generator is described in S19.

S8 Split variance: full tables and verifier checks

Source: `docs/results/SPLIT_VARIANCE.md`; `docs/manuscript/FIGURES.md` §2 (clipped `baseline_rad`); `outputs/results/analyses/split_variance/{summary,summary_clipped,per_split_auc_clipped}.csv`.

Design. Seeds 0–499 of PRISM v1's split search gave 500 distinct test sets (82–94 test scans, mean 86.1; 10–33 positives). The search favours patients with many scans: all 157 patients of PRISM v1's case table appear, and the most frequent is in 71% of splits. Cohort, label and

features are PRISM v1's (315 scans / 157 patients; naive label, equal to ours on all 314 shared scans). Rank = share of splits with strictly lower AUC than the model's own seed-62 value re-run here.

Supplementary Table S13. Naive AUC over 500 splits.

Model	Mean ± SD	2.5–97.5%	Seed 62 (rank)	≥ 0.802	Beats baseline	Mean Δ vs baseline
Initial radiomics pipeline, re-selected per split	0.720 ± 0.055	0.610–0.819	0.801 (92%)	7.2%	12%	−0.081
Initial radiomics model, shipped features refitted (optimistic)	0.815 ± 0.044	0.726–0.887	0.802 (38%)	62%	58%	+0.014
Baseline ridge	0.801 ± 0.052	0.694–0.895	0.871 (91%)	50%	–	–
Baseline + legacy radiomics (clipped)	0.799 ± 0.053	0.674–0.892	0.862 (89%)	50.2%	51.6%	−0.002

Supplementary Table S14. IPCW AUC over 475 splits.

Model	Mean ± SD	Seed 62 (rank)	Beats baseline	Mean Δ
Initial radiomics pipeline, re-selected	0.727 ± 0.058 (0.602–0.832)	0.816 (94%)	25%	−0.042
Shipped features refitted (optimistic)	0.825 ± 0.045 (0.737–0.901)	0.817 (40%)	81%	+0.055
Baseline ridge	0.769 ± 0.056 (0.654–0.876)	0.851 (92%)	–	–
Baseline + legacy radiomics (clipped)	0.774 ± 0.058	0.845 (92.0%)	56.8%	+0.005

Per split, clipped minus pre-clip naive AUC of the radiomics arm: mean −0.002, SD 0.009. Gap between honest and optimistic versions of the initial radiomics pipeline: 0.095 ± 0.040.

Composition. Baseline ridge per-split AUC correlates with the pre-RT event rate (Spearman −0.61 naive, −0.57 IPCW), the AUC of the "post-RT" rule (+0.57, +0.51), the later post-RT share (−0.26) and the pre-RT share (+0.22). Nine composition variables explain CV R^2 0.42 and predict 0.851 for seed 62 (observed 0.871, mean 0.801). For the initial radiomics pipeline: CV R^2 0.10 (naive), 0.15 (IPCW); strongest correlates IDH-mutant fraction (−0.30) and the "post-RT" rule (+0.22); predicted 0.773 for seed 62 (observed 0.801, mean 0.720).

Verifier checks (earlier per-split table; not repeated on the final run). (1) On seed 62 and 15 other splits, redoing the ranking at full cost (10 folds × 100 permutation repeats) gave 0.718 ± 0.054 on the 15 splits (cheap ranking 0.726 ± 0.057; cheap − full +0.008 ± 0.033, r = 0.83). Re-running the shipped Optuna search as well gave 0.654 ± 0.054 (−0.064 ± 0.055 against full ranking with the shipped C); its inner-CV objective averaged 0.865 (optimism ≈ 0.21), and on seed 62 it gave 0.762, not 0.802. (2) Against 300 fully random 30-patient splits, the design splits had more test scans (85.5 vs 59.9) and a narrower baseline AUC spread (SD 0.054

vs 0.069), so spread under the design underestimates what a fresh split would show (Varoquaux, 2018).

S9 Estimator and evaluation details

Source: `docs/results/FIX_EVALUATION.md`.

- **Headline estimator.** Mean over the 5 repeats of the per-repeat pooled-OOF IPCW AUC (and Brier, O/E, slope). The earlier estimator — the AUC of the OOF prediction averaged over repeats — is an ensemble of five models per scan and is optimistic, more so for unstable nested pipelines. On intermediate inputs, the nested radiomics ΔAUC moved from $+0.009$ (repeat-averaged) to -0.002 (mean over repeats) under the change.
- **Bootstrap.** Patient-cluster bootstrap, 1,000 draws; each draw resamples patients once and recomputes the mean-over-repeats statistic of every model on that draw, so differences are paired. ΔAUC is the point estimate (not the bootstrap mean) with the bootstrap CI.
- **Calibration.** O/E (Kaplan–Meier risk / mean prediction, or IPCW-weighted), calibration-in-the-large with a logit offset, calibration slope by IPCW-weighted logistic regression on the logit, flexible calibration curves (restricted cubic spline, 3 knots) and ICI (Van Calster et al., 2019).
- **External estimator.** Every LUMIERE arm is scored on the same 5×5 patient-grouped LUMIERE folds (per-fold IPCW AUC averaged over 25 folds, censoring weights from the whole LUMIERE cohort); locked models are also scored pooled. An earlier pooled score of differently recalibrated folds produced an artefactual AUC of 0.414 for recalibrated ridge; recalibration leaves discrimination unchanged by construction.
- **Selection inside folds.** Feature selection, PCA, ICC filtering, stacking and hyperparameter choice are all inside the training fold, with inner patient-grouped CV for every choice (Varma & Simon, 2006; Cawley & Talbot, 2010; Demircioğlu, 2021).

S10 Feature-group search with permutation control

Source: `docs/results/C2_feature_search.md`; `docs/results/FINAL_RESULTS.md`.

Six groups — clinical, timing, volumes, morphology, change, legacy radiomics (top 20 in-fold; unclipped) — give 63 combinations $\times$ 2 learners (ridge; fixed shallow boosting). The treatment block is not part of this search. The honest estimate selects one of the 126 candidates per outer fold by inner CV. The permutation control moves each patient's whole outcome block to

another patient with the same number of scans, keeps features and folds, and re-runs the full nested pipeline 100 times; p = (1 + #{perm ≥ obs}) / 101.

	Grade 4	All gliomas
Post-hoc best configuration (optimistic)	boosting, clinical + timing: 0.714 [0.643, 0.785]	ridge, clinical + timing + volumes + morphology + radiomics: 0.744 [0.679, 0.804]
Nested over 126 (honest)	0.664 [0.601, 0.722], slope 0.49	0.707 [0.647, 0.766]
Permutation null mean (95th percentile)	0.548 (0.636)	0.528
Permutation p	0.050 (4/100 ≥ observed)	0.010 (0/100)

Single groups, grade 4 (ridge / boosting): radiomics 0.675 / 0.628; morphology 0.650 / 0.538; volumes 0.611 / 0.573; clinical 0.585 / 0.612; timing 0.559 / 0.649; change 0.537 / 0.565. The permutation null is centred above 0.5 because selecting the best of 126 configurations is optimistic even without signal. Boosting was chosen in 15/25 outer folds; selected boosting models had large train–test gaps (for example training AUC 0.961 for the best grade 4 configuration).

S11 Phase-of-care ladder: full results

Source: `docs/results/PHASE_LADDER.md`; `outputs/results/analyses/phase_ladder/{results, cohorts,permutation, subset_of_full,nested_choices}.csv`; `docs/manuscript/figures/fig4_phase.csv`.

Phase definitions (grade 4 / all-glioma scans): pre-RT, `rt_started == 0`, including patients with no RT documented (76 / 115); during RT, started but not completed (6 / 8); early post-RT, completed and ≤ 84 d after RT end (98 / 104); later post-RT, > 84 d (69 / 81). RT without a documented start date counts as started; RT without an end date is never "completed".

Supplementary Table S15. Cohorts.

Population · cohort	Scans	Patients	Events	Non-events	Censored	EPP R2 (10 params)	EPP R2 + I2 (54)
Grade 4 · full	249	126	93	117	39	9.3	1.7
Grade 4 · post-RT	167	84	83	83	1	8.3	1.5
Grade 4 · later post-RT	69	40	36	32	1	3.6	0.7
All · full	308	155	100	156	52	10.0	1.9
All · post-RT	185	92	84	96	5	8.4	1.6
All · later post-RT	81	46	37	40	4	3.7	0.7

By the Riley criteria (anticipated R^2cs 15% of maximum), the primary post-RT cohort would need about 750 scans (375 events) for R2 and about 4,040 scans (2,022 events) for R2 + I2. Nearly all censoring is in pre-RT and early scans. In post-RT cohorts two phase dummies are constant, so R2 has about 8 effective parameters (exploratory note).

Supplementary Table S16. Primary and key secondary results (grade 4, post-RT, ridge).

Model	AUC (95% CI)	Δ vs R2 (95% CI)	Slope	O/E
R2 (clinical + phase)	0.549 (0.458–0.642)	—	0.76	0.99
R2 + I2 (primary)	0.610 (0.529–0.688)	+0.061 (−0.038, +0.156)	0.68	0.99
R2 + I3 (key secondary)	0.609 (0.543–0.676)	+0.061 (−0.041, +0.157)	0.44	0.96
R2 + I1	0.593	+0.044 (−0.054, +0.137)	0.60	0.98
R2 + I4	0.625	+0.076 (−0.041, +0.192)	0.30	0.97
R3 (baseline), reference	0.604 (0.520–0.690)	—	0.85	0.99

Permutation test of the primary contrast: p = 0.208 (20 of 100 permutations ≥ +0.061; p = 21/101); null mean +0.028, median +0.025, range −0.104 to +0.162. The nested radiomics choice was unstable (most common configuration in 12 of 25 outer folds).

Supplementary Table S17. Bases by cohort (grade 4, ridge; AUC [95% CI]) (`fig4_phase.csv`).

Base	Full	Post-RT	Later post-RT
R0 clinical	0.585 [0.496, 0.666]	0.552 [0.456, 0.646]	0.343 [0.281, 0.418]
R1 + timing	0.644 [0.560, 0.721]	0.572 [0.485, 0.667]	0.388 [0.291, 0.503]
R2 + phase	0.664 [0.569, 0.753]	0.549 [0.458, 0.642]	0.376 [0.294, 0.470]
R3 baseline	0.698 [0.615, 0.774]	0.604 [0.520, 0.690]	0.472 [0.339, 0.617]

Full grade 4 cohort: R1 − R0 +0.059 (−0.016, +0.133); R2 − R0 +0.080 (+0.005, +0.152); R3 − R0 +0.113 (+0.026, +0.208); R3 − R2 +0.034 (−0.027, +0.090). R2 + I1 0.696 vs R3 0.698 (Δ −0.002, −0.081 to +0.074). Imaging on top of each base (full grade 4; I1 / I2 / I3 / I4): R0 +0.074 / +0.071 / +0.065 / +0.063; R1 +0.030 / +0.026 / +0.008 / +0.018; R2 +0.031 / +0.023 / +0.002 / +0.020; R3 +0.004 / −0.007 / −0.032 / −0.004; no CI excludes 0. Post-RT grade 4 (I1 / I2 / I3 / I4): R0 +0.049 / +0.058 / +0.046 / +0.079; R1 +0.030 / +0.038 / +0.042 / +0.060; R2 +0.044 / +0.061 / +0.061 / +0.076; R3 +0.008 / +0.017 / +0.005 / +0.039; no CI excludes 0. All gliomas, full: R0 0.619, R1 0.693, R2 0.700, R3 0.720. Boosting (secondary): grade 4 full R2 0.698, R1 0.693; imaging never helped boosting (−0.027 to +0.007); post-RT R2 + I2 vs R2 +0.012 (−0.069, +0.102).

Sensitivity analyses. Later post-RT only (grade 4: 69 scans, 40 patients, 36 events): bases below chance (R0 0.343, calibration slope −2.0), the known negative bias of CV for near-null models with few events; large ΔAUCs there (e.g. R2 + I3 +0.250) are measured against a below-chance reference and are not evidence. Full-cohort models evaluated on post-RT scans only: grade 4 R2 0.570, R2 + I2 0.620, Δ +0.050 (−0.061, +0.157); R3 0.638, R3 + I2 Δ −0.014. Exploratory: age alone 0.586 in grade 4 post-RT; age + I2 vs age +0.022 (−0.103, +0.145). LUMIERE could not be used: no RT dates; the rater flag marks 27 of 132 scans and leaves 83 scans with unknown phase.

S12 Secondary analyses

S12.1 Literature baselines and tuning

Source: `docs/results/BASELINES.md`.

Supplementary Table S18. MU ladder (reference = baseline ridge).

Model	Grade 4 AUC [95% CI]	Δ [95% CI]	All AUC [95% CI]	Δ [95% CI]
Baseline ridge (reference)	0.698 [0.615, 0.774]	—	0.720 [0.638, 0.792]	—
Clinical only	0.585 [0.496, 0.666]	−0.113 [−0.208, −0.026]	0.619 [0.540, 0.696]	−0.101 [−0.177, −0.031]
Timing only	0.559 [0.475, 0.637]	−0.139 [−0.214, −0.070]	0.607 [0.528, 0.684]	−0.112 [−0.175, −0.059]
Clinical + timing	0.644 [0.560, 0.721]	−0.054 [−0.103, −0.012]	0.693 [0.617, 0.766]	−0.027 [−0.061, +0.007]
B0 log ET volume	0.636 [0.541, 0.723]	−0.062 [−0.183, +0.043]	0.673 [0.591, 0.752]	−0.046 [−0.142, +0.054]
B0 ΔET vs previous	0.546 [0.479, 0.614]	−0.153 [−0.241, −0.056]	0.531 [0.465, 0.590]	−0.189 [−0.285, −0.101]
B1 volumetric RANO-style rule	0.551 [0.480, 0.620]	−0.147 [−0.239, −0.062]	0.554 [0.488, 0.615]	−0.166 [−0.258, −0.087]
B1 rule, RANO 2.0 window	0.508 [0.440, 0.577]	−0.191 [−0.289, −0.098]	0.509 [0.448, 0.568]	−0.210 [−0.301, −0.119]
B4 Hashim-style Cox	0.628 [0.538, 0.711]	−0.070 [−0.179, +0.031]	0.678 [0.596, 0.755]	−0.041 [−0.120, +0.039]

The B1 rule: ET $\geq 40\%$ and ≥ 1 cc above the past-only nadir (measurable nadir ≥ 0.1 cc), or ≥ 1 cc from a $\sim$0 cc nadir; positive in 25% of grade 4 scans. Nested hyperparameter grids: every tuned arm within ± 0.025 of the baseline, all CIs include 0 (e.g. nested over ridge + boosting -0.024 [-0.061, $+0.011$], grade 4); choices were unstable.

S12.2 Sensitivity of the primary comparison

Source: `docs/results/SENSITIVITY.md`.

Sensitivity	Scans / patients	Events / censored	Baseline AUC [95% CI]	+ radiomics (nested)	ΔAUC [95% CI]
Primary: grade 4, 120 d	249 / 126	93 / 39	0.698 [0.615, 0.774]	0.666	−0.032 [−0.084, +0.017]
Horizon 90 d	249 / 126	62 / 35	0.663 [0.577, 0.739]	0.664	+0.001 [−0.057, +0.063]
Horizon 180 d	249 / 126	131 / 43	0.667 [0.573, 0.752]	0.636	−0.032 [−0.092, +0.022]
Composite (progression or death)	249 / 126	107 / 25	0.642 [0.566, 0.713]	0.629	−0.013 [−0.073, +0.052]
Alternative progression date	244 / 124	92 / 39	0.670 [0.578, 0.757]	0.648	−0.023 [−0.066, +0.019]
Imaging-only follow-up	249 / 126	93 / 43	0.685 [0.600, 0.761]	0.667	−0.018 [−0.070, +0.033]
All gliomas	308 / 155	100 / 52	0.720 [0.638, 0.792]	0.747	+0.028 [−0.017, +0.071]
Excluding 12-week post-RT scans	151 / 90	46 / 39	0.725 [0.602, 0.834]	0.727	+0.002 [−0.064, +0.067]
Excluding patient 0145 (conflicting dates)	247 / 125	93 / 39	0.691 [0.602, 0.776]	0.687	−0.004 [−0.058, +0.048]

S12.3 Class imbalance

Source: `docs/results/IMBALANCE.md`. Event shares are 44% (grade 4) and 39% (all gliomas), a mild imbalance. Over learners (ridge, boosting, XGBoost, LightGBM, nested variants) × re-balancing (none, class weights, patient-aware oversampling, SMOTE with neighbours from other patients), ΔAUC against the ridge baseline ranged from −0.031 to +0.023; 4 of 190 CIs excluded 0 (ridge, all gliomas, +0.009 to +0.014). Re-balancing shifted predicted risk upward as expected from a prior shift (median O/E 1.01 without re-balancing vs 0.88–0.89 with it) and raised Brier. Grade 4 ridge with class weights: ΔAUC +0.008 [−0.001, +0.019], O/E 1.04 → 0.94. Fixed small-data XGBoost: +0.006 [−0.060, +0.072] (grade 4); nested-tuned XGBoost had calibration slope about 0.6 and transported worse to LUMIERE (0.562 vs 0.608). No re-balancing is used in any reported model.

S12.4 Schedule coupling

Source: `docs/results/SCHEDULE_COUPLING.md`. The interval to the next MRI is unknown at prediction time and partly caused by the outcome (scans are brought forward when progression is suspected), so conditioning on it is a conservative test. Grade 4: positive rate 0.54 when the next scan is $\leq$ 60 d away and 0.10 when $>$ 180 d; interval alone AUC 0.639 [0.555, 0.721]. Baseline AUC within interval tertiles 0.718 / 0.762 / 0.617; conditional AUC (pairs within a tertile) 0.687 [0.595, 0.768]; logit coefficient per SD 0.91 → 0.86 [0.43, 1.41] after adjusting for log interval; baseline features cannot predict the interval (grouped-CV R^2 −0.05). Excluding early post-RT scans and scans whose next scan is $>$ 120 d away: 0.777 [0.653, 0.891] (85 scans). Nested radiomics conditional AUC 0.654, below the baseline. In LUMIERE the interval alone scores 0.788 [0.703, 0.868], and the + volumes boosting model's coefficient is no longer distinguishable from 0 after adjustment (0.37 [−0.14, 0.89]).

S12.5 Feature conflicts

Source: `docs/results/FEATURE_CONFLICTS.md`. No feature group measurably hurts: every leave-one-group-out contrast that would mean "group hurts" has a CI including 0; adding any imaging group to the baseline changes AUC by -0.004 to $+0.019$ (CIs span 0). The conflict is redundancy: timing and treatment encode the same RT-status information (interaction -0.066 to -0.073, CI excluding 0 in 3 of 4 population $\times$ learner cells). Within the baseline, 9 design columns are exactly collinear (VIF $= \infty$) and 16 have VIF > 10; "RT completed" appears in at least five columns. Pre-stated pruning evaluated nested: ΔAUC -0.018 to $+0.011$, all CIs include 0; the same rule applied once globally gives $+0.009$ to $+0.018$, which is selection optimism.

S12.6 SHAP attributions

Source: `docs/results/SHAP.md`. Ridge: exact linear SHAP on in-fold standardised features; boosting: TreeSHAP (path-dependent), additivity checked to 10^{-6}; explanations out of fold, 25 fits. Group share = mean |per-scan group sum| / sum over groups.

Model (grade 4)	OOF AUC	Group shares
Baseline ridge	0.698	treatment 58%, clinical 25%, timing 17%
Baseline boosting	0.687	treatment 51%, clinical 35%, timing 14%
Full non-radiomics ridge	0.692	treatment 34%, morphology 21%, change 15%, clinical 13%, volumes 10%, timing 8%
Full non-radiomics boosting	0.694	treatment 26%, morphology 23%, clinical 21%, change 13%, volumes 13%, timing 3%
Baseline + radiomics score (post hoc configuration)	0.721	treatment 48%, clinical 20%, timing 16%, radiomics 16%

Time-relative features hold 35% (ridge) / 47% (boosting) of feature-level credit in the baseline. Imaging takes 45% / 51% of credit in the full models without an AUC gain. On LUMIERE, the locked ridge's under-prediction is driven by `idh_unknown` (mean -0.71 logits; mean |SHAP| 0.77 on LUMIERE vs 0.13 on MU), by lower DeepBraTumIA ET volumes (-0.32 logits) and by a missingness indicator that is never set in LUMIERE.

S12.7 Calibration, decision curves, added value and sample size

Source: `docs/results/CLINICAL_UTILITY.md`.

Model (grade 4)	AUC	O/E (weighted)	CITL [95% CI]	Slope [95% CI]	ICI
Baseline ridge	0.698	0.96	−0.09 [−0.42, 0.28]	0.89 [0.50, 1.43]	0.032
Baseline boosting	0.687	0.94	−0.12 [−0.47, 0.23]	0.82 [0.44, 1.31]	0.039
Baseline + radiomics (nested)	0.666	0.94	−0.13 [−0.46, 0.25]	0.60 [0.33, 0.93]	0.063
Baseline + deep (nested D1)	0.694	0.94	−0.15 [−0.52, 0.28]	0.55 [0.28, 0.91]	0.085

Net benefit (grade 4, IPCW)	10%	20%	30%
Treat all	0.369	0.291	0.189
Baseline ridge	0.372	0.300	0.232
Δ radiomics (nested)	−0.000 [−0.004, 0.004]	−0.001 [−0.013, 0.011]	−0.020 [−0.038, −0.004]
Δ boosting	−0.002 [−0.007, 0.001]	−0.007 [−0.021, 0.006]	−0.011 [−0.039, 0.013]
Δ deep (nested D1)	−0.002 [−0.013, 0.006]	+0.002 [−0.014, 0.018]	+0.001 [−0.023, 0.026]

Added value vs baseline (continuous NRI; IDI): radiomics +0.17 [−0.05, 0.41]; +0.000 [−0.030, 0.029]. Deep +0.48 [0.24, 0.73]; +0.036 [−0.009, 0.082]. The positive NRI reflects upward risk shifts in events by over-fitted models; ΔAUC and Δ net benefit are null (Pencina et al., 2008). All gliomas: radiomics Δ net benefit +0.011 [0.004, 0.020] at 10% and +0.014 [0.001, 0.027] at 20%, with ΔAUC +0.028 [−0.017, +0.071]; nothing similar in grade 4. LUMIERE locked boosting: net benefit ≈ treat-all (0.218 vs 0.222 at 20%).

Sample size (Riley et al., 2020; grade 4). 210 verified scans, 93 events, outcome proportion 0.443, R^2cs 0.119 (from AUC 0.698). Baseline with 31 design columns: 3.0 events per parameter; required 2,190 scans (970 events) by the shrinkage criterion. Adding 10 radiomics features: 2,896; the full radiomics pool (2,477 candidate parameters): 174,942. Requirements assume independent scans, so the shortfall is understated.

S13 External evaluation on LUMIERE: full results

Source: `docs/results/EXTERNAL_LUMIERE.md`; `outputs/results/external_lumiere/results.csv`.

Shared features: age, sex, IDH and MGMT (with unknown), log days since surgery, < 3 months post-RT flag, log ET / necrosis + cavity / SNFH volumes (DeepBraTumIA; 14 of 132 index scans without usable volumes, imputed with MU training medians inside the locked pipeline).

Supplementary Table S19. Models trained on MU grade 4. Pooled locked AUC [95% CI] and O/E; other columns are per-fold means.

Features	Learner	MU CV	Locked, pooled	O/E	Locked, per fold	Fine-tuned (shrink)	Fine-tuned (pooled)	LUMIERE only
Clinical	ridge	0.585	0.610 [0.487, 0.737]	1.15	0.558	0.544	0.505	0.396
Clinical	boosting	0.625	0.627 [0.514, 0.731]	0.86	0.624	—	0.560	0.520
Clinical + timing	ridge	0.612	0.610 [0.497, 0.737]	0.98	0.603	0.591	0.603	0.586
Clinical + timing	boosting	0.664	0.632 [0.501, 0.748]	0.84	0.643	—	0.632	0.627
Volumes	ridge	0.611	0.545 [0.410, 0.653]	1.00	0.521	0.531	0.567	0.569
Volumes	boosting	0.591	0.529 [0.389, 0.644]	1.00	0.515	—	0.513	0.518
Clinical + timing + volumes	**ridge**	0.667	**0.608 [0.469, 0.739]**	**1.33**	0.598	0.606	0.617	0.580
Clinical + timing + volumes	**boosting**	0.675	**0.620 [0.495, 0.733]**	**0.99**	0.608	—	0.658	0.610

Recalibration equals locked per fold (pooled O/E 0.93–0.97). No adaptation arm has a ΔAUC CI above 0; LUMIERE-only is significantly worse for clinical ridge (Δ −0.162 [−0.238, −0.066]). Cohort normalisation of volumes (rank) raised the pooled locked AUC of every volume model (+0.007 to +0.069); clinical + timing + volumes boosting 0.620 → 0.659 [0.529, 0.767], per-fold Δ +0.047 [−0.002, +0.110]. This is transductive and a sensitivity analysis only. All-glioma-trained locked clinical + timing + volumes: ridge 0.599 [0.462, 0.728] (O/E 1.30), boosting 0.619 [0.477, 0.740] (O/E 1.04).

Deep features on LUMIERE (118 scans with deep features; `deep/lumiere/locked.csv`): grade 4 clinical + timing 0.622 → + deep 0.679 (Δ +0.057 [−0.132, +0.235], O/E 2.08); random encoder same configuration 0.667 (+0.045). With volumes: 0.635 → 0.725 (+0.089 [−0.026, +0.215]). All gliomas: +0.064 [−0.011, +0.144]; with volumes +0.050 [+0.002, +0.109]. Pre-trained vs random, same configuration: +0.011 to +0.069, no CI above 0. A classifier separates LUMIERE from MU on deep features with AUC 0.82 / 0.79 / 0.81 (BraTS 2021 / BrainSeg-Founder / random).

PRISM's initial radiomics model, as shipped (legacy features re-extracted, 118 scans; files dated 2026-10-09, before the final LUMIERE label definition): raw 0.613 [0.500, 0.729] pooled (O/E 1.24); recalibrated O/E 0.98; fine-tuned 0.570 pooled.

Literature baselines on LUMIERE (locked; `BASELINES.md`): clinical + timing 0.610 [0.497, 0.737]; Hashim-style Cox 0.640 [0.508, 0.754] (Δ +0.030 [−0.080, +0.127]); volumetric RANO-style rule 0.559.

Harmonisation note. DeepBraTumIA ET and SNFH medians (grade 4: 0.61 and 19.11 cc) are far below MU's (5.45 and 42.85 cc). The harmonised necrosis + cavity definition is correct but did not improve transfer in a development run (`FIX_FEATURES.md` §1).

S14 Imaging-defined label, imaging diagnostics and re-segmentation

Source: `docs/results/CLEAN_LABEL.md`; `docs/results/DIAGNOSTICS_IMAGING.md`; `docs/results/RESEGMENTATION.md`; `outputs/results/analyses/{clean_label,diagnostics_imaging,reseg}/`.

Imaging-defined label. MU codes the type of first progression: imaging, clinical only, or both. Censoring the clinical-only progressions at their progression date changed grade 4 events from 93 (44 / 12 / 37 by type) to 81 and censored scans from 39 to 51; no non-event changed. Primary (grade 4, full, ridge): clinical 0.563 (0.474–0.642); baseline 0.696 (0.610–0.776); imaging (volumes + change) 0.628 (0.549–0.701), Δ −0.068 (−0.167, +0.033); baseline + imaging 0.722 (0.639–0.799), Δ +0.027 (−0.033, +0.082). On the original label, baseline + imaging was 0.694 (Δ −0.004, −0.060 to +0.051). Secondary contrasts (post-RT, all gliomas, boosting) gave baseline

+ imaging Δ +0.008 to +0.028 with all CIs including 0; imaging alone in all gliomas was worse than the baseline (−0.090, −0.176 to −0.007). Within-LUMIERE CV with the expert RANO label: every model at or below chance (ridge clinical + timing 0.466), so LUMIERE cannot answer the label question. Rule not met.

Volume cross-check against the dataset's own volume table (pass). Pearson r: ET 0.9998 (exact voxel match 98.3%), SNFH 0.9994 (99.7%), RC 0.9993 (94.1%), NETC 0.9957 (83.7%); disagreements are concentrated in seven patients with extra timepoints or a second mask version in the table. Measurement volumes equal mask voxel counts exactly. Label mapping: diagonal agreement 0.94–0.97, off-diagonal ≤ 0.003. Intensities agree with labels (ET/brain on T1c median 1.45, > 1 in 98% of scans; SNFH/brain on FLAIR median 1.44, > 1 in 100%).

Concurrent detection (positive control; fail). Progression scan = first MRI with a mask on or after the documented progression date (120 scans, all gliomas).

Comparison (imaging only, 5 × 5 grouped CV; ridge / boosting)	All gliomas	Grade 4
Progression scans vs all index scans	0.62 [0.57, 0.67] / 0.63	0.62 / 0.63
Progression scans vs verified 120-d negatives	0.68 [0.62, 0.74] / 0.70	0.65 / 0.66
Both groups with a prior scan	0.62 / 0.61	0.60 / 0.62

Progression scan vs the same patient's previous scan (74 pairs): ET larger in 49% (Wilcoxon p = 0.54); lesion larger in 65% (p = 0.033); lesion growth from nadir larger in 79% (n = 42, p = 0.0003); ET growth from nadir 50% (p = 0.95). By progression type: imaging (40 pairs) ET larger in 50%; imaging + clinical (27) 52%; clinical only (7) 29%.

Why the control fails. Between consecutive scans ≤ 14 days apart (n = 25), the median change was 3.2-fold in ET and 1.8-fold in lesion volume, larger than over 45–90 day gaps; across 389 pairs the size of the change was unrelated to the gap (Spearman −0.01, p = 0.83). Some short-gap jumps straddle a re-resection; others (four patients) are implausible.

Label–date alignment (pass). The progression date coincides with an MRI for 62% of progressors (±7 d: 69%; 30 d: 73%); 22% of progressors have no MRI on or after the progression date. Positives: median time to progression 66 d (IQR 44–92); 76% are the last scan before progression. Agreement between the 120-d label and "the next scan is the progression scan": 79% ($\kappa = 0.53$).

Horizon curve (grade 4, IPCW AUC(H); imaging / clinical + phase / both). 60 d: 0.63 / 0.55 / 0.65; 90 d: 0.62 / 0.63 / 0.68; 120 d: 0.60 / 0.66 / 0.69; 180 d: 0.58 / 0.59 / 0.63; 270 d: 0.59 / 0.55 / 0.63. Imaging stays at about 0.6 at every horizon; 30-d estimates (16 events) are not interpretable.

Data issues to report to the data owners. Three scan pairs from different patients have identical masks (0018_T5 / 0032_T5, 0060_T2 / 0062_T2, 0113_T1 / 0132_T1) and four

further pairs are near-identical; a few implausible within-days volume jumps; 15 of 96 recorded deaths precede a dated MRI (S3).

Re-segmentation. BraTS 2021 SwinUNETR and BrainSegFounder were run on all 594 complete scans (sliding window, ROI 128^3, overlap 0.5; labels written in the MU convention; RC taken from the MU mask where the new map is background, because BraTS models have no cavity class and label most of the cavity as NETC, so tumour core is not comparable).

Agreement with MU masks	Dice ET median [IQR]	Dice WT	Pearson log-volume ET / WT
BraTS 2021 SwinUNETR	0.80 [0.49, 0.91]	0.85 [0.74, 0.93]	0.92 / 0.91
BrainSegFounder	0.81 [0.56, 0.91]	0.86	0.94 / 0.92

Short-gap consistency ($\leq$ 30 d, 54 pairs): median ET fold change 2.57 $\rightarrow$ 2.17 (Δ median |log ratio| -0.17 [-0.46, -0.02]); WT unchanged. Positive controls with the new masks: primary detection (progression scan vs verified negatives, ridge) 0.681 $\rightarrow$ 0.665, Δ -0.016 [-0.057, $+0.025$] (BrainSegFounder 0.677); ET larger at progression 49% $\rightarrow$ 45% (BrainSegFounder 41%). Forward 120-d task, imaging only: all gliomas 0.644 $\rightarrow$ 0.675, Δ $+0.031$ [-0.017, $+0.084$]; clinical + phase + imaging 0.712 $\rightarrow$ 0.740, Δ $+0.029$ [-0.010, $+0.069$]. An exploratory plain-AUC boosting gain ($+0.059$ [$+0.014$, $+0.107$]) was one of more than 20 comparisons, not pre-specified and not confirmed by the IPCW ridge estimate. Verdict by the pre-specified rule: the new segmentation does not help.

S15 The auditable layer: full results and an example audit record

Source: `docs/results/AUDIT_SYSTEM.md`; `outputs/results/analyses/audit_system/`. Run: 2026-10-15, `python -m prism audit --workers 24`, 114 s. Checks (`checks.csv`): CV predictions reproduce the baseline OOF predictions (max $|\Delta|$ 1.7×10^{-16}); intercept + Σ contributions = logit (max error 1.3×10^{-15}). Code changes after the plan were confined to an as-of rule for demographics, a NaN rule for one-class strata, and a shared estimator for referral subsets; none changed a model or a rule.

Locked artifact and model card. Grade 4: 249 training scans, 126 patients, 210 verified, 93 events; C = 0.1; conformal threshold q at $\alpha = 0.2$ / 0.1: 0.613 / 0.683 (all gliomas 0.529 / 0.620). Out of scope: pseudoprogression diagnosis; cohorts without treatment fields (including LUMIERE); other institutions without validation.

Supplementary Table S20. Example audit records (out-of-fold, repeat 0, grade 4; label shown for evaluation only).

Scan	Phase	Risk [95% bootstrap]	Conformal set $\alpha = 0.2$	Flags	120-d label
0201_T3	pre-RT	0.04 [0.01, 0.27]	{0}	OOD (MGMT unknown)	0
0136_T2	during RT	0.47 [0.36, 0.77]	{0}	none	1
0117_T1	early post-RT	0.50 [0.39, 0.61]	{0, 1}	(i) 26 d after RT; (iii)	1
0059_T1	early post-RT	0.48 [0.22, 0.77]	{0, 1}	(i); (ii) rt_start_day below training 1st percentile; (iii)	0
0089_T5	later post-RT	0.45 [0.34, 0.72]	{0, 1}	(ii) rt_start_day above 99th percentile; (iii)	1
0014_T4	later post-RT	0.63 [0.44, 0.86]	{1}	none	0 (set misses)

Example record (PatientID_0117_Timepoint_1, day 85 from diagnosis), abridged from `records.md`.

- Phase of care: early post-RT ($\leq$ 84 d after RT end).
- Risk of documented progression within 120 d: 0.50 (95% bootstrap interval 0.39–0.61); conformal set at $\alpha = 0.2$ {0, 1}.
- Referral: REFER — early post-RT window (scan 26 d after RT end); uncertain (the $\alpha = 0.2$ set contains both classes).
- Log-odds: intercept -0.33 + clinical $+0.22$ + timing -0.14 + treatment $+0.24 = -0.01$ (check error 5.6×10^{-17}).
- Inputs known by scan day: yes, all. Examples: `rt_started` = 1 (source event day 21 $\leq$ scan day); `rt_completed` = 1 (day 59); `days_since_rt_end` = 26; prescription 60 Gy in 30 fractions (visible from day 21); `surgery_day` -4.
- Largest terms: `early_post_rt` -0.12; `visit_index` -0.12; `previous_brain_tumor` $+0.07$; `rt_started` $+0.07$; `male` -0.07; `mgmt_unknown` $+0.07$ (log-odds).
- Evaluation only: 120-d label 1 (progression at 111 d).

The displayed risk (model fitted on the whole training fold) and the conformal set (model fitted on 70% of training patients) can disagree in direction (e.g. 0136_T2: risk 0.47, set {0}); this occurred in 1.8% / 2.9% of singleton sets (exploratory). Median width of the 95% bootstrap interval: 0.37 (IQR 0.32–0.44) in grade 4 and 0.31 in all gliomas.

Supplementary Table S21. P1, conformal coverage at $\alpha = 0.2$ (`conformal.csv`).

Population	Overall [95% CI]	Pre-RT	During RT	Early post-RT	Later post-RT	Events / non-events	Singleton rate
Grade 4	0.841 [0.790, 0.885]	0.868 (0/5 events covered)	0.567 (6 scans)	0.871	0.806	0.824 / 0.855	0.595
All gliomas	0.823 [0.776, 0.865]	0.850 (0/10)	0.550 (8 scans)	0.852	0.790	0.698 / 0.904	0.696

Per-patient coverage 0.828 / 0.796. At $\alpha = 0.1$: 0.949 [0.916, 0.975] / 0.928 [0.896, 0.955], singleton rate 0.296 / 0.422. Conformal model AUC 0.667 / 0.709 (cost of the 70/30 split). Displayed-risk AUC on singleton scans 0.732 / 0.723 (non-singletons 0.628 / 0.594).

Supplementary Table S22. P2 and referral rules (`p2.csv`, `referral.csv`, `referral_within_phase.csv`).

	Realised coverage	AUC retained [95% CI]	Composition-matched	Random removal	P2 [95% CI]
Grade 4	0.671	0.731 [0.635, 0.812]	0.725	0.698	+0.006 [−0.023, +0.036]
All gliomas	0.685	0.735 [0.654, 0.820]	0.733	0.720	+0.003 [−0.031, +0.036]

Grade 4 rule	Referred	AUC kept	AUC referred	Random removal	Composition-matched
(i) early post-RT	39.4% (98 scans, 47 events)	0.723	0.663	0.699 [0.673, 0.723]	0.723 (the rule itself)
(ii) out of distribution	29.8%	0.671	0.665	0.698	0.695
(iii) uncertain	40.5%	0.732	0.628	0.699	0.734 [0.709, 0.760]
any of (i)–(iii)	72.6%	0.735	0.687	0.698	0.753

Within-phase AUC (grade 4 / all): pre-RT 0.555 / 0.493; during RT 0.448 / 0.671; early post-RT 0.663 / 0.678; later post-RT 0.597 / 0.644. On OOD-referred scans the model over-predicts (O/E 0.70, slope 0.66).

Shortcut audit. Unknown marker share by 120-d label (progression / no progression / censored), grade 4: IDH unknown 0/93, 6/117, 5/39; MGMT unknown 6/93, 25/117, 8/39. Dropping both unknown indicators: grade 4 0.676 [0.592, 0.752], Δ −0.023 [−0.047, −0.001]; also dropping `rt_date_unknown` Δ −0.022 [−0.045, −0.001]; all gliomas Δ −0.011 [−0.031, +0.008]. Timing only 0.559, Δ −0.139 [−0.214, −0.070]; treatment only 0.701 [0.622, 0.776], Δ +0.003 [−0.050, +0.048] (all gliomas 0.710, Δ −0.009).

Supplementary Table S23. Subgroups (grade 4; † fewer than 10 events, descriptive).

Stratum	Scans / events	AUC [95% CI]	Slope	O/E [95% CI]
GBM	214 / 86	0.68 [0.58, 0.76]	0.78	1.10 [0.90, 1.31]
Non-GBM grade 4 †	35 / 7	0.72 [0.54, 0.95]	1.34	0.66 [0.30, 1.00]
MGMT methylated	88 / 34	0.64 [0.52, 0.77]	0.62	0.99
MGMT unmethylated	122 / 53	0.72 [0.61, 0.84]	1.11	1.14
IDH wild type	208 / 87	0.66 [0.56, 0.77]	0.78	1.09
Male / female	119 / 44, 130 / 49	0.67 [0.55, 0.79] / 0.72 [0.61, 0.83]	0.85 / 0.94	1.02 / 1.06
Age $\leq$ 56.7 / (56.7, 67] / $>$ 67	109 / 30, 68 / 35, 72 / 28	0.71 / 0.64 / 0.68	0.89 / 0.73 / 0.89	0.81 / 1.39 [1.07, 1.77] / 1.06
Early post-RT	98 / 47	0.66 [0.55, 0.76]	1.06	0.95
Later post-RT	69 / 36	0.60 [0.45, 0.74]	0.50	1.03

The middle age tertile is under-predicted (O/E CI excludes 1); uncorrected for 19 strata. Field strength, scanner and race could not be audited (no acquisition metadata in MU).

LUMIERE transport audit (shared-feature ridge, grade-4-trained; reproduces AUC 0.608, O/E 1.33). Largest standardised mean differences MU → LUMIERE: `idh_unknown` +0.97 (4.4% → 40.9%), `log_et_cc` −0.84, `early_post_rt` −0.72, `log_snfh_cc` −0.56. Mean contribution shift: `idh_unknown` −0.68 logits; net linear-predictor shift −0.64. Refit without the molecular unknown indicators: LUMIERE O/E 1.33 → 0.89, slope 0.38 → 0.73, AUC 0.608 → 0.614 (all gliomas O/E 1.30 → 0.96). MU-calibrated conformal coverage on LUMIERE ($\alpha = 0.2$): 0.715 (events 0.340, non-events 0.947).

Supplementary Table S24. Explanation stability (`stability_summary.csv`).

	Grade 4: 25 CV fits	Grade 4: 200 bootstraps	All: CV	All: bootstrap
Kendall τ-b, |coef| vs locked, median (IQR)	0.63 (0.56–0.67)	0.42 (0.36–0.49)	0.54 (0.48–0.60)	0.39 (0.32–0.46)
Top-5 overlap, mean (all 5 identical)	0.67 (4%)	0.52 (1.5%)	0.65 (4%)	0.52 (2%)
Minimum sign agreement over the top 5	1.00	0.92	1.00	0.97
Features with sign agreement $\geq$ 0.9	61%	55%	68%	61%

Locked top 5 (grade 4): `rt_start_day`, `previous_brain_tumor`, `idh_unknown`, `rt_started`, `biopsy_before_resection`. Group shares of mean |contribution|: treatment 51%, timing 25%, clinical 24%. On a LUMIERE refit of the shared-feature model, 9 of 11 coefficients keep the MU sign; `early_post_rt` and the necrosis + cavity volume flip.

S16 Reproducibility

Every stage is a command of one Python package (`python -m prism <stage>`), wrapped by numbered scripts in `scripts/` (`run_all.sh` with a `--to STEP` option). Shared fold assignments are in `outputs/data/folds.csv`; every analysis writes its outputs, a `run_info.json` and a log under `outputs/`. Seeds are fixed (bootstrap, permutations, conformal splits). The audit layer writes a `MANIFEST.json` with SHA-256 hashes of inputs, code and outputs, the git state and package versions (one recorded drift: scipy 1.17.1 installed against 1.18.1 in the lock file). Pretrained weights are downloaded locally and not redistributed. Unit tests cover label rules, as-of-scan treatment features, feature fixes and the repeated-CV estimator. Commands for each analysis are listed at the end of the corresponding result document in `docs/results/`. [PENDING: public code release and tag.]

S17 Reporting checklists cross-referenced to the manuscript

Item wording: TRIPOD+AI (Collins et al., 2024), `docs/manuscript/CHECKLIST_TRIPOD_AI.md`; CLEAR (Kocak et al., 2023 [CHECK]), `docs/manuscript/CHECKLIST_CLEAR.md`. "M" = main text section; "S" = Supplement section.

TRIPOD+AI.

Item	Topic	Where reported
1	Title	Title (development + external evaluation, population, outcome)
2	Abstract	Structured abstract
3a–c	Background, intended use, inequalities	M1; intended use in Key points and S15 model card; inequalities: race not auditable (M4 Limitations, S15)
4	Objectives	M1 final paragraph
5a–b	Data sources, dates	M2.3; accrual dates: [PENDING — relative days only in both datasets; take from dataset papers]
6a–c	Setting, eligibility, treatments	M2.3–2.4, S2; as-of treatment handling M2.5–2.6, S4
7	Data preparation	M2.5, S4, S5, S14
8a–c	Outcome, assessors, blinding	M2.4, S3; MU assessor not documented, LUMIERE expert RANO; no blinding (M4 Limitations)
9a–c	Predictors	M2.5, S4–S6; MU masks from the dataset, LUMIERE automatic
10	Sample size	Fixed by public datasets; Riley calculation S12.7, M3.8
11	Missing data	In-fold imputation with indicators (M2.5, S4)
12a–g	Analysis methods	M2.1 (architecture), M2.6–2.13, S8–S9, S13 (recalibration, locked predictions)
13	Class imbalance	None used; S12.3
14	Fairness	Subgroup audit S15; limits M4
15	Model output	Probabilities, audit record and grounded report (M2.12–2.13, S15, S19); decision curves S12.7; no threshold chosen
16	Development vs evaluation differences	M2.11, S13
17	Ethics	Declarations [PENDING]
18a–f	Funding, conflicts, protocol, registration, data, code	Declarations [PENDING]; plans S1; code S16
19	Patient and public involvement	Declarations (none)
20a–c	Flow, characteristics, comparison with development data	Table 1, S2 (Figure S1, Table S2)
21	Participants and events per analysis	Tables 1, S4, S15; each result table
22	Full model specification	Locked model files and model card (S15) [PENDING: release coefficients]
23a–b	Performance with CIs, subgroups, heterogeneity	M3, Tables 2–3, S11–S15; heterogeneity only MU vs LUMIERE
24	Model updating	S13 (recalibration, fine-tuning)
25–26	Interpretation, limitations	M4
27a–c	Usability, input quality, next steps	S15 (OOD flags, as-of checks), M4

CLEAR (selected items; full table in `CHECKLIST_CLEAR.md`).

Items	Topic	Where reported
1–6	Title, abstract, keywords, background, rationale, objectives	Title, Abstract, Keywords, M1
7–11	Guidelines, ethics, sample size, study nature, eligibility	M2.2, Declarations, S12.7, M2.4, S2
13–18	Data source, overlap, split, imaging protocol, non-radiomic predictors, reference standard	M2.3, S6 (BraTS pretraining overlap risk), M2.7–2.8, S4, S3; MU protocol not distributed
19–24	Segmentation, pre-processing, resampling, discretisation, image types	M2.3, S5, S14
25–28	Extraction method, classes, number, defaults	S5
29–33	Missing data, imbalance, reliability, scaling, dimension reduction	S4, S12.3, S5 (ICC filter), S5
34–37	Algorithm, tuning, confounders, model selection	M2.8, S5, S12.4 (schedule), M2.9 (phase of care)
38–43	Testing, metrics, uncertainty, comparisons, explainability	M2.8, M2.11, S9, S12.6
44–48	Characteristics, flow, feature statistics, performance, comparison	Table 1, S2, S5, M3, Table 3
49–52	Discussion items	M4
53–58	Open science	Declarations, S16 [PENDING: release]

S18 CFB-GBM diffusion analysis

Source: `docs/results/CFB_ADC.md`; `outputs/results/analyses/cfb_adc/{results.csv, rule.json}`; dataset facts `docs/literature/DATASETS.md`.

Data. TCIA CFB-GBM clinical v03, treatment and MRI availability v02, RANO v04, PyRadiomics features v03 on the GTV; no images used. t0 is the RT-planning MRI; t1 lies a median 14 weeks after t0, about 6–8 weeks after the end of RT, inside the 12-week pseudoprogression window. t1→t2 labels: 30 progressions among 107 labelled patients. No IDH, MGMT or progression dates. The RANO label is computed from the GTV volume (RANO sizes equal the PyRadiomics GTV mesh volume, Spearman 1.00).

Design. One row per patient; outcome t1→t2 progressive disease. Feature sets (a) clinical (age, sex, WHO performance status, WHO edition, RT dose, fractions, delay to RT, weeks t0→t1); (b) + log GTV volume at t1 and change t0→t1; (c) + T1-Gd and FLAIR first-order and texture at t1 plus GTV shape; (d) + ADC first-order and texture at t1. Each radiomics block: in-fold 1% clipping, median imputation, variance filter, scaling, top-k univariate F ($k \in \{5, 10\}$); ridge with $C \in \{0.01 \ldots 3\}$; k and C by inner stratified 3-fold CV. 20 repeats of stratified 5-fold CV, identical splits for all sets; patient bootstrap CIs (1,000); 1,000 permutations of the ADC block for the primary contrast.

Supplementary Table S25. Results (AUC = mean over 20 repeats of pooled-OOF ROC AUC).

Set	Conventional cohort (107 patients, 30 events)	ADC cohort (77 patients, 19 events)
(a) clinical	0.501 (0.403–0.596)	0.565 (0.437–0.689)
(b) + GTV volume and change	0.685 (0.569–0.781)	0.713 (0.589–0.814)
(c) + conventional radiomics	0.676 (0.575–0.769)	0.691 (0.576–0.788)
(d) + ADC radiomics	0.696 (0.593–0.785)[a]	0.698 (0.577–0.800)

Contrast	Conventional cohort	ADC cohort
(b) vs (a)	+0.185 (+0.064, +0.293)	+0.148 (+0.031, +0.266)
(c) vs (b)	−0.009 (−0.067, +0.051)	−0.022 (−0.074, +0.025)
(d) vs (c)	+0.019 (−0.006, +0.045)	**+0.007 (−0.019, +0.035)** (primary; permutation p = 0.151; minimal detectable Δ 0.038)

[a] ADC block imputed for 30 patients without ADC at t1; secondary only. **Rule not met.** Clinical features are at chance (no molecular markers and no phase variation: every t1 scan is at the same post-RT stage). Volume is the only informative block and is partly circular with the volumetric label. Conventional texture adds nothing beyond volume; ADC adds nothing measurable beyond conventional radiomics. Gains larger than about 0.04 are unlikely; smaller gains cannot be excluded. Perfusion was not available.

S19 PRISM report generator: design and faithfulness evaluation

Source: `docs/results/REPORT_AGENT.md` [PENDING: REPORT_AGENT results]. The generator takes one audit record (S15) as its only input; every statement in the report carries a pointer to the record field it states. [PENDING: REPORT_AGENT results — generator and prompt, record-to-report schema, faithfulness criteria and scoring, number of records, results table, example reports including failure cases.]

Supplementary figures

- **Figure S1.** Cohort flow (`docs/manuscript/cohort_flow.png`).
- **Figure S2.** Split-variance histograms per model, original analysis figure (`outputs/results/analyses/split_variance/split_variance.png`; pre-clip radiomics arm — superseded by main Figure 2).

- **Figure S3.** Phase-ladder overview (`outputs/results/analyses/phase_ladder/ladder.png`).
- **Figure S4.** Sensitivity forest plot (`outputs/results/analyses/sensitivity/forest.png`).
- **Figure S5.** Calibration curves and decision curves (`outputs/results/analyses/utility/calibration_dca.png`).
- **Figure S6.** Imaging diagnostics: label–date alignment and horizon curve (`outputs/results/analyses/diagnostics_imaging/{alignment,horizon_curve}.png`); visual QA panels (`qa/*.png`).
- **Figure S7.** SHAP summaries for the baseline and full models (`outputs/results/analyses/shap/grade4/*_{bar,beeswarm}.png`).
- [PENDING: export to the journal format.]

References

P. Blanche, J.-F. Dartigues, and H. Jacqmin-Gadda. Estimating and comparing time-dependent areas under receiver operating characteristic curves for censored event times with competing risks. *Statistics in Medicine*, 32(30):5381–5397, 2013. doi:10.1002/sim.5958.

G. C. Cawley and N. L. C. Talbot. On over-fitting in model selection and subsequent selection bias in performance evaluation. *Journal of Machine Learning Research*, 11(70):2079–2107, 2010. URL `https://jmlr.org/papers/v11/cawley10a.html`.

G. S. Collins, K. G. M. Moons, P. Dhiman, et al. TRIPOD+AI statement: Updated guidance for reporting clinical prediction models that use regression or machine learning methods. *BMJ*, 385:e078378, 2024. doi:10.1136/bmj-2023-078378.

J. Cox et al. BrainSegFounder [full title to be completed]. *Medical Image Analysis*, 2024. doi:10.1016/j.media.2024.103301. CHECK.

A. Demircioğlu. Measuring the bias of incorrect application of feature selection when using cross-validation in radiomics. *Insights into Imaging*, 12(1):172, 2021. doi:10.1186/s13244-021-01115-1.

B. Kocak, B. Baessler, S. Bakas, et al. CheckList for EvaluAtion of Radiomics research (CLEAR): A step-by-step reporting guideline for authors and reviewers endorsed by ESR and EuSoMII. *Insights into Imaging*, 14:75, 2023. doi:10.1186/s13244-023-01415-8. CHECK.

M. J. Pencina, R. B. D'Agostino, Sr., R. B. D'Agostino, Jr., and R. S. Vasan. Evaluating the added predictive ability of a new marker: From area under the ROC curve to reclassification and beyond. *Statistics in Medicine*, 27(2):157–172, 2008. doi:10.1002/sim.2929.

R. D. Riley, J. Ensor, K. I. E. Snell, et al. Calculating the sample size required for developing a clinical prediction model. *BMJ*, 368:m441, 2020. doi:10.1136/bmj.m441.

H. Uno, T. Cai, L. Tian, and L. J. Wei. Evaluating prediction rules for t-year survivors with censored regression models. *Journal of the American Statistical Association*, 102(478):527–537, 2007. doi:10.1198/016214507000000149.

B. Van Calster, D. J. McLernon, M. van Smeden, L. Wynants, and E. W. Steyerberg. Calibration: The Achilles heel of predictive analytics. *BMC Medicine*, 17:230, 2019. doi:10.1186/s12916-019-1466-7.

J. J. M. van Griethuysen, A. Fedorov, C. Parmar, et al. Computational radiomics system to decode the radiographic phenotype. *Cancer Research*, 77(21):e104–e107, 2017. doi:10.1158/0008-5472.CAN-17-0339.

S. Varma and R. Simon. Bias in error estimation when using cross-validation for model selection. *BMC Bioinformatics*, 7:91, 2006. doi:10.1186/1471-2105-7-91.

G. Varoquaux. Cross-validation failure: Small sample sizes lead to large error bars. *NeuroImage*, 180:68–77, 2018. doi:10.1016/j.neuroimage.2017.06.061.

A. Zwanenburg, M. Vallières, M. A. Abdalah, et al. The Image Biomarker Standardization Initiative: Standardized quantitative radiomics for high-throughput image-based phenotyping. *Radiology*, 295(2):328–338, 2020. doi:10.1148/radiol.2020191145.