

# Non-Injective Projections and the Structural Failure of Bell Factorization

Jérôme Beau 

Independent Researcher; jerome.beau@cosmochrony.org

\* Correspondence: jerome.beau@cosmochrony.org

## Abstract

Violations of Bell inequalities are commonly interpreted as evidence for nonlocal influences or as constraints on realist descriptions. We show that the failure of Bell-type factorizability arises generically whenever observable outcomes are obtained through a non-injective mapping from an underlying configuration space. In this framework, probabilistic factorization fails not because of superluminal dynamics or hidden causal influences, but as a structural consequence of information loss under projection. Observable outcomes correspond to equivalence classes of underlying configurations, preventing the assignment of independent local variables. Such non-injective mappings arise naturally when effective observables are defined as projections of higher-dimensional relational configuration spaces with non-trivial fiber structure. We demonstrate explicitly, via a toy model, that this mechanism reproduces Bell–CHSH violations while preserving operational no-signalling and statistical independence of measurement settings. The framework preserves locality at the underlying level, introduces no additional hidden-variable dynamics, and does not modify quantum mechanics. It identifies a minimal structural condition for Bell violations and clarifies how classical factorizability is recovered in regimes where the effective mapping becomes approximately injective.

**Keywords:** Bell inequalities; quantum correlations; non-injective projection; probabilistic factorization; locality; quantum foundations

## 1. Introduction

The experimental violation of Bell inequalities stands as one of the most robust and conceptually challenging features of quantum theory. Repeated tests have confirmed that correlations observed between spacelike separated measurement outcomes cannot be reproduced by any local hidden-variable model satisfying statistical factorization. This empirical fact is commonly interpreted as requiring either nonlocal dynamics, the abandonment of realism, or more radical departures from classical causal intuition.

Despite these conclusions, it is important to recall that Bell inequalities do not constrain physical dynamics directly. They constrain the logical and probabilistic structure of effective descriptions, specifically the assumption that joint outcome probabilities factorize into independent local contributions conditioned on a complete specification of relevant variables. The failure of Bell inequalities therefore indicates a breakdown of effective factorization, but does not, by itself, mandate superluminal influences or causal nonlocality.

In this work, we explore a general and minimal mechanism by which Bell-type correlations may arise without invoking nonlocal dynamics, retrocausality, or hidden variables.

Received:

Revised:

Accepted:

Published:

**Copyright:** © 2026 by the authors.

Submitted to *Entropy* for possible open access publication under the terms and conditions of the [Creative Commons Attribution \(CC BY\)](https://creativecommons.org/licenses/by/4.0/) license.

The central observation is that effective physical descriptions typically arise through many-to-one mappings from underlying configurations to observable outcomes. When this mapping is non-injective, distinct underlying configurations correspond to the same observable description, leading generically to a loss of factorization at the level of observable probabilities.

We further argue that such non-injectivity is not an arbitrary feature, but the natural consequence of a finite descriptive capacity of effective observables. Observable states correspond not to individual microscopic configurations, but to equivalence classes whose multiplicity reflects a bounded resolution of the effective description. In this sense, non-injectivity can be understood as the manifestation of a finite projective capacity, which limits the amount of distinguishable information that can be encoded in observable variables.

Within this framework, joint probabilities arise from averaging over entire classes of underlying configurations. Even if the underlying dynamics is local and deterministic, the induced probability structure at the observable level does not admit a factorized representation. The violation of Bell inequalities is thus traced to a structural property of the descriptive mapping itself, rather than to any dynamical nonlocality.

The effective loss of information associated with this projection is not arbitrary. In physical realizations, it is controlled by a finite resolution scale, set in quantum theory by Planck's constant  $\hbar$ , which determines the minimal distinguishability of underlying configurations. Quantum states therefore correspond to finite-capacity equivalence classes rather than sharply defined microscopic states. The breakdown of factorization reflects this irreducible coarse-graining, not an epistemic lack of knowledge.

This perspective also provides a natural characterization of the quantum-to-classical transition. In regimes where the effective descriptive capacity is large compared to the occupied state space, the mapping becomes approximately injective. In this low-occupancy regime, observable probabilities recover an approximately factorized form and Bell violations are suppressed. Classical statistical independence thus appears as a limiting case of effective injectivity, rather than as a fundamental principle.


The scope of the present work is deliberately restricted. We do not propose a modification of quantum dynamics, introduce additional physical degrees of freedom, or advance a specific underlying dynamical model. Our analysis is structural in nature and concerns the logical conditions under which Bell-type inequalities apply to effective descriptions. By isolating non-injective projection, understood as a finite-capacity effect, as a sufficient mechanism for the failure of factorization, we aim to clarify the conceptual origin of quantum correlations while preserving both operational locality and the empirical content of quantum theory.

The non-injective projection considered here is not introduced ad hoc, but arises naturally in relational frameworks in which spacetime and observables emerge as effective descriptions from an underlying configuration space, as discussed in [1].

## 2. Non-Injective Projection and the Failure of Bell Factorization

We consider a minimal abstract setting in which observable descriptions arise from an underlying configuration space through a generally many-to-one mapping. Let  $\Omega$  denote the space of underlying configurations. This space represents a structural configuration space underlying the effective description, rather than a physical state space evolving in spacetime. Observable outcomes are defined through a projection

$$\Pi : \Omega \rightarrow \mathcal{O}, \quad (1)$$



**Figure 1.** Schematic non-injective projection from the underlying configuration space  $\Omega$  to the observable outcome space  $O_A \times O_B$ . Multiple underlying configurations correspond to the same observable pair  $(a, b)$ , defining equivalence classes. This many-to-one identification induces a generic failure of probabilistic factorization at the observable level without requiring nonlocal dynamics.

where  $\mathcal{O}$  is the space of observable states or measurement outcomes. Crucially,  $\Pi$  is assumed to be non-injective, so that multiple distinct elements of  $\Omega$  correspond to the same observable description.

Such non-injective mappings arise generically when observable descriptions have a finite descriptive capacity. In this case, the effective space  $\mathcal{O}$  cannot encode all distinctions present in  $\Omega$ , and observable states correspond to equivalence classes of underlying configurations rather than to individual elements.

Here  $\Omega$  denotes an underlying configuration space from which effective observables arise through projection. Such configuration spaces appear in various approaches where observable descriptions are obtained through coarse-graining of a more detailed state space. Examples arise in relational or spectral reconstruction approaches [1]. This configuration space should not be interpreted as a hidden physical state space or as an ontological completion of quantum mechanics, but rather as a structural space of admissible configurations underlying the effective description.

Figure 1 provides a schematic illustration of this many-to-one projection mechanism. A concrete toy model making this structure explicit is presented in Appendix B.

Let  $A$  and  $B$  denote two spacelike separated measurement settings, with corresponding observable outcomes  $a \in \mathcal{O}_A$  and  $b \in \mathcal{O}_B$ . In standard local hidden-variable models, one assumes the existence of a variable  $\lambda$  such that the joint probability distribution factorizes as

$$P(a, b \mid A, B, \lambda) = P(a \mid A, \lambda) P(b \mid B, \lambda). \quad (2)$$

Bell inequalities follow from this factorization together with statistical independence assumptions on  $\lambda$ .

In the present framework, no such factorization is generically available. Because the observable outcomes are obtained through the projection  $\Pi$ , the probability distribution accessible at the effective level is given by

$$P(a, b \mid A, B) = \sum_{\omega \in \Pi^{-1}(a, b)} \mu(\omega), \quad (3)$$

where  $\mu$  is a probability measure on  $\Omega$ , and  $\Pi^{-1}(a, b)$  denotes the set of underlying configurations compatible with the joint observable outcome  $(a, b)$ .

When  $\Pi$  is non-injective, the preimage  $\Pi^{-1}(a, b)$  does not decompose into independent subsets associated with  $A$  and  $B$ . The same underlying configuration  $\omega \in \Omega$  may therefore contribute simultaneously to the preimages of several observable outcome pairs, preventing

the introduction of response functions  $A(A, \omega)$  and  $B(B, \omega)$  whose joint statistics reproduce  $P(a, b \mid A, B)$  through a factorizable form. As a result, the joint probability  $P(a, b \mid A, B)$  does not admit a representation as a product of marginal probabilities conditioned on any set of effective variables. The failure of factorization arises from the structure of the projection itself, not from any dynamical interaction between the measurement regions.

Importantly, locality at the underlying level is preserved. The measure  $\mu(\omega)$  may be defined over configurations with strictly local causal structure, and no superluminal influence or signal exchange is required. The violation of Bell inequalities therefore reflects a mismatch between the factorizable structure assumed in Bell-type derivations and the actual structure of effective descriptions induced by non-injective mappings.

This observation identifies non-injectivity as a sufficient condition for the failure of Bell factorization. Bell inequalities constrain only those effective models in which the mapping from underlying configurations to observables is injective or admits an equivalent factorizable representation. When this condition fails, Bell-type correlations may arise without violating local causality at the underlying level.

#### *Non-injectivity as a finite-capacity effect*

The non-injective character of the projection  $\Pi : \Omega \rightarrow \mathcal{O}$  should not be interpreted as an abstract or ad hoc assumption. It arises naturally when the effective description  $\mathcal{O}$  has a finite capacity to encode distinctions present in the underlying configuration space  $\Omega$ .

In such regimes, multiple distinct configurations  $\omega \in \Omega$  become indistinguishable within the effective description, leading to an objective loss of injectivity of the projection. This loss is structural rather than epistemic: it reflects a limitation of the effective representation itself, not an incomplete specification of underlying variables.

The multiplicity of pre-images  $\Pi^{-1}(o)$  for a given observable outcome  $o \in \mathcal{O}$  naturally defines a structural entropy associated with the projection,

$$S_{\Pi} \equiv - \sum_{o \in \mathcal{O}} \nu(o) \log \nu(o), \quad \nu(o) = \mu(\Pi^{-1}(o)),$$

where  $\mu$  denotes the measure on  $\Omega$  induced by the underlying dynamics. This *projection entropy* quantifies the degree of non-injectivity of the effective description and is independent of any notion of epistemic ignorance.

From this perspective, the breakdown of probabilistic factorization can be understood as a consequence of operating in a regime where the effective capacity is not large compared to the set of accessible configurations. In such finite-capacity regimes, distinct underlying configurations necessarily overlap at the level of observables, and factorization fails.

Conversely, when the effective capacity is large compared to the occupied portion of configuration space, the mapping becomes approximately injective and standard factorizable descriptions are recovered.

Violations of Bell-type inequalities thus correspond to the minimal regime in which finite-capacity effects become operationally observable. They can be interpreted as the weakest observable signature of a non-zero projection entropy  $S_{\Pi}$ , without invoking non-local dynamics, retrocausal influences, or hidden-variable completions.

### **3. Bell Inequalities Revisited: Structural vs Dynamical Assumptions**

Bell's theorem is commonly presented as a no-go result for local realist theories. More precisely, it establishes that no model satisfying a specific set of probabilistic assumptions can reproduce all quantum correlations. It is therefore essential to distinguish carefully between the assumptions entering the derivation of Bell inequalities and the physical principles they are often taken to represent.

The derivation of Bell inequalities relies on three key assumptions. First, outcome probabilities are assumed to be well-defined and conditioned on a complete specification of relevant variables. Second, joint probabilities are assumed to factorize for spacelike separated measurements, reflecting the absence of direct causal influence between distant measurement settings. Third, the conditioning variables are assumed to be statistically independent of the measurement choices. Violations of Bell inequalities imply that at least one of these assumptions fails.

In many discussions, the failure of Bell inequalities is interpreted as evidence for a breakdown of locality [2]. However, locality in Bell's sense is a condition on the factorization of effective probabilities, not a direct statement about the causal structure of the underlying theory. Bell inequalities constrain models in which observable outcomes can be represented as functions of local settings and a shared set of variables that fully determine those outcomes. They do not directly constrain the existence or absence of superluminal interactions.

Within the framework considered here, the failure of Bell inequalities originates from the breakdown of effective factorization. As shown in the previous section, non-injective mappings between underlying configurations and observable outcomes generically arise when the effective description has a finite capacity to encode underlying distinctions. In such regimes, no complete set of outcome-defining variables exists at the observable level. Although the underlying description may be local and deterministic, the observable description does not admit a factorized probabilistic representation.

Such non-injective projections need not be regarded as purely abstract constructions. They may arise when the effective observable space is obtained by projecting a higher-dimensional configuration space onto a lower-dimensional description. Geometric realizations, for instance based on fibered structures such as Hopf-type projections, provide concrete examples in which loss of injectivity is controlled by structural invariants associated with the projection.

This distinction clarifies the sense in which Bell's theorem applies. Bell inequalities are valid only for effective descriptions in which observable outcomes can be parameterized by a set of variables that renders the projection from underlying configurations injective, or at least approximately injective, and therefore factorizable. When this condition is not met, the assumptions required for the derivation of Bell inequalities are violated independently of any consideration of causal nonlocality.

From this perspective, violations of Bell inequalities do not compel the introduction of nonlocal dynamics or retrocausal mechanisms. They instead signal a structural limitation of effective descriptions in which observable outcomes are treated as independently specifiable local properties. Nonlocal correlations arise because joint outcomes must be globally consistent with a single non-factorizable effective description, not because information propagates between distant regions.

The present analysis therefore reframes Bell's theorem as a constraint on a class of descriptive models rather than as a direct statement about the ontology or dynamics of physical reality. Bell inequalities test the compatibility of observed correlations with injective and factorizable effective descriptions. When non-injective projections are involved, especially in finite-capacity regimes, Bell-type correlations may arise while preserving locality at the underlying level.

#### 4. Measurement, Decoherence, and the Emergence of Classicality

The structural origin of Bell-type correlations identified above naturally raises the question of their apparent absence in classical macroscopic phenomena. Everyday physical systems exhibit effectively local and statistically independent behavior, despite being

ultimately composed of quantum constituents. Any satisfactory account of Bell violations must therefore explain not only their existence, but also their suppression in the classical limit.

Within the present framework, this transition is understood as a change in the effective properties of the projection from underlying configurations to observable descriptions. While the projection is generally non-injective at microscopic scales, the effective descriptive capacity associated with observable states becomes large compared to the set of configurations that are actually explored. In this regime, the mapping becomes approximately injective.

Decoherence plays an important but secondary role in this process. Through continuous interaction with uncontrolled environmental degrees of freedom, the system is dynamically driven toward states that are stable under coarse-grained observation. This evolution effectively reduces the overlap between distinct underlying configurations contributing to the same observable outcome, thereby pushing the system toward a regime in which the projection behaves as approximately one-to-one.

From the perspective of observable descriptions, this corresponds to a regime in which the effective capacity of the description exceeds the occupied portion of configuration space. Global consistency constraints associated with non-injective projections become negligible, and observable outcomes can be treated as effectively independent.

In this regime, joint probabilities admit an approximate decomposition of the form

$$P(a, b \mid A, B) \approx P(a \mid A) P(b \mid B), \quad (4)$$

up to corrections that are exponentially small in the strength of environmental coupling or the degree of coarse-graining. Bell inequalities are therefore not violated in practice, not because the underlying structure has changed, but because the effective description operates in an approximately injective regime.

Importantly, this recovery of classical behavior does not require the introduction of additional collapse postulates or modifications of quantum dynamics. It follows from the same structural mechanism responsible for Bell violations, applied in a regime where the effective descriptive capacity is large compared to the occupied state space. Classicality thus corresponds to a limit in which observable descriptions admit an approximately complete specification, rendering joint probabilities effectively factorizable.

From this viewpoint, the quantum-to-classical transition reflects a change in descriptive accessibility rather than a change in fundamental physical laws. Bell-type correlations are suppressed when finite-capacity effects become negligible in the relevant regime. The classical world emerges as a limit in which effective injectivity is restored, and with it, the validity of classical notions of separability and locality.

## 5. Discussion and Conceptual Positioning

The analysis presented in this work proposes a structural reinterpretation of Bell inequality violations that avoids introducing nonlocal dynamics, hidden variables, or modifications of quantum mechanics. It is therefore important to clarify how this approach relates to existing interpretations and foundational proposals addressing quantum nonlocality.

Unlike local hidden-variable models, the present framework does not assume the existence of a set of accessible variables that fully determine observable outcomes [3]. No completion of the quantum state is postulated, and no additional degrees of freedom are introduced. The failure of Bell inequalities arises not from incomplete knowledge of underlying variables, but from the non-injective nature of the mapping between underlying configurations and observable descriptions.



The present approach is also distinct from superdeterministic models [4]. While superdeterminism denies statistical independence between measurement settings and underlying variables, no such assumption is required here. Measurement choices remain free and uncorrelated with the underlying configuration space. Bell violations arise solely from the structural properties of effective descriptions, not from conspiratorial correlations between settings and states.

Retrocausal interpretations similarly invoke influences propagating backward in time to account for nonlocal correlations. By contrast, the mechanism discussed here is entirely atemporal at the descriptive level. No causal influence, forward or backward, is required between distant measurement events. Correlations emerge from the global consistency constraints imposed by non-factorizable effective descriptions, rather than from dynamical signaling.

The present framework is compatible with standard accounts of quantum contextuality, but shifts the emphasis from contextual measurement dependence to descriptive non-injectivity. Rather than attributing Bell violations to context-dependent properties of observables, we identify a more general structural origin: the impossibility of assigning independent local outcomes within a single effective description when the underlying mapping is many-to-one.

Finally, we stress that the present work does not advocate a specific ontological interpretation of quantum mechanics. It neither endorses nor rejects realism, operationalism, or informational approaches. Our aim is more modest: to isolate a minimal and sufficient structural condition under which Bell-type inequalities fail. Non-injective projection, understood as a finite-capacity effect, provides such a condition, clarifying how quantum correlations can arise without abandoning locality at the underlying level.

In physical realizations, this non-injectivity can be understood as a consequence of finite descriptive capacity. In particular, Planck's constant  $h$  sets a fundamental bound on the distinguishability of underlying configurations, thereby controlling the effective size of the equivalence classes  $[\omega]$  that define observable outcomes.

In this sense, Bell inequality violations may be viewed not as evidence for exotic dynamics, but as indicators of intrinsic limitations of finite-capacity effective descriptions [5]. They reveal the boundaries of applicability of classical probabilistic reasoning, rather than the presence of nonlocal causal mechanisms.

## 6. Conclusion

### 6.1. Relation to local hidden-variable models

Violations of Bell inequalities are commonly interpreted as ruling out local hidden-variable models. More precisely, Bell's theorem constrains a class of effective descriptions in which measurement outcomes are assumed to be determined by a set of underlying variables that renders joint probabilities factorizable for spacelike separated measurements.

The analysis presented here clarifies that Bell inequalities do not directly constrain underlying physical dynamics. They constrain the logical structure of effective probabilistic descriptions. When observable outcomes admit an injective or effectively factorizable representation in terms of underlying variables, Bell-type inequalities apply. When this condition fails, Bell-type correlations may arise even if the underlying description is local, deterministic, and causally well behaved.

Within the framework considered here, the failure of Bell inequalities originates from the non-injective nature of the mapping between underlying configurations and observable outcomes. Such non-injectivity arises generically when the effective description has a finite capacity to encode distinctions present at the underlying level. Because multiple underlying configurations correspond to the same effective description, no set of outcome-

determining variables exists at the observable level that would allow joint probabilities to be decomposed into independent local contributions.

Bell inequalities are therefore violated not because of nonlocal dynamics, but because the assumptions required for probabilistic factorization are not satisfied in finite-capacity effective descriptions.

### 6.2. *Non-injective projection versus superdeterminism and free choice*

A frequent objection to any violation of statistical independence in Bell-type analyses is that it implicitly relies on superdeterministic assumptions, namely correlations between measurement settings and underlying physical states. In such scenarios, the apparent violation of Bell inequalities is attributed to a conspiratorial coordination between experimental choices and microscopic conditions.

The mechanism discussed here is fundamentally different. No correlation is postulated between measurement settings and underlying configurations, and no restriction is imposed on the freedom of experimental choices at the level of observable descriptions. Measurement settings are assumed to be freely and independently chosen within the effective description, in full accordance with standard experimental practice.

The loss of statistical independence arises instead from the non-injective structure of the mapping from underlying configurations to observable outcomes. Such non-injectivity arises generically in regimes where the effective description has a finite capacity to encode distinctions present at the underlying level. Because distinct underlying configurations are identified at the effective level, conditioning on observable variables does not isolate independent subsystems. The resulting failure of probabilistic factorization is therefore structural and geometric in origin, rather than causal or conspiratorial.

In this sense, the present framework preserves operational free choice while violating the assumptions required for Bell-type factorizations. The breakdown of statistical independence reflects intrinsic limitations of finite-capacity effective descriptions, not a fine-tuned coordination of initial conditions.

### 6.3. *Measurement, contextuality, and effective descriptions*

Recent work in quantum foundations has emphasized that nonlocal and contextual correlations may be understood as obstructions to constructing globally consistent probabilistic models from local measurement contexts [6]. Within ontological-model frameworks, observable statistics are likewise interpreted as coarse-grained descriptions of an underlying ontic state space [7]. These approaches identify contextuality as a fundamental feature of quantum theory, reflecting the impossibility of assigning outcome values independently of the measurement context.

The mechanism discussed here is complementary in spirit, but identifies a more primitive structural origin for the failure of Bell factorizability. Rather than attributing contextuality to context-dependent properties of observables, we trace it to the non-injective mapping between underlying configurations and observable descriptions. When the effective description has a finite capacity to encode distinctions present at the underlying level, distinct configurations are necessarily identified, and no globally consistent assignment of independent local outcomes is possible.

From this perspective, contextuality appears as a manifestation of descriptive non-injectivity. The impossibility of constructing non-contextual models reflects the fact that observable variables do not provide a complete specification of the underlying configuration. Bell-type correlations and contextuality therefore share a common structural origin in the finite-capacity nature of effective descriptions.



In regimes where the effective descriptive capacity is large compared to the explored configuration space, the mapping becomes approximately injective. In this limit, contextuality effects are suppressed, and observable outcomes may be treated as effectively independent. Classical statistical behavior thus emerges as a regime of approximate injectivity, rather than as a fundamentally distinct domain.

The present work does not propose a modification of quantum mechanics or a new ontological framework. Its contribution is to isolate a minimal and sufficient structural condition for the failure of Bell inequalities and the emergence of contextual correlations. Non-injective projection, understood as a finite-capacity effect, provides such a condition while preserving both operational locality and empirical adequacy.

We conclude that Bell inequality violations and contextuality need not be interpreted as evidence for fundamental nonlocality, but may instead reflect intrinsic limitations of finite-capacity effective descriptions in which observable outcomes are treated as independently specifiable local properties.

#### 6.4. Conclusion

In conclusion, we have shown that violations of Bell inequalities do not require dynamical nonlocality, nor the abandonment of realism, but can instead be understood as a generic consequence of non-injective projection in finite-capacity effective descriptions. Within this framework, Bell-type correlations arise not from superluminal influences or hidden causal mechanisms, but from structural features of observable descriptions in which multiple underlying configurations correspond to the same outcomes.

This analysis is purely structural and does not rely on any modification of quantum dynamics or on the introduction of additional hidden variables. The failure of Bell-type factorizability is traced to the descriptive level itself, and reflects the impossibility of assigning independent local outcomes within a single effective representation when the underlying mapping is non-injective.

This perspective is naturally embedded in relational approaches in which spacetime and observables are not taken as fundamental, but emerge as effective descriptions from an underlying configuration space. In particular, the non-injective projection considered here arises generically in relational and spectral frameworks where spacetime geometry itself is reconstructed from more primitive relational structures [1]. Concrete realizations may involve configuration spaces with non-trivial fiber structure, such as Hopf-type projections from  $S^3$  onto effective observable spaces, where the loss of injectivity is controlled by geometric invariants of the projection.

Quantum correlations then cease to appear as instances of mysterious “action at a distance” and instead reflect global consistency constraints imposed by the non-injective mapping between underlying configurations and the domain of localized measurements.

From this viewpoint, the empirical success of relativistic locality and quantum statistics is preserved at the effective level, while their apparent tension is resolved by recognizing the intrinsic limitations of finite-capacity effective descriptions. Classical locality and probabilistic independence emerge as limiting cases in which the effective mapping becomes approximately injective, corresponding to regimes where the available descriptive capacity exceeds the occupied configuration space. Bell-type correlations may therefore be interpreted as the minimal observable signature of a non-zero projection entropy  $S_{\Pi}$  associated with the non-injective character of the effective description.

## Appendices

### Appendix A Formal Conditions for the Failure of Probabilistic Factorization

This appendix provides a formal clarification of the structural conditions under which probabilistic factorization fails in the presence of non-injective projections. The purpose is not to introduce new physical assumptions, but to make explicit the logical content underlying the arguments presented in the main text.

Let  $(\Omega, \Sigma, \mu)$  be a probability space describing underlying configurations  $\omega \in \Omega$ , equipped with a normalized measure  $\mu$ . Observable outcomes are obtained through a measurable mapping

$$\Pi : \Omega \rightarrow \mathcal{O}, \quad (\text{A5})$$

where  $\mathcal{O}$  denotes the space of observable descriptions. We assume that  $\Pi$  is non-injective, so that there exist distinct  $\omega_1 \neq \omega_2$  such that  $\Pi(\omega_1) = \Pi(\omega_2)$ . Such non-injectivity arises generically when the effective description  $\mathcal{O}$  has a finite capacity to encode distinctions present in  $\Omega$ .

Let  $A$  and  $B$  denote two spacelike separated measurement settings, and let  $a \in \mathcal{O}_A$  and  $b \in \mathcal{O}_B$  denote the corresponding observable outcomes. The joint probability distribution accessible at the effective level is given by

$$P(a, b \mid A, B) = \mu\left(\Pi^{-1}(a, b)\right) = \sum_{\omega \in \Pi^{-1}(a, b)} \mu(\omega). \quad (\text{A6})$$

Suppose that a factorizable representation of the joint distribution exists. Then there must exist a measurable variable  $\lambda$  and a probability density  $\rho(\lambda)$  such that

$$P(a, b \mid A, B) = \int d\lambda \rho(\lambda) P(a \mid A, \lambda) P(b \mid B, \lambda). \quad (\text{A7})$$

This representation implicitly assumes that  $\lambda$  provides a complete and non-redundant parameterization of the relevant underlying configurations for the observable outcomes.

However, when  $\Pi$  is non-injective, no such variable  $\lambda$  can generically exist at the effective level. Any candidate  $\lambda$  constructed from observable data must identify entire equivalence classes

$$[\omega] = \{\omega' \in \Omega \mid \Pi(\omega') = \Pi(\omega)\}, \quad (\text{A8})$$

rather than individual underlying configurations. As a result, conditioning on  $\lambda$  does not isolate independent contributions associated with  $A$  and  $B$ .

More formally, if  $\lambda$  is a measurable function of  $\Pi(\omega)$ , then  $\lambda$  is constant over each equivalence class  $[\omega]$ . The joint probability conditioned on  $\lambda$  therefore takes the form

$$P(a, b \mid A, B, \lambda) = \sum_{\omega \in [\omega_\lambda]} \mu(\omega \mid \lambda), \quad (\text{A9})$$

which does not decompose into a product of independent marginal terms unless additional constraints are imposed that effectively restore injectivity. In generic finite-capacity regimes, such constraints are absent, and factorization fails.

This establishes that non-injectivity of the mapping from underlying configurations to observable descriptions is sufficient to obstruct any representation of the form Eq. (A7). The failure of Bell-type factorization thus follows from a purely structural property of the descriptive mapping, independently of any assumptions about dynamical nonlocality, retrocausality, or hidden variables.

The result emphasizes that Bell inequalities apply only to effective descriptions admitting an injective or equivalently factorizable representation. When observable outcomes arise through non-injective projections, violations of Bell inequalities are a natural and generic consequence of finite-capacity effective descriptions.

## Appendix B A Toy Model Illustrating Non-Injective Projection

We present a simple toy model intended to illustrate how Bell-type correlations may arise from non-injective projection without invoking nonlocal dynamics, hidden variables, or superdeterministic assumptions. The model is deliberately minimal and does not aim to reproduce the full structure of quantum theory. Its purpose is solely to make explicit the logical mechanism underlying the failure of probabilistic factorization.

In particular, the model illustrates how such non-injectivity arises when the effective description has a finite capacity to encode distinctions present at the underlying level, and how this leads generically to non-factorizable observable statistics.

### Appendix B.1 Underlying configuration space

Let  $\Omega$  denote a finite set of underlying configurations. Each configuration  $\omega \in \Omega$  represents a complete specification of a relational structure. No spatiotemporal interpretation is assumed at this level, and no notion of subsystem decomposition is imposed.

We consider a situation in which two measurement regions, denoted  $A$  and  $B$ , are identified only at the level of observable descriptions. At the level of  $\Omega$ , the configurations need not decompose into independent components associated with  $A$  and  $B$ , reflecting the fact that the effective description may have a finite capacity to encode such distinctions.

### Appendix B.2 Non-injective projection

Observable outcomes are defined through a projection

$$\Pi : \Omega \rightarrow O_A \times O_B,$$

where  $O_A$  and  $O_B$  denote the sets of possible outcomes for measurements performed in regions  $A$  and  $B$ , respectively.

The projection  $\Pi$  is assumed to be non-injective. Distinct underlying configurations may therefore correspond to the same pair of observable outcomes  $(a, b)$ . Such non-injectivity arises naturally when the effective description  $O_A \times O_B$  has a finite capacity to encode distinctions present in  $\Omega$ .

Observable states are thus identified with equivalence classes

$$[\omega] = \{\omega' \in \Omega \mid \Pi(\omega') = \Pi(\omega)\}.$$

### Appendix B.3 Emergence of correlations

Let  $\mu(\omega)$  be a probability distribution defined on  $\Omega$ . The joint probability distribution accessible at the observable level is given by

$$P(a, b) = \sum_{\omega \in \Pi^{-1}(a, b)} \mu(\omega).$$

Because the preimage  $\Pi^{-1}(a, b)$  does not generically decompose into independent subsets associated with  $A$  and  $B$ , the joint probability  $P(a, b)$  does not admit a factorized representation of the form  $P(a)P(b)$  or  $P(a \mid \lambda)P(b \mid \lambda)$  for any effective variable  $\lambda$ .

This non-factorizability arises because conditioning on observable outcomes does not isolate independent subsets of underlying configurations. Distinct configurations

contributing to a given outcome pair  $(a, b)$  may also contribute to other outcome pairs, preventing a decomposition into independent response functions associated with  $A$  and  $B$ .

The resulting correlations therefore arise not from any interaction or information exchange between  $A$  and  $B$ , but from the structure of the non-injective mapping itself. In particular, they reflect the fact that the effective description has a finite capacity to encode distinctions present in  $\Omega$ , so that multiple configurations are necessarily identified at the observable level.

From the perspective of the effective description, the two outcomes appear correlated even when the corresponding measurements are spacelike separated.

#### *Appendix B.4 Free choice and absence of superdeterminism*

Measurement settings at  $A$  and  $B$  are assumed to be freely and independently chosen within the effective description. Formally, this is expressed by the statistical independence condition

$$\mu(\omega \mid A, B) = \mu(\omega),$$

for all  $\omega \in \Omega$  and all choices of settings  $(A, B)$ .

No correlation between measurement settings and underlying configurations is required or assumed. In particular, the underlying probability measure  $\mu$  is defined independently of any experimental choices.

The violation of statistical independence at the observable level arises solely from the non-injective nature of the projection  $\Pi$ . Because observable outcomes do not provide access to the full underlying configuration, conditioning on observable variables does not isolate independent degrees of freedom. The effective description therefore fails to admit a factorized representation, even though measurement settings remain statistically independent of the underlying state.

The resulting breakdown of probabilistic factorization is thus structural rather than conspiratorial. It reflects the loss of injectivity of the descriptive mapping, not any fine-tuned correlation between experimental choices and underlying configurations.

#### *Appendix B.5 Explicit CHSH violation from non-injective projection*

We make the Bell–CHSH violation explicit by exhibiting an observable probability model generated by a non-injective projected description.

Let measurement settings be  $x \in \{0, 1\}$  at  $A$  and  $y \in \{0, 1\}$  at  $B$ , and outcomes be  $a, b \in \{-1, +1\}$ . Consider the no-signalling correlations defined by

$$P(a, b \mid x, y) = \begin{cases} \frac{1}{2} & \text{if } ab = (-1)^{xy}, \\ 0 & \text{otherwise,} \end{cases} \quad (\text{A10})$$

for which the marginals are uniform:  $P(a \mid x) = P(b \mid y) = 1/2$ .

The correlators are therefore

$$E_{xy} = \sum_{a, b = \pm 1} ab P(a, b \mid x, y) = (-1)^{xy}, \quad (\text{A11})$$

and the CHSH expression is

$$S = |E_{00} + E_{01} + E_{10} - E_{11}| = 4 > 2. \quad (\text{A12})$$

*Realization as a projected description.* Define an underlying finite configuration space  $\Omega \equiv \{(a_0, a_1, b_0, b_1) \mid a_x, b_y \in \{-1, +1\}\}$  equipped with a normalized measure  $\mu$  supported only on the subset satisfying the global constraint

$$a_0 b_0 = a_0 b_1 = a_1 b_0 = +1, \quad a_1 b_1 = -1. \quad (\text{A13})$$

This constraint encodes a non-factorizable relational structure at the level of  $\Omega$ , consistent with the absence of subsystem decomposition assumed in Appendix B.1–B.3.

Observable pairs are obtained through a projection

$$\Pi : \Omega \times \{0, 1\}^2 \rightarrow O_A \times O_B, \quad \Pi(\omega, x, y) = (a_x, b_y). \quad (\text{A14})$$

This formulation makes explicit that the underlying configuration  $\omega$  is independent of the measurement settings, which merely select which components are read out. The dependence on  $(x, y)$  therefore does not introduce contextual hidden variables, but reflects the operational definition of observables.

For fixed  $(x, y)$ , the induced mapping  $\Pi_{x,y} : \omega \mapsto (a_x, b_y)$  is generally non-injective. Many distinct underlying configurations correspond to the same observable pair  $(a, b)$ , so that the observable probabilities take the projected form

$$P(a, b \mid x, y) = \mu(\Pi_{x,y}^{-1}(a, b)). \quad (\text{A15})$$

For a symmetric choice of  $\mu$  on the constrained subset Eq. (A13), this reproduces exactly the distribution Eq. (A10).

This construction shows explicitly that a maximal CHSH violation can arise from a non-injective projection applied to an underlying configuration space with global constraints, without requiring any superluminal influence. The violation reflects the incompatibility between Bell-type factorizability and projected descriptions in which observable outcomes are defined only up to equivalence classes of underlying configurations.

## Appendix B.6 Interpretation

This toy model illustrates how correlations between observables may arise when they correspond to different projections of a single underlying configuration. In this setting, joint outcomes are not independently specifiable, but must satisfy global consistency constraints inherited from the underlying structure.

Bell-type correlations can therefore be understood as a manifestation of non-factorizability at the level of effective descriptions, rather than as evidence for nonlocal causal influences. The apparent nonlocality reflects the fact that observable outcomes are defined only up to equivalence classes of underlying configurations, and cannot be assigned independently within a single effective representation.

The role of the toy model is not to reproduce quantum theory in detail, but to isolate a minimal structural mechanism sufficient to generate Bell–CHSH violations. In particular, the use of PR-box correlations highlights that the strength of the violation is not tied to any specific dynamics, but to the failure of injectivity in the mapping from underlying configurations to observables.

More realistic realizations may impose additional constraints that restrict the set of admissible correlations (for instance to the quantum Tsirelson bound), but the structural origin of the correlations remains the same: a non-injective projection from an underlying configuration space that does not admit a subsystem factorization.

**Acknowledgments:** The author acknowledges the use of large language models as a supportive tool for refining language, structure, and internal consistency during the development of this manuscript.

All conceptual contributions, theoretical choices, and interpretations remain the sole responsibility of the author.

**Conflicts of Interest:** The author declares no conflicts of interest.

## References

1. Beau, J. Relational Reconstruction of Spacetime Geometry from Graph Laplacians. *Zenodo* **2026**. Companion paper, <https://doi.org/10.5281/zenodo.18356037>.
2. Bell, J.S. On the Einstein Podolsky Rosen paradox. *Physics Physique Fizika* **1964**, *1*, 195.
3. 't Hooft, G. *The Cellular Automaton Interpretation of Quantum Mechanics*; Vol. 185, *Fundamental Theories of Physics*, Springer, 2016. <https://doi.org/10.1007/978-3-319-41285-6>.
4. Hall, M.J.W. Local deterministic model of singlet state correlations based on relaxing measurement independence. *Physical Review Letters* **2010**, *105*, 250404. <https://doi.org/10.1103/PhysRevLett.105.250404>.
5. Spekkens, R.W. Evidence for the epistemic view of quantum states. *Physical Review A* **2007**, *75*, 032110. <https://doi.org/10.1103/PhysRevA.75.032110>.
6. Abramsky, S.; Brandenburger, A. The Sheaf-Theoretic Structure of Non-Locality and Contextuality. *New Journal of Physics* **2011**, *13*, 113036. <https://doi.org/10.1088/1367-2630/13/11/113036>.
7. Spekkens, R.W. Evidence for the Epistemic View of Quantum States: A Toy Theory. *Physical Review A* **2007**, *75*, 032110. <https://doi.org/10.1103/PhysRevA.75.032110>.