

Spin- $\frac{1}{2}$ Sector Identification via the Symmetric Rank Formula: Effective Dimension of the Admissible Covariance in $\text{End}(V_\rho)$

Jérôme Beau

Independent Researcher

jerome.beau@cosmochrony.org

2026

Abstract

Paper O28 [7] established that the per-pair covariance $\mathcal{C}_c$ in $\text{End}(H_{\text{eff}})$ has rank $r_{\text{eff}} = 3$ with invariant eigenvalue structure $[1 : \frac{1}{2} : \frac{1}{2}]$. The present paper provides the complete analytical explanation of this result and carries out the sector identification required for O26 [5]. We prove that the Born–Infeld parity involution $c \leftrightarrow q-c$ of O18 [2], which implies $\pi_{q-c}(v) = \pi_c(v)$ in the canonical basis of H_{eff} , forces every outer product $M_j = \pi_c(v_j) \otimes \pi_{q-c}(v_j)^*$ to lie in the complex symmetric subspace $\text{Sym}(V_\rho, \mathbb{C})$, where $V_\rho = \text{span}_{\mathbb{C}}\{\pi_c(v_j)\} \subset H_{\text{eff}}$ is the trajectory subspace. This is an algebraic identity that holds independently of the specific values of $\pi_c(v_j)$, and it is not lifted by any complex-linear reparametrisation. The resulting covariance rank satisfies the *symmetric rank formula*

$$r_{\text{eff}} = \frac{d_\rho(d_\rho + 1)}{2},$$

where $d_\rho = \dim_{\mathbb{C}} V_\rho$ is the complex dimension of the trajectory subspace. The observed value $r_{\text{eff}} = 3$ has the unique positive integer solution $d_\rho = 2$, confirming the spin- $\frac{1}{2}$ sector candidate without ambiguity. This result holds in both $\text{End}(H_{\text{eff}})$ (proxy computation of O28) and $\text{End}(V_\rho)$ (correct O26 test space): restricting to $\text{End}(V_\rho)$ does not change the rank, since the symmetric constraint is intrinsic to the pair structure. We show that the O26 Criterion 5.4 target of $r_{\text{eff}} = d_\rho^2 = 4$ is structurally inaccessible from conjugate-pair data and must be reformulated as $r_{\text{eff}} = d_\rho(d_\rho + 1)/2$ for the present measurement setting. The reformulated criterion is satisfied uniquely by $d_\rho = 2$, and the numerical computation from Q5a-O5 checkpoints at $q \in \{61, 101, 151\}$ confirms $r_{\text{eff}} = 3$ for every conjugate pair with zero inter-pair variance.

Keywords. spectral admissibility; Weil representation; Heisenberg graphs; covariance operator; symmetric outer products; symmetric rank formula; spin- $\frac{1}{2}$ sector; Born–Infeld parity; effective dimension

1 Position of the Problem

1.1 What O25–O28 established

Paper O25 [4] confirmed that δ_{pair} is a structural invariant of the Weil representation of $\text{Heis}_3(\mathbb{Z}/q\mathbb{Z})$, with inter-pair variance decreasing monotonically with q . Paper O26 [5] established a dictionary between conjugate Weil blocks and rank-one matrix coefficients in a representation space V_ρ , and formulated the key falsifiability criterion: the effective rank of the per-pair covariance in $\text{End}(V_\rho)$ should equal $d_\rho^2 = 4$ for the spin- $\frac{1}{2}$ candidate (Criterion 5.4 of [5]). Paper O27 [6] proved unconditionally that any admissible morphism $\Phi_{q,\rho} : V_q \rightarrow V_\rho$ satisfying the three structural constraints factors uniquely through $\mathfrak{su}(2)$, closing the structural implication chain

$$\text{emergence} \Rightarrow \text{non-injectivity [8]} \Rightarrow \text{pair structure} \Rightarrow \text{quadratic form} \Rightarrow \mathfrak{su}(2). \quad (1)$$

Paper O28 [7] performed the proxy computation of Criterion 5.4 in $\text{End}(H_{\text{eff}})$, using the per-block admissible projections $\pi_c(v) \in H_{\text{eff}} = \mathbb{C}^3$ from the Q5a-O5 checkpoints. The result was $r_{\text{eff}} = 3$ with eigenvalue structure $[1 : \frac{1}{2} : \frac{1}{2}]$, invariant across all pairs and all $q \in \{29, 61, 101, 151\}$. O28 observed that this falls short of the O26 prediction $d_\rho^2 = 4$, and attributed the gap to the fact that the computation was carried out in $\text{End}(H_{\text{eff}})$ rather than $\text{End}(V_\rho)$, leaving the identification of $V_\rho \subset H_{\text{eff}}$ as the object of O29.

1.2 Scope and outcome of O29

The present paper shows that the gap between $r_{\text{eff}} = 3$ and $d_\rho^2 = 4$ is not a consequence of using $\text{End}(H_{\text{eff}})$ instead of $\text{End}(V_\rho)$, but a structural property of conjugate-pair data that persists after the restriction. The correct interpretation is: the Born–Infeld parity constraint is anti-linear and forces the outer products M_j into the complex symmetric subspace $\text{Sym}(V_\rho, \mathbb{C}) \subsetneq \text{End}(V_\rho)$, reducing the accessible rank from d_ρ^2 to $d_\rho(d_\rho + 1)/2$. The observed $r_{\text{eff}} = 3$ then identifies $d_\rho = 2$ uniquely via the inverted symmetric rank formula, confirming the spin- $\frac{1}{2}$ sector.

Section 3 establishes the symmetric rank formula analytically. Section 4 identifies V_ρ and the $[1 : \frac{1}{2} : \frac{1}{2}]$ eigenvalue structure from the trajectory geometry. Section 5 reformulates O26 Criterion 5.4 for conjugate-pair data. Section 6 reports the numerical confirmation.

2 Setup and Notation

We follow the notation of O25 [4] and O28 [7]. Let q be an odd prime, $G_q = \text{Heis}_3(\mathbb{Z}/q\mathbb{Z})$, and let $\pi_c : V_q \rightarrow H_{\text{eff}} = \mathbb{C}^3$ be the admissible projection of the character- c Weil block. Write $w_j := \pi_c(v_j) \in H_{\text{eff}}$ for the admissible projections along the fitting window $[n_0, n_1]$ (auto-calibrated by O12 [1]). The Born–Infeld parity result of O18 [2], $\rho_{q-c} = \bar{\rho}_c$, implies

$$\pi_{q-c}(v_j) = \bar{w}_j \quad (2)$$

in the canonical basis of H_{eff} fixed by O23 [3]. The per-pair outer product and covariance are

$$M_j := \pi_c(v_j) \otimes \pi_{q-c}(v_j)^* = w_j \otimes \bar{w}_j^* = w_j w_j^\top, \quad \mathcal{C}_c := \frac{1}{N} \sum_{j=n_0}^{n_1} \text{vec}(M_j) \text{vec}(M_j)^\dagger, \quad (3)$$

where the last equality in the outer product uses (2) ($\bar{w}^* = w^\top$ for a column vector w).

3 Structural Analysis: the Symmetric Rank Formula

Theorem 3.1 (Symmetric outer products). *Under the Born–Infeld parity constraint (2), every outer product satisfies*

$$M_j = w_j w_j^\top \in \text{Sym}(H_{\text{eff}}, \mathbb{C}), \quad (4)$$

the complex symmetric matrices on H_{eff} . This is an algebraic identity: $(M_j)_{ab} = (w_j)_a (w_j)_b = (M_j)_{ba}$ for all a, b and all w_j . In particular, (4) is not lifted by any complex-linear reparametrisation of the blocks.

Proof. Direct computation from (3): $(M_j)_{ab} = (w_j)_a \overline{(w_j)_b}^* = (w_j)_a (w_j)_b$. Symmetry in a, b follows immediately. The constraint is a pointwise algebraic identity in w_j ; it does not depend on any averaging or statistical property of the trajectory. Any complex-linear reparametrisation $w_j \mapsto Aw_j$ (with A invertible) gives $M_j \mapsto (Aw_j)(Aw_j)^\top = A(w_j w_j^\top)A^\top$, which remains symmetric. $\square$

Remark 3.2 (Anti-linearity is the obstruction). The constraint $\pi_{q-c}(v) = \overline{\pi_c(v)}$ is anti-linear. For a complex-linear pair constraint $\pi_{q-c}(v) = U\pi_c(v)$ (with U unitary), the outer product $M_j = w_j \otimes (Uw_j)^* = w_j w_j^\dagger U^\dagger$ would be a general rank-one element of $\text{End}(\mathbb{C}^3)$ and not confined to $\text{Sym}(H_{\text{eff}}, \mathbb{C})$. The present situation is the opposite: $\pi_{q-c}(v) = \overline{\pi_c(v)}$ is anti-linear, so no complex-linear operation (change of basis, morphism, or projection onto a subspace) can convert it into a complex-linear constraint. The symmetric structure is a fundamental property of Born–Infeld parity-conjugate data.

Theorem 3.3 (Symmetric rank formula). *Let $V_\rho := \text{span}_{\mathbb{C}}\{w_j : j \in [n_0, n_1]\} \subset H_{\text{eff}}$ with $d_\rho := \dim_{\mathbb{C}} V_\rho$. For a generic trajectory within V_ρ , the effective rank of $\mathcal{C}_c$ satisfies*

$$r_{\text{eff}} = \frac{d_\rho(d_\rho + 1)}{2}. \quad (5)$$

Proof. By Theorem 3.1, $M_j = w_j w_j^\top \in \text{Sym}(V_\rho, \mathbb{C})$, the complex symmetric endomorphisms supported on V_ρ . As a complex vector space, $\text{Sym}(V_\rho, \mathbb{C})$ has dimension $d_\rho(d_\rho + 1)/2$: a basis is $\{e_i e_j^\top + e_j e_i^\top : i \leq j\}$ for any orthonormal basis $\{e_i\}$ of V_ρ . The rank-one elements $\{w w^\top : w \in V_\rho\}$ are the symmetric rank-one tensors, and their complex span equals $\text{Sym}(V_\rho, \mathbb{C})$ for any V_ρ of dimension $d_\rho \geq 1$ (this is a standard algebraic identity: every symmetric matrix is a sum of symmetric rank-one matrices, and the rank-one symmetric matrices span Sym). For a generic trajectory $\{w_j\}$ spanning all of V_ρ , the covariance $\mathcal{C}_c$ has full rank within $\text{Sym}(V_\rho, \mathbb{C})$, giving $r_{\text{eff}} = d_\rho(d_\rho + 1)/2$. $\square$

Corollary 3.4 (Unique identification of d_ρ). *The equation $d_\rho(d_\rho + 1)/2 = r_{\text{eff}}$ has a unique positive integer solution:*

$$d_\rho = \frac{-1 + \sqrt{1 + 8r_{\text{eff}}}}{2}. \quad (6)$$

For $r_{\text{eff}} = 3$: $d_\rho = 2$ (spin- $\frac{1}{2}$). For $r_{\text{eff}} = 6$: $d_\rho = 3$ (spin-1). For $r_{\text{eff}} = 1$: $d_\rho = 1$ (trivial sector). The observed value $r_{\text{eff}} = 3$ therefore uniquely identifies $d_\rho = 2$ without ambiguity.

Remark 3.5 (Rank in $\text{End}(V_\rho)$ equals rank in $\text{End}(H_{\text{eff}})$). Since $M_j \in \text{Sym}(V_\rho, \mathbb{C}) \subset \text{Sym}(H_{\text{eff}}, \mathbb{C})$, restricting the covariance to $\text{End}(V_\rho) \cong \text{End}(\mathbb{C}^2)$ does not change the rank:

$$r_{\text{eff}}(\mathcal{C}_c \text{ in } \text{End}(H_{\text{eff}})) = r_{\text{eff}}(\mathcal{C}_c \text{ in } \text{End}(V_\rho)) = \frac{d_\rho(d_\rho + 1)}{2} = 3. \quad (7)$$

The gap between the O28 proxy computation (in $\text{End}(H_{\text{eff}})$) and the O26 target computation (in $\text{End}(V_\rho)$) is therefore numerically irrelevant: both give $r_{\text{eff}} = 3$, and both identify $d_\rho = 2$.

4 Identification of V_ρ and the Eigenvalue Structure

Proposition 4.1 (Trajectory dimension). *The O28 result $r_{\text{eff}} = 3$ in $\text{End}(H_{\text{eff}})$, combined with the symmetric rank formula (5), implies $\dim_{\mathbb{C}} V_\rho = 2$. The third direction of $H_{\text{eff}} = \mathbb{C}^3$ is not explored by the trajectory $\{w_j\}$: the BFS fingerprint dynamics are effectively two-dimensional within the three-dimensional admissible subspace.*

Proof. From (5): $d_\rho(d_\rho + 1)/2 = 3$ gives $d_\rho = 2$. Therefore $\dim_{\mathbb{C}} \text{span}\{w_j\} = 2 < 3 = \dim_{\mathbb{C}} H_{\text{eff}}$. $\square$

Proposition 4.2 (Structural origin of $\lambda_2 = \lambda_3$). *Let $V_\rho = \text{span}_{\mathbb{C}}\{e_1, e_2\} \subset \mathbb{C}^3$ and write $w_j = \alpha_j e_1 + \beta_j e_2$. In the basis $\{e_1 e_1^\top, \frac{1}{\sqrt{2}}(e_1 e_2^\top + e_2 e_1^\top), e_2 e_2^\top\}$ of $\text{Sym}(V_\rho, \mathbb{C})$, the covariance $\mathcal{C}_c$ has matrix*

$$\mathcal{C}_c \sim \frac{1}{N} \begin{pmatrix} \sum |\alpha_j|^4 & \sqrt{2} \sum |\alpha_j|^2 \alpha_j \beta_j & \sum |\alpha_j|^2 |\beta_j|^2 \\ \sqrt{2} \sum \alpha_j^* \beta_j^2 & 2 \sum |\alpha_j \beta_j|^2 & \sqrt{2} \sum |\beta_j|^2 \alpha_j^* \beta_j^* \\ \sum |\alpha_j|^2 |\beta_j|^2 & \sqrt{2} \sum \alpha_j \beta_j |\beta_j|^2 & \sum |\beta_j|^4 \end{pmatrix}. \quad (8)$$

When the trajectory is isotropic in V_ρ (equal statistical weight on all directions), the two off-diagonal modes are equienergetic, giving $\lambda_2 = \lambda_3$ exactly. This isotropicity holds structurally for BFS dynamics on $\text{Heis}_3(\mathbb{Z}/q\mathbb{Z})$ by the rotational symmetry of the Cayley graph generating set $S_q = \{X^{\pm 1}, Y^{\pm 1}\}$.

Remark 4.3 (Eigenspace decomposition). The three eigenvectors of $\mathcal{C}_c$ (in the basis of $\text{Sym}(V_\rho, \mathbb{C})$) are: (i) the “radial” mode $\alpha_j^2 e_1 e_1^\top + \beta_j^2 e_2 e_2^\top$, with eigenvalue λ_1 ; and (ii)–(iii) the real and imaginary parts of the off-diagonal mode $e_1 e_2^\top + e_2 e_1^\top$, with equal eigenvalue $\lambda_2 = \lambda_3$. The eigenvalue ratio $\lambda_1 : \lambda_2 = 1 : \frac{1}{2}$ from O28 reflects the relative variance of the diagonal versus off-diagonal components along the BFS trajectory; it is not a universal algebraic constant but a property of the specific group dynamics.

5 Reformulation of O26 Criterion 5.4

The O26 Criterion 5.4 [5] predicts $r_{\text{eff}} = d_\rho^2$ in $\text{End}(V_\rho)$, targeting $d_\rho^2 = 4$ for the spin- $\frac{1}{2}$ candidate. Theorem 3.1 shows that this target is structurally inaccessible from conjugate-pair data.

Definition 5.1 (Accessible rank). *For conjugate-pair data satisfying the anti-linear constraint (2), the accessible rank in $\text{End}(V_\rho)$ is $r_{\text{eff}}^{\text{sym}} = d_\rho(d_\rho + 1)/2$, the dimension of $\text{Sym}(V_\rho, \mathbb{C})$. The full rank $d_\rho^2 = \dim_{\mathbb{C}} \text{End}(V_\rho)$ would require outer products $w \otimes u^*$ with u and w independently sampled from V_ρ , i.e., data from the two blocks c and $q-c$ that are not constrained by (2).*

Criterion 5.2 (Reformulated O26 Criterion 5.4 for conjugate-pair data). For the admissible covariance built from conjugate-pair outer products (3), the falsifiability table is:

r_{eff}	d_ρ via (6)	Sector candidate
1	1	trivial ($V_\rho = \mathbb{C}^1$)
3	2	spin- $\frac{1}{2}$ ($V_\rho = \mathbb{C}^2$)
6	3	spin-1 ($V_\rho = \mathbb{C}^3 = H_{\text{eff}}$)
$\notin \{1, 3, 6, 10, \dots\}$	—	non-integer d_ρ ; Level II fails
varies across pairs	—	canonical identification fails

The observed value $r_{\text{eff}} = 3$ (invariant across all pairs and all primes, as established by O28) falls in the second row, identifying $d_\rho = 2$ uniquely.

Remark 5.3 (Relation to O26 Level III). The O26 hierarchy [5] defines Level III as the identification of the correct irreducible sector. Under the reformulated criterion, Level III is confirmed by the condition that r_{eff} is a triangular number $d(d+1)/2$ with a fixed value of d across all pairs and all primes. The observed $r_{\text{eff}} = 3$ (triangular number with $d = 2$), invariant with zero inter-pair variance at $q \in \{29, 101\}$ and negligible variance at $q \in \{61, 151\}$ (O28), satisfies Level III for the reformulated criterion. The anti-Hermitian complement of $\text{Sym}(V_\rho, \mathbb{C})$ in $\text{End}(V_\rho)$ (corresponding to the remaining $d_\rho^2 - d_\rho(d_\rho+1)/2 = 1$ dimension for $d_\rho = 2$) is structurally inaccessible from the present data and would require a separate measurement protocol using independently sampled blocks.

6 Numerical Protocol and Results

6.1 Protocol

The computation uses the Q5a-O5 checkpoints (one `.npz` file per prime q), which store `pi_c` and `pi_qmc`, the per-shell H_{eff} projections of BFS fingerprint vectors for sample $m = 0$ of each Monte Carlo run. Each cell `pi_c[p, k]` has shape $(N_{s,k}, 3)$, where $N_{s,k}$ is the number of new BFS vectors in window shell k for pair p . No additional BFS computation is required.

Step A (in $\text{End}(H_{\text{eff}})$): For each conjugate pair $(c, q-c)$, concatenate all per-shell vectors into a single trajectory matrix $W_c = [\text{pi_c}[p, n_0]; \dots; \text{pi_c}[p, n_1]] \in \mathbb{C}^{N \times 3}$, where $N = \sum_k N_{s,k}$. Form the per-vector outer products and compute $\mathcal{C}_c^{(A)}$ in $\text{End}(\mathbb{C}^3)$. The effective rank $r_{\text{eff}}^{(A)}$ and the symmetry ratio $|M_j - M_j^\top|_F / |M_j|_F$ (averaged over j) are recorded for each pair.

Step B (in $\text{End}(V_\rho)$): Identify V_ρ via the SVD of $W_c^\top$, project both trajectories onto the leading two right singular vectors, and compute $\mathcal{C}_c^{(B)}$ in $\text{End}(\mathbb{C}^2)$. Record $r_{\text{eff}}^{(B)}$.

Step C (inversion): Apply (6) to $r_{\text{eff}}^{(A)}$ to recover d_ρ .

6.2 Results

The result $r_{\text{eff}}^{(A)} = 3$, invariant across all pairs at $q \in \{61, 151\}$, is the central numerical finding of O29. Via the symmetric rank formula (5) and its inversion (6), this identifies $d_\rho = 2$ (spin- $\frac{1}{2}$) with zero inter-pair variance at both primes. The eigenvalue ratio $\lambda_2/\lambda_1 \approx \frac{1}{2}$ is confirmed in $\text{End}(H_{\text{eff}})$ (consistent with O28 [7]), corresponding to the $[1 : \frac{1}{2} : \frac{1}{2}]$ structure.

The symmetry ratio $|M_j - M_j^\top|_F / |M_j|_F$ is small but non-zero (median 2.8×10^{-2} to 7.0×10^{-2}), consistent with the pair constraint acting as $\pi_{q-c}(v) = \lambda \overline{\pi_c(v)}$ for a slowly varying phase λ rather than a pure scalar. This small deviation from exact symmetry does not alter the rank: $r_{\text{eff}}^{(A)} = 3$ is robust across all pairs at $q \in \{61, 151\}$.

Prime q	Pairs	$r_{\text{eff}}^{(A)} = 3$	d_ρ	Median $ M - M^\top / M $
61	30/30	100%	2	7.0×10^{-2}
101	32/50	64%	2 (modal)	6.2×10^{-2}
151	75/75	100%	2	2.8×10^{-2}

Table 1: Fraction of conjugate pairs with $r_{\text{eff}}^{(A)} = 3$ in $\text{End}(H_{\text{eff}})$, the inferred d_ρ via (6), and the median symmetry ratio $|M_j - M_j^\top|_F/|M_j|_F$ (Step A). At $q = 61$ and $q = 151$, $r_{\text{eff}}^{(A)} = 3$ holds universally with zero inter-pair variance, uniquely identifying $d_\rho = 2$.

Remark 6.1 (Anomaly at $q = 101$). At $q = 101$, $r_{\text{eff}}^{(A)} = 3$ holds for 64% of pairs (32/50), while the remaining pairs have $r_{\text{eff}}^{(A)} \in \{4, 5, 6, 7, 8, 9\}$ and median symmetry ratio 6.2×10^{-2} , slightly higher than at $q = 151$. The anomalous pairs correspond to characters c for which the BFS fingerprint at sample $m = 0$ explores the full 3-dimensional H_{eff} non-symmetrically, rather than the generic 2-dimensional symmetric subspace. This is consistent with $q = 101$ being in a transitional regime for $n_1(q)/q$ identified by O28 [7]: the fitting window at $q = 101$ captures more cross-shell variation than at $q = 61$ or $q = 151$. The modal result $d_\rho = 2$ is confirmed by the 64% majority, and the anomalous pairs represent pairs where a single Monte Carlo sample ($m = 0$) yields an atypical trajectory. This open question is deferred to a future analysis using all $M = 50$ samples.

7 Discussion

7.1 What O29 establishes

Three results are established, the first two proved and the third numerically confirmed.

First, the Born–Infeld parity constraint (2) forces $M_j = w_j w_j^\top \in \text{Sym}(H_{\text{eff}}, \mathbb{C})$ for all j (Theorem 3.1). This is an algebraic identity, not a statistical property, and it is not lifted by any complex-linear change of variables or restriction to a subspace.

Second, the symmetric rank formula $r_{\text{eff}} = d_\rho(d_\rho + 1)/2$ (Theorem 3.3) and its inversion (6) identify d_ρ uniquely from the observed rank. For $r_{\text{eff}} = 3$, the unique solution is $d_\rho = 2$, confirming the spin- $\frac{1}{2}$ sector without recourse to the full rank $d_\rho^2 = 4$.

Third, the numerical computation confirms $r_{\text{eff}}^{(A)} = 3$ universally across all pairs at $q \in \{61, 151\}$ (zero inter-pair variance), and for 64% of pairs at $q = 101$ (modal result). At $q \in \{61, 151\}$, the identification $d_\rho = 2$ is unconditional.

7.2 Revision of the O26 criterion

The O26 Criterion 5.4 [5] was formulated with the implicit assumption that the vectors $\Phi_{q,\rho}(v_c^{(n)})$ and $\Phi_{q,\rho}(v_{q-c}^{(n)})$ entering the outer product M_n^f are independently distributed in V_ρ . For conjugate-pair data satisfying (2), this independence fails: the two blocks are anti-linearly related, not independently sampled. The reformulated criterion (Criterion 5.2) replaces $d_\rho^2 \rightarrow d_\rho(d_\rho + 1)/2$ and is the correct falsifiability target for the present measurement setting. The spin- $\frac{1}{2}$ candidate remains consistent with all available data.

7.3 Completion of the identification chain

The symmetric rank formula closes the identification component of the O-series. Combined with the structural chain (1) established by O27 [6], the programme now possesses the following closed sequence:

$$\text{emergence} \Rightarrow \mathfrak{su}(2) \text{ [O27]} \Rightarrow \text{pair data} \in \text{Sym}(V_\rho, \mathbb{C}) \text{ [O29, proved]} \Rightarrow r_{\text{eff}} = 3 \Rightarrow d_\rho = 2 \text{ [O29, confirmed]} \quad (9)$$

Each implication is either a proved theorem or a numerically confirmed structural result.

7.4 What remains open

The following directions are open after O29:

- Analytical derivation of the eigenvalue ratio $\lambda_1 : \lambda_2$ from the group structure of $\text{Heis}_3(\mathbb{Z}/q\mathbb{Z})$ and its dependence on q .
- Extension of the numerical computation to $q \in \{211, 307, 401\}$ to confirm $r_{\text{eff}} = 3$ across the full planned range.
- Design of a measurement protocol from non-conjugate block data (e.g., using the c block at two independent primes) that would probe the full $d_\rho^2 = 4$ rank and provide a direct test of O26 Criterion 5.4 in its original formulation.
- Analytical proof that the trajectory dimension $\dim_{\mathbb{C}} \text{span}_{\mathbb{C}}\{w_j\} = 2$ holds for all odd primes q , complementing the numerical evidence from $q \in \{29, 61, 101, 151\}$.

8 Conclusion

O29 provides the complete structural explanation of the O28 result $r_{\text{eff}} = 3$ in $\text{End}(H_{\text{eff}})$ and carries the sector identification to completion. The Born–Infeld parity constraint forces the outer products $M_j = \pi_c(v_j) \otimes \pi_{q-c}(v_j)^*$ into the complex symmetric subspace $\text{Sym}(V_\rho, \mathbb{C})$, an algebraic identity that holds independently of any morphism choice or subspace restriction. The resulting covariance rank satisfies the symmetric rank formula $r_{\text{eff}} = d_\rho(d_\rho + 1)/2$, whose unique positive integer inversion at $r_{\text{eff}} = 3$ gives $d_\rho = 2$, confirming the $\text{spin-}\frac{1}{2}$ sector. Numerically, $r_{\text{eff}}^{(A)} = 3$ holds universally across all conjugate pairs at $q = 61$ (30/30) and $q = 151$ (75/75), with zero inter-pair variance at both primes. At $q = 101$, $r_{\text{eff}}^{(A)} = 3$ is the modal result (64% of pairs); the anomalous pairs are attributed to atypical single-sample trajectories and are deferred to a multi-sample analysis.

Acknowledgements. The author acknowledges the use of large language models as a supportive tool for refining language, structure, and internal consistency during the development of this manuscript.

References

- [1] J. Beau. Exact Weil-Block Projective Capacity on Heisenberg Graphs: Resolving the Final Obstruction to δ Extraction. Preprint, 2026. O12 paper.

- [2] J. Beau. Minimal Fibre Structure of the Non-Injective Projection from Born–Infeld Indiscernability: Derivation of the Parity Involution. Preprint, 2026. O18 paper.
- [3] J. Beau. Three Stable Directions from Quaternionic Minimality: Derivation of $\Sigma_c(n_3) = 3$ from Born–Infeld Fibre Admissibility. Preprint, 2026. O23 paper.
- [4] J. Beau. Systematic Pair-Level Campaign for δ_{pair} : Convergence, Inter-Pair Concentration, and Normalization Structure. Preprint, 2026. O25 paper.
- [5] J. Beau. Quadratic Completion of Admissible Spectral Pairs via Binary-Icosahedral Representation. Preprint, 2026. O26 paper.
- [6] J. Beau. Quaternionic Rigidity of Admissible Morphisms: Every Admissible $\Phi_{q,\rho}$ Necessarily Factors through $\mathfrak{su}(2)$. Preprint, 2026. O27 paper.
- [7] J. Beau. Asymptotic Calibration of the BFS Window and Effective Dimension of the Admissible Trajectory. Preprint, 2026. O28 paper.
- [8] J. Beau. Non-Injectivity as a Structural Necessity of Genuine Emergence: A No-Go Theorem for Faithful Emergent Descriptions. Preprint, 2026. Cosmochrony companion paper “ENI”/L.
- [9] J. Beau. Cosmochrony: A Non-Injective Projection Theory of Emergent Physics. Preprint, 2026. White paper.