

The Fine-Structure Constant as a Projective Invariant: Lamb Shift and Schwinger Effect in a Pre-Geometric Framework

Jérôme Beau^{1*}

^{1*}Independent Researcher, France.

Corresponding author(s). E-mail(s): jerome.beau@cosmochrony.org;

Abstract

The Lamb shift and the Schwinger effect are among the most precise and conceptually challenging predictions of quantum electrodynamics. They are conventionally interpreted as consequences of vacuum fluctuations, radiative corrections, and non-perturbative instabilities of a dynamical quantum vacuum. While this interpretation achieves remarkable quantitative success, it leaves open fundamental questions regarding the ontological status of the vacuum and the origin of renormalization procedures.

In this work, we propose a unified reinterpretation of these phenomena within the Cosmochrony framework, in which all effective physical observables emerge from a non-injective projection of a single pre-geometric relational substrate. From this perspective, vacuum-related effects do not arise from physical excitations of an underlying field-theoretic vacuum, but from intrinsic limitations of the projection mapping relational configurations to effective spacetime descriptions.

We show that the Lamb shift can be understood as projective spectral noise, reflecting the finite resolvability of highly localized interactions and the coarse-grained influence of unresolved relational modes. Similarly, the Schwinger effect is reinterpreted as a saturation phenomenon associated with the bounded capacity of the projection to transport relational flux under extreme field conditions, leading to a breakdown of injectivity and the emergence of new effective degrees of freedom.

A central result of this analysis is a unified structural interpretation of the fine-structure constant. Rather than appearing as an unexplained fundamental parameter, it emerges as an invariant ratio between projective resolution and relational flux capacity, governing both atomic-scale spectral corrections and strong-field instabilities.

This reinterpretation preserves the quantitative predictions of quantum electrodynamics while providing a coherent ontological account of its most subtle effects.

It demonstrates that precision QED phenomena can be consistently embedded within a pre-geometric relational framework, supporting the view that quantum field theories operate as effective descriptions of deeper projective constraints rather than as fundamental theories of the vacuum.

1 Introduction and Motivation

Quantum Electrodynamics (QED) stands as one of the most accurate physical theories ever constructed, achieving extraordinary agreement between theoretical predictions and experimental measurements across a wide range of phenomena [1]. Among its most emblematic successes are the Lamb shift in hydrogenic atoms and the non-perturbative prediction of electron-positron pair production in ultra-strong electric fields, known as the Schwinger effect [2, 3].

Despite their empirical precision, these phenomena raise persistent conceptual questions regarding the ontological status of the quantum vacuum. In the standard formulation, both effects are attributed to vacuum fluctuations and radiative corrections arising from the interaction of charged particles with quantized electromagnetic fields. The vacuum is thus treated as a dynamical entity, endowed with structure, energy, and virtual excitations, whose effects must be regularized and renormalized to yield finite physical predictions [4].

Within a relational pre-geometric approach, the quantum vacuum is not regarded as a fundamental physical medium. Instead, all effective physical observables arise from a non-injective projection Π of a single pre-geometric relational substrate χ onto an effective spacetime description. As developed in previous work, the non-injectivity of this projection implies that distinct relational configurations may correspond to identical effective observables, leading to intrinsic limits in localization, resolution, and transport [5].

From this perspective, phenomena traditionally attributed to vacuum fluctuations may be reinterpreted as manifestations of projective limitations rather than dynamical excitations of an underlying field. In regimes of high informational density or extreme field intensity, the projection Π ceases to be effectively injective, and observable corrections emerge as structural effects associated with bounded resolution and saturation of relational flux.

The present work applies this projective interpretation to two cornerstone effects of QED: the Lamb shift and the Schwinger pair production mechanism. We argue that the Lamb shift can be understood as a form of projective spectral noise, arising from the finite resolvability of interactions localized at atomic scales. In this view, the energy level shifts of s -states relative to p -states do not reflect electron self-energy in a fluctuating vacuum, but the sensitivity of highly localized states to the spectral structure of the underlying relational Laplacian.

Similarly, the Schwinger effect is reinterpreted as a saturation phenomenon associated with a bounded capacity of the relational substrate to transport effective electromagnetic flux. When this capacity is exceeded, the projection becomes unstable, and the system relaxes through a local re-stratification of the substrate, manifesting

as the creation of particle-antiparticle pairs. Pair production is therefore not viewed as extraction from the vacuum, but as an inevitable consequence of non-injective projection under extreme field conditions.

A central outcome of this unified interpretation is a revised understanding of the fine-structure constant α . Rather than appearing as an unexplained fundamental parameter, α emerges as an invariant ratio between projective resolution and relational flux capacity, governing both the magnitude of atomic-level corrections and the threshold behavior of strong-field phenomena.

The goal of this paper is not to challenge the quantitative success of QED, but to provide an alternative ontological interpretation of its most subtle predictions. By reframing the Lamb shift and the Schwinger effect as projective and saturation phenomena, respectively, this work aims to demonstrate that precision quantum electrodynamics can be consistently embedded within a pre-geometric relational description. This embedding offers a coherent account of vacuum-related effects without invoking a physically populated vacuum, and supports the view that a unified relational and projective interpretation can consistently account for phenomena across disparate physical regimes.

2 Projective Origin of Vacuum Effects

In standard quantum field theory, vacuum-related effects are commonly interpreted as arising from fluctuations of quantized fields around a lowest-energy state. These fluctuations are treated as physical, albeit virtual, entities whose observable consequences emerge through radiative corrections and loop processes. While this interpretation has proven extraordinarily successful at the computational level, it implicitly attributes a rich dynamical structure to the vacuum itself.

Within a relational pre-geometric approach, no such physically populated vacuum is postulated. Instead, all effective physical phenomena originate from a single pre-geometric relational substrate χ , whose configurations are mapped to effective spacetime observables through a projection $\Pi : \Omega \rightarrow \mathcal{O}$. As established in earlier work, this projection is generically non-injective: distinct relational configurations may correspond to identical effective descriptions [5].

Non-injectivity of the projection has direct and unavoidable physical consequences. In particular, it implies that the effective description cannot resolve arbitrarily fine relational distinctions. This limitation introduces a fundamental bound on localization, timing, and interaction resolution, independent of any dynamical fluctuations. Observable deviations from idealized point-like behavior therefore arise as structural features of the projection itself.

We refer to this limitation as *projective resolution*. At sufficiently low energies and moderate field intensities, the projection Π behaves as effectively injective, and standard quantum and classical descriptions remain accurate. However, in regimes characterized by extreme localization or high relational flux, the projection saturates. Beyond this regime, additional relational information cannot be faithfully transported into the effective spacetime description.

The saturation of projective resolution manifests observationally as corrections to otherwise degenerate or symmetric configurations. Crucially, these corrections do not signal the presence of additional degrees of freedom, but rather the breakdown of perfect correspondence between relational configurations and effective observables. In this sense, so-called vacuum effects are reinterpreted as *projection artifacts*, encoding the finite capacity and resolution of the mapping from χ to spacetime.

This perspective provides a unified reinterpretation of phenomena traditionally treated as conceptually distinct. Corrections attributed to vacuum polarization, self-energy, or zero-point fluctuations are recast as manifestations of bounded projectability. They reflect the fact that effective fields and particles are not fundamental entities, but emergent descriptions constrained by the structural properties of the underlying relational substrate.

In the following sections, this projective interpretation is applied to two paradigmatic quantum electrodynamical phenomena. The Lamb shift is shown to arise from the sensitivity of highly localized atomic states to projective spectral noise. The Schwinger effect is interpreted as a saturation-induced instability, in which excess relational flux forces a local reconfiguration of the substrate. Together, these effects illustrate how precision QED phenomena can be consistently understood within a unified relational and projective description.

3 The Lamb Shift as Projective Spectral Noise

The Lamb shift constitutes one of the most precise and conceptually significant tests of quantum electrodynamics. Experimentally observed as a lifting of the degeneracy between the $2s_{1/2}$ and $2p_{1/2}$ levels in hydrogen, it played a decisive role in the development of renormalized QED [2]. In the standard interpretation, this shift is attributed to radiative corrections, notably electron self-energy and vacuum polarization effects.

From the projective perspective introduced in Section 2, the Lamb shift admits a different interpretation. Rather than signaling an interaction with fluctuating vacuum degrees of freedom, the observed energy correction reflects a limitation in the spectral resolvability of highly localized bound states under the non-injective projection $\Pi : \Omega \rightarrow \mathcal{O}$. The Lamb shift thus appears as a manifestation of projective spectral noise.

Atomic bound states correspond, in this framework, to admissible localized relaxation modes of the relational substrate. While the linear effective description predicts degeneracies between states of equal principal quantum number, this degeneracy is lifted once non-linear saturation and projectability constraints are taken into account. The lifting is not uniform across orbital configurations, but depends sensitively on the degree to which a given state probes the inner structure of the projection. This finite spectral smearing is consistent with the non-injective character of the projection previously identified in the context of Bell-type correlations, where effective interactions remain delocalized even in stationary configurations [6].

States with nonzero probability density at the origin, such as s -states ($\ell = 0$), probe regions of elevated relational density, where coupling between localized modes and the global relaxation flow is strongest. In these regions, Born–Infeld-type saturation effects and spectral frustration are maximal. By contrast, p -states ($\ell = 1$), whose

wavefunctions vanish at the nucleus, remain less sensitive to these inner-core constraints. The observed s - p splitting therefore reflects a differential sensitivity to unresolved relational structure rather than a dynamical self-interaction of the electron.

At the effective level, this mechanism induces a finite upward shift of the s -state energy. Dimensional considerations suggest a characteristic scale of the form

$$\Delta E_{\text{Lamb}} \sim \kappa \alpha^5 m_e c^2, \quad (1)$$

where α denotes the invariant ratio between projective resolution and relational flux capacity, and κ is a numerical factor of order unity encoding details of the local spectral structure. This estimate yields an energy shift of order 10^{-5} eV, consistent with the observed magnitude of the Lamb splitting. Crucially, the correction is intrinsically finite, reflecting the bounded nature of the underlying relational dynamics.

From this viewpoint, renormalization in QED can be reinterpreted as an effective procedure compensating for the coarse-grained representation of unresolved spectral structure. The quantitative success of standard radiative corrections is preserved, but their ontological interpretation is revised. The Lamb shift does not arise from vacuum fluctuations, but from spectral frustration induced by the finite resolution and saturation properties of the projection.

This spectral-probe logic extends naturally to other fine and hyperfine corrections, which similarly depend on the proximity of localized electronic states to regions of maximal projective compression. Precision spectroscopy thus provides a direct observational window into the spectral limits of emergent spacetime dynamics.

4 The Schwinger Effect as Projective Flux Saturation

The Schwinger effect provides a paradigmatic example of a non-perturbative phenomenon in quantum electrodynamics. It predicts the spontaneous creation of electron-positron pairs in the presence of a sufficiently strong electric field, with a characteristic exponential suppression below a critical field strength [3]. Unlike radiative corrections such as the Lamb shift, this effect cannot be captured by any finite-order perturbative expansion.

In the standard interpretation, Schwinger pair production is described as a vacuum instability, often visualized as a tunneling process through the energy barrier separating negative- and positive-energy states. While this picture successfully reproduces the observed rate, it relies on the notion of a dynamically populated vacuum subject to extreme excitation.

Within a relational pre-geometric description, the Schwinger effect admits a structural reinterpretation. Effective electromagnetic fields correspond to directed transport of relational relaxation flux through the non-injective projection $\Pi : \Omega \rightarrow \mathcal{O}$. This transport is subject to a finite capacity, reflecting the bounded ability of the projection to sustain smooth relaxation under increasing field intensity, consistent with the saturation mechanisms identified in a previous Born–Infeld-type effective analysis [7].

At moderate field strengths, relaxation proceeds homogeneously and the projection remains effectively injective. As the imposed electric field increases, the associated relaxation flux approaches a saturation threshold. Beyond this threshold, smooth

transport becomes inadmissible: additional flux cannot be conveyed without loss of injectivity, and the homogeneous relaxation regime becomes unstable.

The onset of Schwinger pair production corresponds precisely to this saturation point. Rather than extracting particles from a pre-existing vacuum, the system restores admissibility by activating additional effective modes of the projection. These modes manifest as particle-antiparticle pairs, which redistribute excess relaxation flux into stable, projectable structures. Global neutrality is preserved, ensuring charge conjugation symmetry.

From this perspective, pair production acts as a dissipation mechanism by structure creation. When the transport capacity of the effective field is exceeded, the relational substrate reorganizes locally, enlarging the space of admissible configurations. The exponential form of the Schwinger rate reflects the probabilistic breakdown of effective injectivity under extreme flux conditions, rather than tunneling through an energy barrier of the vacuum.

This interpretation naturally aligns with Born–Infeld-type effective dynamics, in which field invariants are bounded and divergences are avoided through saturation. Here, such behavior is not imposed at the level of an effective Lagrangian, but emerges from the finite transport capacity of the relational substrate itself.

The Schwinger effect thus exemplifies a general principle: when effective descriptions are driven beyond the limits of projectability, stability is restored not by divergence, but by reconfiguration. Matter creation appears as a universal relaxation channel, ensuring the continued consistency of the effective spacetime description under extreme conditions.

5 Unified Interpretation via the Fine-Structure Constant

The fine-structure constant α occupies a singular position in modern physics. It governs the strength of electromagnetic interactions, controls the magnitude of radiative corrections in atomic systems, and sets the scale of non-perturbative phenomena such as the Schwinger effect. Despite its ubiquity and empirical precision, its origin remains unexplained within standard quantum electrodynamics, where it enters as a fundamental dimensionless input parameter.

Within a relational pre-geometric description, α acquires a unified geometric and relational interpretation. Rather than characterizing a coupling between elementary fields, it emerges as an invariant ratio between two structural properties of the projection $\Pi : \Omega \rightarrow \mathcal{O}$: the resolution with which relational configurations can be mapped to effective observables, and the maximum flux of relational tension that can be transported without loss of injectivity.

These two properties manifest differently depending on the physical regime. In atomic systems, where localization and spectral resolution are the dominant constraints, α controls the sensitivity of energy levels to unresolved relational structure. As shown in Section 3, the Lamb shift reflects the response of highly localized states to projective spectral noise. The magnitude of this response is set by the ratio between the

characteristic atomic scale and the fundamental projective resolution, encapsulated by α .

In contrast, in strong-field regimes such as those associated with the Schwinger effect, the dominant limitation is not spectral resolution but transport capacity. As discussed in Section 4, α governs the threshold at which relational flux saturates, forcing a breakdown of injectivity and the emergence of new effective degrees of freedom. The same invariant ratio that determines the scale of atomic corrections thus fixes the critical field strength for pair production.

This dual role of α explains why phenomena as disparate as the Lamb shift and Schwinger pair creation are controlled by the same numerical constant within standard QED. From the projective standpoint, this is not coincidental. Both effects probe complementary limits of the same underlying structure: the bounded capacity of the projection to faithfully encode relational information into effective spacetime observables.

Importantly, this interpretation does not require a modification of the numerical value of α or a departure from the quantitative predictions of QED. Instead, it provides an ontological explanation for the universality of α across perturbative and non-perturbative regimes. Renormalized coupling constants and effective field strengths appear as different manifestations of a single structural invariant.

The emergence of α as a projective invariant also clarifies its apparent scale-independence. While effective couplings may run with energy due to the structure of the effective description, the underlying ratio between projective resolution and flux capacity remains fixed. This explains why α consistently governs both low-energy precision spectroscopy and extreme-field phenomena without invoking additional fundamental parameters.

In this unified view, the fine-structure constant is neither arbitrary nor mysterious. It encodes a balance condition between distinguishability and transport at the level of the relational substrate. The fact that this balance permeates the full range of electromagnetic phenomena supports the view that quantum electrodynamics operates as an effective description of deeper projective constraints rather than as a fundamental theory of the vacuum.

6 Experimental Constraints and Observational Outlook

Any alternative interpretation of precision quantum electrodynamics must confront the stringent experimental constraints imposed by modern measurements. In this respect, the projective description developed in this work is not exempt from empirical scrutiny. On the contrary, the reinterpretation of vacuum-related effects as manifestations of bounded projectability leads to concrete and falsifiable implications.

Precision spectroscopy of hydrogenic and hydrogen-like systems provides the most direct constraints on projective spectral noise. Measurements of the Lamb shift, hyperfine splittings, and transition frequencies in light atoms agree with QED predictions at the level of parts per billion. Within a relational pre-geometric description, this agreement implies that the effective projective resolution scale lies well below currently

accessible atomic length scales. Any additional contribution arising from unresolved relational structure must therefore remain subdominant within the precision of existing data.

However, the projective interpretation suggests that deviations from standard QED predictions, if present, should exhibit characteristic patterns. Rather than appearing as uniform shifts, such deviations would preferentially affect states with enhanced localization, higher nuclear charge, or increased sensitivity to short-distance structure. Precision comparisons between transitions with different orbital character, as well as systematic studies across isoelectronic sequences, offer potential avenues to isolate such effects.

Strong-field phenomena provide a complementary testing ground. While direct observation of Schwinger pair production in static electric fields remains experimentally challenging, rapidly advancing laser technologies are approaching regimes where non-perturbative effects become accessible. In this context, the present description predicts no deviation in the existence or order of magnitude of the critical field, but allows for controlled modifications in the functional form of the pair production rate near saturation.

In particular, departures from the standard exponential dependence may arise as the relational flux approaches its transport bound. Such deviations would manifest as changes in the pre-exponential factors or as environment-dependent corrections in inhomogeneous or time-dependent fields. These signatures differ qualitatively from those expected from conventional higher-order QED corrections and may therefore serve as discriminants between dynamical vacuum and projective saturation interpretations.

An important aspect of the present description is that it does not introduce additional free parameters at the effective level. The same invariant controlling atomic-scale corrections governs strong-field instabilities. As a result, constraints derived from precision spectroscopy and strong-field experiments are not independent, but jointly restrict the admissible structure of the projection. This interdependence enhances the falsifiability of the description.

Beyond laboratory experiments, the projective interpretation may also be relevant in astrophysical environments characterized by extreme electromagnetic fields, such as magnetars or relativistic plasma configurations. While such settings introduce additional modeling uncertainties, they offer access to regimes unattainable on Earth and may provide indirect constraints on saturation mechanisms.

Overall, the present relational and projective description remains fully compatible with existing experimental data, while making distinctive qualitative predictions regarding the structure of corrections in extreme regimes. Future advances in precision spectroscopy, strong-field laser physics, and high-intensity plasma experiments will therefore play a crucial role in assessing whether vacuum-related phenomena are more naturally interpreted as dynamical field effects or as manifestations of bounded projectability in an emergent spacetime description.

Beyond isolated laboratory tests, the projective interpretation naturally suggests a hierarchy of falsification across physical scales. Precision atomic spectroscopy probes the resolution limit of the projection through spectral corrections such as the Lamb shift. Strong-field experiments test the transport capacity of relational flux via the Schwinger

threshold. If the underlying saturation mechanism is universal, analogous signatures are expected to arise in more extreme astrophysical and cosmological environments, where effective descriptions are likewise driven toward their projective limits. This multi-scale structure enhances the falsifiability of the description by linking phenomena traditionally treated in isolation through a common set of structural constraints.

7 Discussion and Conceptual Implications

The reinterpretation of the Lamb shift and the Schwinger effect developed in this work has implications that extend beyond the specific phenomena considered. Together, these effects probe complementary limits of effective quantum electrodynamics. The Lamb shift explores the regime of extreme localization and spectral resolution, while the Schwinger effect probes the limits of flux transport under intense fields. Their unification within a single projective description suggests that they are not independent curiosities, but structural manifestations of the same underlying constraints.

A central conceptual outcome of this analysis concerns the status of the quantum vacuum. In the standard formulation of QED, the vacuum is treated as a physically active entity, endowed with fluctuations, virtual excitations, and renormalized energies. While this picture is operationally successful, it leaves open deep questions about the ontological nature of these entities and the origin of the procedures required to render predictions finite.

Within a relational pre-geometric description, vacuum-related effects acquire a different status. They do not arise from the dynamics of an underlying field-theoretic vacuum, but from the structural limitations of the projection mapping a pre-geometric relational substrate to effective spacetime observables. Renormalization, in this view, is not a correction to an ill-defined physical medium, but an effective bookkeeping device compensating for the finite resolution and transport capacity of the emergent description.

This shift in perspective clarifies why quantum electrodynamics exhibits both extreme precision and persistent conceptual tension. The theory operates remarkably well within its domain of effective applicability, yet its interpretational difficulties signal the presence of deeper constraints not expressible within a purely spacetime-based formalism. A relational and projective description provides a setting in which these constraints can be made explicit, without undermining the empirical content of QED.

An important implication of this work is that the apparent diversity of quantum corrections conceals a high degree of structural unity. Spectral shifts, non-perturbative instabilities, and coupling strengths are shown to be controlled by a small number of invariant projective properties. The fine-structure constant emerges as a central organizing parameter, encoding the balance between distinguishability and transport at the relational level. Its universality across disparate regimes ceases to be mysterious once this balance is recognized as fundamental.

The results presented here reinforce the viability of a unified relational and projective interpretation of precision quantum electrodynamics. Related analyses have addressed gravitation, cosmology, and quantum non-locality within comparable principles. By embedding precision QED phenomena within the same class of structural constraints,

the present work extends this interpretive strategy to the core of atomic and particle physics. This extension is essential for assessing whether such descriptions can provide a coherent effective account across a broad range of physical regimes.

Several open questions remain. A more explicit characterization of the relational Laplacian spectrum and its coarse-grained impact on atomic dynamics would allow for quantitative refinements of the present interpretation. Similarly, a detailed modeling of flux saturation in time-dependent or inhomogeneous fields could sharpen predictions for strong-field experiments. These developments are left for future work.

In conclusion, the Lamb shift and the Schwinger effect need not be regarded as evidence for a physically populated vacuum. They can instead be understood as signatures of bounded projectability in an emergent spacetime description. This reinterpretation preserves the empirical success of quantum electrodynamics while providing a coherent ontological account of its most subtle phenomena. As such, it supports the view that precision quantum field theories are effective manifestations of deeper relational structures rather than fundamental descriptions of physical reality.

Funding

The author declares that no external funding was received for this research.

Acknowledgements. The author acknowledges the use of large language models as a supportive tool for refining language, structure, and internal consistency during the development of this manuscript. All conceptual contributions, theoretical choices, and interpretations remain the sole responsibility of the author.

References

- [1] Weinberg, S.: The Quantum Theory of Fields. Volume I: Foundations. Cambridge University Press, Cambridge (1995)
- [2] Lamb, W.E., Retherford, R.C.: Fine structure of the hydrogen atom by a microwave method. *Physical Review* **72**(3), 241–243 (1947) <https://doi.org/10.1103/PhysRev.72.241>
- [3] Schwinger, J.: On gauge invariance and vacuum polarization. *Physical Review* **82**(5), 664–679 (1951) <https://doi.org/10.1103/PhysRev.82.664>
- [4] Peskin, M.E., Schroeder, D.V.: An Introduction to Quantum Field Theory. Addison-Wesley, Reading, MA (1995)
- [5] Beau, J.: Emergent spacetime geometry from bounded relational relaxation. Zenodo (2026) <https://doi.org/10.5281/zenodo.18407505>
- [6] Beau, J.: Bell inequality violations from non-injective projection. Zenodo (2026) <https://doi.org/10.5281/zenodo.18371173>

- [7] Beau, J.: Emergent spacetime geometry from bounded relational relaxation. Zenodo (2026) <https://doi.org/10.5281/zenodo.18407505>