

The Spectral Admissibility Sub-Programme

Presentation Note 1

Jérôme Beau

Independent Researcher

`jerome.beau@cosmochrony.org`

2026

Abstract

The spectral admissibility sub-programme of the Cosmochrony corpus addresses a single central question: which spectral sectors of the Weil representation of $\text{Heis}_3(\mathbb{Z}/q\mathbb{Z})$ survive the Born–Infeld bounded-flux constraint? The answer organises the entire O-series (papers O1–O32 and five precursor papers) into a logically closed derivation chain:

bounded flux $\implies$ admissibility envelope $\implies$ Weil-sector capacity $\implies \delta_{\text{pair}} \implies \beta^* \approx 0.126$.

This note maps the constituent papers, identifies four internal phases, records the status of every result as proved, structural, or open, and states the two remaining open deliverables: the pointwise analytical proof of hypothesis [H-color] (partially resolved at three levels in O32) and the extension of the numerical campaign to $q = 401$.

Keywords. spectral admissibility; Born–Infeld constraint; Weil representation; Heisenberg group; pair observable; capacity exponent; lepton mass hierarchy; $\text{SU}(3)$; colour triplet; non-injective projection; presentation note.

Contents

1	Motivation and Central Question	3
2	Position in the Cosmochrony Programme	3
3	Logical Chain of the Sub-Programme	3
4	Constituent Papers	4
4.1	Precursor papers: SU(2) admissibility sub-programme	4
4.2	O-series core: LPS phase (O1–O8)	4
4.3	O-series core: Heisenberg transition (O9–O15)	5
4.4	Pair observable and transfer chain (O16–O24)	5
4.5	Numerical campaign and closure papers (O25–O32)	6
5	Inputs from Upstream Results	6
6	Outputs to Downstream Branches	7
7	Status of Results	8
7.1	Proved results	8
7.2	Structural results	8
7.3	Numerical evidence	9
7.4	Open hypotheses	9
8	Open Deliverables	9
9	Conclusion	10

1 Motivation and Central Question

The spectral admissibility sub-programme answers the following question.

Central question. The non-injective projection Π of the Cosmochrony framework acts on the Weil representation of $\text{Heis}_3(\mathbb{Z}/q\mathbb{Z})$, decomposed into irreducible blocks V_c indexed by characters $c \in (\mathbb{Z}/q\mathbb{Z})^\times$. The Born–Infeld saturation constraint bounds the projective flux carried by each mode: $A_n \leq A_n^{\max} := c_\chi / \sqrt{\lambda_n}$, where λ_n is the Laplacian eigenvalue at BFS depth n and c_χ is the BI saturation constant. *Which combinations of Weil sectors remain admissible under this constraint, and what capacity exponent δ_{pair} do they carry?*

The practical importance of this question is that δ_{pair} governs the cascade exponent β^* via the structural relation $\beta^* \approx 1/(\delta_{\text{pair}} + \frac{1}{2})$ established in the O-series. The value $\beta^* \approx 0.126$ matches the phenomenological window $\beta^* \in (0.09, 0.13)$ derived from the charged-lepton mass ratios $m_e : m_\mu : m_\tau$, making this sub-programme the primary quantitative connection between the projective framework and Standard Model observables.

The sub-programme is also the hardest to navigate: it spans over thirty papers written over a period of successive obstruction identifications and resolutions. This note is intended as a structured entry point, not a summary of results.

2 Position in the Cosmochrony Programme

The Cosmochrony corpus is organised into three branches. Branch I establishes the axiomatic primitive: four axioms (A1–A4) suffice to derive $\text{Heis}_3(\mathbb{Z}/q\mathbb{Z})$ and its Weil representation as theorems [4, 15, 14]. Branch II is the spectral admissibility sub-programme (this note). Branch III derives quantum mechanics, spacetime geometry, gauge structure, and fermionic matter from Branch I axioms and Branch II spectral data.

The spectral admissibility sub-programme therefore occupies a pivotal position: it is the *computational engine* of the corpus. It takes the algebraic output of Branch I (the Weil representation and its decomposition into irreducible blocks) and produces the two inputs that Branch III depends on:

1. the identification of the admissible sector as the spin- $\frac{1}{2}$ representation of $\text{SU}(2)$ embedded via the admissibility thread $Q_8 \subset 2I \subset \text{SU}(2)$;
2. the numerical value $\beta^* \approx 0.126$ and its representation-theoretic interpretation as the scaling exponent of Hilbert–Schmidt norm growth along trajectories in the minimal admissible non-abelian sector $\mathfrak{su}(2)$.

Remark 1 (Vocabulary note). Throughout this note, the term *admissibility constraint* replaces the earlier “relaxation” vocabulary of the preliminary Cosmochrony formulation. The substrate χ is static: no substrate dynamics or relaxation mechanism is postulated at the foundational level. Observable evolution is an induced ordering of admissible projections, not a dynamics of χ itself.

3 Logical Chain of the Sub-Programme

The sub-programme is organised around the following derivation chain.

$$\underbrace{c_\chi}_{\text{BI saturation}} \implies \underbrace{A_n^{\max} = c_\chi / \sqrt{\lambda_n}}_{\text{admissibility envelope}} \implies \underbrace{\sigma_{\text{pair}}^{\text{can}}(n)}_{\text{pair capacity}} \implies \underbrace{\delta_{\text{pair}}}_{\text{capacity exponent}} \implies \underbrace{\beta^* \approx 0.126}_{\text{cascade exponent}} . \tag{1}$$

Each arrow in (1) corresponds to a distinct analytical or numerical sub-problem, resolved at a different stage of the sub-programme. The chain is *unconditional* with respect to the fibre structure of Π (O24, [36]), meaning that no free parameter is adjusted to produce β^* . The sole remaining conditionality concerns the extension of the $SU(3)$ sector: this requires hypothesis [H-color] (Section 8).

Three structural facts underlie the chain and should not be confused.

1. The *parity involution* $c \leftrightarrow q - c$, derived from Born–Infeld evenness in O18 [12], is the structural origin of the pair observable $\sigma_{\text{pair}}^{\text{can}}(n) = \sigma_c(n) \cdot \sigma_{q-c}(n)$. The factor-of-two relation $\delta_{\text{pair}} = 2\delta_c$ is a consequence of this fibre pairing, not a normalisation artefact.
2. The *observable hierarchy*: the pair capacity $\sigma_c(n)$ (normalised BFS span in the Weil block V_c) is strictly finer than the Markov spectrum of the Weil operators. Spectral equality of characters does not imply capacity profile equality. This distinction is essential for hypothesis [H-color].
3. The *admissibility thread* $Q_8 \subset 2I \subset SU(2)$ is an output of the sub-programme, not an input. Q_8 is the minimal non-abelian finite subgroup of $SU(2)$ compatible with bounded-flux admissibility; $2I$ is the maximal admissible finite subgroup; the LPS family provides the Ramanujan approximation to $SU(2)$ in the large- q limit.

4 Constituent Papers

The sub-programme comprises five precursor papers and thirty-two O-series papers. Table 1 provides an overview organised by internal phase.

4.1 Precursor papers: $SU(2)$ admissibility sub-programme

Four papers precede the O-series proper and explore spectral admissibility on Cayley graphs of finite $SU(2)$ subgroups.

SpectralAdmissibility [29] carries out the spectral admissibility programme on the chain $Q_8 \subset 2I \subset \text{PSL}(2, \mathbb{F}_q)$. It derives the admissibility hierarchy $A_n^{\text{max}} = c_\chi / \sqrt{\lambda_n}$ in the linearised regime and shows the non-linear DBI correction preserves it. For the $SU(3)$ chain $\Delta(27) \subset \Sigma(168)$, it establishes the co-admissibility of the fundamental and anti-fundamental representations via character theory.

SpectralCapacity [30] introduces the capacity functional $\mathcal{C}_\alpha(G, S)$ and proves the binary-polyhedral maximality conjecture: $2I$ uniquely maximises $\mathcal{C}_\alpha(G, S)$ for all $\alpha > 0$ among finite $SU(2)$ subgroups with neutral generating sets. The proof is closed analytically for degrees $d \in \{6, 12, 24\}$ (Theorem 8.1 of that paper).

SpectralGramRigidity [18] shows that the geometry of neutral generating sets on $SU(2)$ is fully encoded in pairwise character values $G_{st} = -\frac{1}{2}\chi_\rho(st)$.

ThreeGenerations [34] derives the three-generation structure from the ADE stratigraphic levels of binary Cayley graphs (structural).

SpectralRelaxation [6] establishes $\Lambda_{\text{proj}} \asymp \lambda_2$ for LPS expander families, providing the entry point for the O-series dynamics.

4.2 O-series core: LPS phase (O1–O8)

The opening O-series papers work on Lubotzky–Phillips–Sarnak expander graphs, which have exponential shell growth $|B_n| \sim p^n$ and therefore represent the most natural large-group setting for bounded-flux dynamics.

O1 [24] shows that LPS saturation is a fixed-valence artefact; growing valence restores the pre-saturation window.

O3 [22] derives the phenomenological target window $\beta^* \in (0.09, 0.13)$ from the charged-lepton mass ratios $m_e : m_\mu : m_\tau$ (structural).

O4 [32] proves the structural upper bound $\beta \leq 1$ from the BI flux constraint and the Cheeger isoperimetric inequality.

O5 [3] establishes the obstruction theorem: the β gap cannot be closed by any purely representation-theoretic or fixed-representation approach.

O6 [11] proves matrix-level confinement of the cascade exponent.

O7 [20] reformulates the target as a capacity exponent $\delta \in [7.4, 10.6]$ on three-step path fingerprints (structural).

O8 [1] establishes the geometric obstruction closing the LPS phase: exponential shell growth compresses the entire pre-saturation window into $O(\log q)$ BFS steps, preventing stable slope estimation.

4.3 O-series core: Heisenberg transition (O9–O15)

The geometric obstruction of O8 forces a change of arena. The Cayley graph $\text{Cay}(\text{Heis}_3(\mathbb{Z}/q\mathbb{Z}), S_q)$ has polynomial ball growth $|B_n| \sim Cn^4$ (Bass–Guivarc’h theorem, homogeneous dimension $D = 4$), restoring an exploitable window of $\Theta(q)$ BFS steps.

O9 [21] resolves the geometric obstruction by switching to Heisenberg graphs. Polynomial growth is confirmed; the window depth scales as $\Theta(q)$.

O10 [19] performs the first large- q computation ($q = 101$, $|G_q| = 1,030,301$) and identifies an algorithmic obstruction: the dense TensorSketch fingerprint is structurally mismatched to the problem, requiring sketch dimension $D \gg q^4$.

O11 [37] resolves the algorithmic obstruction via a bidimensional Fourier-character proxy for the Weil-block decomposition; a stable estimate $\hat{\delta}_{\text{cap}} \approx 3.4$ is extracted at $q = 307$.

O12 [8] implements the exact Weil projection, closes the representation-level ambiguity left by O11, and extracts $\hat{\delta}_{\text{exact}} \approx 4.3\text{--}4.8$ for $q \in \{29, 53, 61\}$ ($R^2 > 0.98$).

O13 [7] confirms asymptotic stability of the exact Weil-block capacity over the extended prime range and establishes monotone decrease of $\hat{\delta}_{\text{exact}}(q)$, definitively refuting the finite-size hypothesis.

O14 [17] corrects the $\delta \mapsto \beta^*$ relation via block normalisation and central-phase contributions.

O15 [16] establishes the block-level no-go: no reweighting of individual block data recovers the admissible window $\delta \in [7.4, 10.6]$ (proved, Corollary 4.8 of that paper). This result forces the introduction of the pair observable in O16.

4.4 Pair observable and transfer chain (O16–O24)

The block-level no-go of O15 is structurally informative: it identifies the wrong observable class, not a defect of the transfer mechanism. The physically correct observable is the *pair capacity* $\sigma_{\text{pair}}^{\text{can}}(n) = \sigma_c(n) \cdot \sigma_{q-c}(n)$ on conjugate pairs $\{c, q - c\}$.

O16 [9] identifies the observable mismatch. The exponent doubling $\delta_{\text{pair}} = 2\delta_c \approx 7.44$ resolves the gap (structural).

O17 [10] proves that $\rho_{q-c} = \bar{\rho}_c$ is forced by fibre-level admissibility; the pair structure is derived, not postulated.

O18 [12] gives the foundational derivation of the parity involution $c \leftrightarrow q - c$. Born–Infeld parity is the only global symmetry of the BI action, ensuring each fibre of Π is exactly $\{\chi, -\chi\}$ (minimal fibre condition, proved, Theorem 4.1).

O19 [2] proves that the residual amplitude factor $r(c, q) \in \{1, 2, 3, 4\}$ does not affect δ_{pair} ; the observable is pipeline-independent.

O20 [23] establishes a dynamical selection criterion for admissible modes via projective persistence (structural).

O21 [28] defines the observable saturation rank n_3^{obs} via the threshold condition $\Sigma_c(n_3) = 3$, constructs the fibre-level transfer $\delta_{\text{pair}} \rightarrow \beta^*$, and resolves the O15 obstruction by identifying it as an observable-class mismatch.

O22 [27] proves that Born–Infeld projection locking forces n_3^{obs} onto a BFS shell of $\text{Cay}(G_q, S_q)$: shell-alignment is a theorem, not an independent geometric hypothesis.

O23 [35] closes the complementary question: the threshold value $\Sigma_c(n_3) = 3$ is not an arithmetic input but a structural consequence of Born–Infeld fibre admissibility. The value three is the unique dimension of the imaginary subspace $\text{Im } \mathbb{H}$ of the minimal admissible associative non-commutative algebra.

O24 [36] is the capstone of the analytical O-series. The Verticality Lemma (“ $\ker \Pi$ is free, $\text{Im } \Pi$ is constrained”) closes the unconditional transfer chain $c_\chi \rightarrow \delta_{\text{pair}} \rightarrow \beta^*$ (Corollary 4.2: $\beta^* \approx 0.126$, proved).

4.5 Numerical campaign and closure papers (O25–O32)

With the analytical chain closed by O24, the post-O24 programme turns to three tasks: numerical validation of δ_{pair} , representation-theoretic identification of the admissible sector, and the $\text{SU}(3)$ colour sector.

O25 [33] carries out the systematic pair-level numerical campaign across all $(q-1)/2$ conjugate pairs with $M = 50$ independent block samples per pair, for $q \in \{29, 61, 101, 151, 211\}$. Results: $\bar{\delta}_{\text{pair}} \in \{8.89, 9.98, 9.55, 9.13, 8.78\}$ with inter-pair standard deviation decreasing from 0.54 to 0.11. After the O14 normalisation correction, values $\delta_{\text{corr}}(q) \in [7.7, 9.0]$ fall consistently within the admissible window $[7.4, 10.6]$ for $q \in \{61, 101, 151, 211\}$.

O26 [25] establishes the quadratic completion of admissible spectral pairs via binary-icosahedral representation, providing the Level III identification criterion for the $\text{spin-}\frac{1}{2}$ sector.

O27 [26] proves that every admissible morphism $\Phi_{q,\rho}$ necessarily factors through $\mathfrak{su}(2)$ (quaternionic rigidity theorem, proved unconditionally). The identification chain becomes:

$$\text{emergence} \Rightarrow \mathfrak{su}(2) \text{ [O27]} \Rightarrow \text{pair data} \in \text{Sym}(V_\rho, \mathbb{C}) \text{ [O29]} \Rightarrow r_{\text{eff}} = 3 \Rightarrow d_\rho = 2 \text{ [O29]}.$$

O28 [5] confirms asymptotic calibration of the BFS window: $\delta_{\text{corr}}(q) \in [7.4, 10.6]$ for all $q \in \{29, 61, 101, 151, 211\}$; the covariance C_c in $\text{End}(H_{\text{eff}})$ has rank exactly 3 for every conjugate pair at $q \in \{61, 101, 151\}$, with invariant eigenvalue structure $[1 : \frac{1}{2} : \frac{1}{2}]$.

O29 [31] identifies the $\text{spin-}\frac{1}{2}$ irreducible sector $V_\rho \subset H_{\text{eff}} = \mathbb{C}^3$ via the symmetric rank formula $r_{\text{eff}} = d_\rho(d_\rho + 1)/2$. The unique solution $d_\rho = 2$ is confirmed numerically for $q \in \{61, 101, 151\}$.

O30 [38] derives analytically the invariant eigenvalue ratio $\lambda_1/\lambda_2 = 2$ of the per-pair covariance: Born–Infeld parity anti-linearity forces an equatorial constraint on the admissible fibre, and the factor of two is purely structural (squared symmetric-tensor normalisation of the Carnot-degree-2 central generator $Z = [X, Y]$, proved).

O31 [40] develops the structural framework for $\text{SU}(3)$ emergence. Metaplectic intertwining of $\text{Heis}_3(\mathbb{Z}/q\mathbb{Z})$ forces triplet co-admissibility unconditionally for the colour-adapted Cayley graph $\text{Cay}(G_q, S_q^{(3)})$. $\text{SU}(3)$ is identified as the symmetry group of the co-admissible colour triplet, conditional on hypothesis [H-color] for the standard generating set. O31 also closes the spectral approach to [H-color]: exact generator-twist isospectrality does not imply colour-triplet spectral degeneracy.

O32 [39] provides the numerical test of [H-color] on the standard graph. Results for $q \in \{61, 151, 211\}$ (test, $q \equiv 1 \pmod{3}$) and $q \in \{29, 101\}$ (control, $q \equiv 2 \pmod{3}$): hypothesis [H-color] passes all three test primes; $q = 307$ confirms ($R_{\text{var}} = 0.7$, all 4 triplets).

5 Inputs from Upstream Results

The spectral admissibility sub-programme takes the following inputs from Branch I.

Table 1: Summary of the O-series by internal phase. Status codes: P = proved; S = structural; N = numerical; O = open.

Phase	Papers	Central output	Status
Precursors	SpAdm, SpCap, SpGram, 3Gen	$Q_8 \subset 2I$, binary maximality	P/S
LPS phase	SpRel, O1–O8	$\beta \leq 1$; geometric obstruction	P
Heisenberg transition	O9–O15	Exact $\hat{\delta}_{\text{exact}} \approx 4.5$; no-go	P
Pair + transfer	O16–O24	$c_\chi \rightarrow \delta_{\text{pair}} \rightarrow \beta^*$, unconditional	P
Numerical + sector	O25–O30	$\delta_{\text{corr}} \in [7.4, 10.6]$; $d_\rho = 2$	P/N
SU(3) / colour	O31–O32	SU(3) conditional on [H-color]	S/N/O

1. **Weil representation.** The fact that $\text{Heis}_3(\mathbb{Z}/q\mathbb{Z})$ and its Weil representation V_ρ are theorems of the axioms A1–A4, not additional postulates. This is established in Foundation [4] and HeisenbergStructure [14].
2. **BI saturation functional.** The Born–Infeld action with saturation constant c_χ is the unique analytic completion compatible with bounded projective transport capacity. The sub-programme takes c_χ as a parameter; its structural origin is in Branch I.
3. **Non-injectivity no-go.** The ENI paper [15] establishes that non-injectivity is a structural necessity of genuine emergence, not an interpretive option. The pair structure and the parity involution $c \leftrightarrow q - c$ are downstream consequences.
4. **Noscale principle.** The absence of an external dimensional scale at the admissibility level [13] ensures that the only dimensionful parameter available to the sub-programme is c_χ , enforcing the internal consistency of the capacity law.

No result from Branch III is an input to the sub-programme. The admissibility thread $Q_8 \subset 2I \subset \text{SU}(2)$, the value δ_{pair} , and the cascade exponent β^* are all outputs.

6 Outputs to Downstream Branches

Table 2 lists the principal outputs of the sub-programme and their downstream consumers.

Table 2: Principal outputs of the spectral admissibility sub-programme and their consumers.

Output	Consumer	Status
$\beta^* \approx 0.126$	SMchargemass, lepton hierarchy	P
Admissibility thread $Q_8 \subset 2I \subset \text{SU}(2)$	Q1–Q3 (quantum sector)	P
Spin- $\frac{1}{2}$ sector $V_\rho \cong \mathbb{C}^2$	Q7–Q11 (geometry)	P
$\Sigma_c(n_3) = 3$ and $\text{Im } \mathbb{H} \cong \mathfrak{su}(2)$	Q7 (spatial identification)	P
SU(3) co-admissible triplet (conditional)	Q6a, Gauge note	S/O
$\delta_{\text{corr}}(q) \in [7.4, 10.6]$ for $q \in \{61, \dots, 211\}$	Open: $q \rightarrow \infty$ limit	N

7 Status of Results

7.1 Proved results

The following results are established as theorems within their stated assumptions.

1. *Binary-polyhedral maximality* (SpectralCapacity [30]): $2I$ uniquely maximises $\mathcal{C}_\alpha(G, S)$ for all $\alpha > 0$ among finite $SU(2)$ subgroups with neutral generating sets. Closed analytically for $d \in \{6, 12, 24\}$ and by Proposition 8.11 for the dicyclic exclusion.
2. *Gram rigidity* (SpectralGramRigidity [18]): the geometry of neutral generating sets on $SU(2)$ is fully encoded in pairwise character values.
3. $\beta \leq 1$ *upper bound* (O4 [32]): the structural upper bound from the BI flux constraint and the Cheeger inequality.
4. *Block-level no-go* (O15 [16]): no reweighting of individual block data recovers $\delta \in [7.4, 10.6]$; the pair observable is necessary.
5. *Parity involution* (O18 [12]): BI parity is the unique global symmetry of the BI action; the minimal fibre condition $\Pi^{-1}(x) = \{\chi, -\chi\}$ is forced.
6. *Pipeline independence of δ_{pair}* (O19 [2]): the residual amplitude factor $r(c, q)$ does not affect the pair exponent.
7. *Shell-alignment theorem* (O22 [27]): Born–Infeld projection locking forces $n_3^{\text{obs}} \in \mathbb{N}$ as the first element of $L_{\text{BI}} \cap N_q$.
8. *Quaternionic minimality* (O23 [35]): the threshold $\Sigma_c(n_3) = 3$ follows from the fact that $\mathbb{H}$ is the minimal admissible associative non-commutative algebra, and $\dim_{\mathbb{R}} \text{Im } \mathbb{H} = 3$.
9. *Unconditional transfer chain* (O24 [36]): the Verticality Lemma closes $c_\chi \rightarrow \delta_{\text{pair}} \rightarrow \beta^* \approx 0.126$ without residual conditionality on the fibre structure.
10. *Quaternionic rigidity* (O27 [26]): every admissible morphism $\Phi_{q,\rho}$ factors through $\mathfrak{su}(2)$.
11. *Eigenvalue ratio* (O30 [38]): the invariant ratio $\lambda_1/\lambda_2 = 2$ of the per-pair covariance is analytically derived from the equatorial admissible fibre constraint.
12. *Generator-twist isospectrality* (O31 [40]): the spectra of $\rho_c(P_{S_q})$ and $\rho_c(P_{\varphi_\omega(S_q)})$ are equal for all c , but this does not imply colour-triplet spectral degeneracy. This closes the spectral route to [H-color].

7.2 Structural results

The following results are derived from structural arguments that are internally consistent but not yet fully formalised as theorems.

1. *Phenomenological window* (O3 [22]): $\beta^* \in (0.09, 0.13)$ from $m_e : m_\mu : m_\tau$.
2. *Capacity exponent target* (O7 [20]): the target $\delta \in [7.4, 10.6]$ on three-step path fingerprints.
3. *Observable-class identification* (O16 [9]): the pair observable resolves the gap by exponent doubling $\delta_{\text{pair}} = 2\delta_c$.
4. *$SU(3)$ emergence framework* (O31 [40]): the metaplectic intertwining of $\text{Heis}_3(\mathbb{Z}/q\mathbb{Z})$ forces triplet co-admissibility unconditionally for the colour-adapted Cayley graph.

7.3 Numerical evidence

The following results are numerically established but not yet derived analytically.

1. $\delta_{\text{corr}}(q) \in [7.7, 9.0]$ (O25 [33]): for $q \in \{61, 101, 151, 211\}$ after O14 normalisation; consistently within the admissible window $[7.4, 10.6]$.
2. *Spin- $\frac{1}{2}$ sector* (O29 [31]): $r_{\text{eff}} = 3$ and $d_\rho = 2$ confirmed for $q \in \{61, 101, 151\}$.
3. *Eigenvalue structure* $[1 : \frac{1}{2} : \frac{1}{2}]$ (O28 [5]): robust across all conjugate pairs and all tested primes.
4. [H-color] *numerical support* (O32 [39]): passes for $q \in \{61, 151, 211, 307\}$ with variance ratio $R_{\text{var}} \leq 1$ relative to the control noise floor.

7.4 Open hypotheses

The following items remain open within the sub-programme.

1. [H-color]_{pointwise}: O32 [39] establishes three of the four levels of the hypothesis analytically (rank equality in O31, average equality to $O(q^{-1})$, effective exponent equality in the $q \rightarrow \infty$ limit). The remaining open level is exact pointwise profile equality at finite q and M . The spectral route to this level is closed (O31): the correct mechanism must act directly on the rank structure of the BFS walk, not on Markov spectra.
2. *Asymptotic stabilisation of δ_∞* : the limit $\delta_\infty = \lim_{q \rightarrow \infty} \delta_{\text{pair}}(q)$ has not been computed analytically. The correct asymptotic variable is $n_1(q)/q$, not q itself.
3. *Extension to $q = 401$* : the campaign at $q \in \{29, 61, 101, 151, 211, 307\}$ is complete; $q = 401$ is pending.

8 Open Deliverables

Two open deliverables define the boundary of the sub-programme as of this note.

Open deliverable 1: Pointwise proof of [H-color]. O32 [39] establishes a three-tier resolution of this hypothesis. (i) [H-color]_{rank}: sector rank equality $r_c(n) = r_{\omega c}(n) = r_{\omega^2 c}(n)$ is proved unconditionally in O31 [40] via intra-coset Vandermonde symmetry. (ii) [H-color]_{avg}: the block-averaged capacity satisfies $\mathbb{E}[\sigma_c(n)] = \mathbb{E}[\sigma_{\omega c}(n)] + O(q^{-1})$, proved in O32 (Proposition 1) via the modulation frequency argument. (iii) [H-color]_{eff}: the effective exponents satisfy $\delta_c = \delta_{\omega c} = \delta_{\omega^2 c}$ in the $q \rightarrow \infty$ limit, proved in O32. What remains open is [H-color]_{pointwise}: exact profile equality at finite q and M , which is consistent with all numerical results but not claimed analytically. The SU(3) identification on the standard graph is therefore analytically established at the level of effective exponents; the remaining open question is the finite- q residual variance $R_{\text{var}}(q) \propto q^{-1}$, to be confirmed at $q = 401$.

Open deliverable 2: Numerical campaign at $q = 401$. The campaign at $q \in \{29, 61, 101, 151, 211, 307\}$ is complete. $q = 307$ confirms [H-color] ($R_{\text{var}} = 1.17 \times 10^{-3}$, all 4/4 triplets, O32 [39]). The extension to $q = 401$ is the sole remaining numerical task: it constrains the rate of convergence of $n_1(q)/q$ and provides the fifth data point for the $R_{\text{var}}(q) \propto q^{-1}$ scaling prediction.

All other open problems of earlier papers (e.g., the rigorous continuum limit of the state law, the exact form of Φ on LPS graphs) are sub-problems within already resolved phases and do not block the downstream branches.

9 Conclusion

The spectral admissibility sub-programme provides the core computational content of the Cosmochrony corpus. Starting from the Born–Infeld bounded-flux constraint on the Weil representation of $\text{Heis}_3(\mathbb{Z}/q\mathbb{Z})$, it derives the cascade exponent $\beta^* \approx 0.126$ unconditionally, identifies the spin- $\frac{1}{2}$ admissible sector, and establishes the admissibility thread $Q_8 \subset 2I \subset \text{SU}(2)$ as a structural output.

The sub-programme is analytically closed for the $\text{SU}(2)$ sector. It is closed numerically for $q \in \{29, 61, 101, 151, 211, 307\}$. The extension to $\text{SU}(3)$ is conditional on hypothesis [H-color], whose analytical proof remains an open problem in Hecke spectral theory.

The outputs of this sub-programme are used throughout Branch III: by the quantum sector (Q1–Q3), by the geometric emergence sub-programme (Q5a–Q11), by the gauge structure sub-programme (Q6a, O31–O32), and by the fermionic matter papers (Q14). Reading the O-series without this map is possible but significantly more difficult; this note is intended to make the logical structure transparent before the technical machinery is engaged.

References

- [1] J. Beau. Three-Step Path Fingerprints on LPS Graphs and the Capacity Growth Exponent, 2025. Spectral admissibility sub-program "O8" paper, doi:10.5281/zenodo.19151769.
- [2] J. Beau. About Canonical Pair Observables in Weil Blocks: Structure of the Residual Amplitude Factor and Pipeline-Independent Formulation, 2026. O19 paper, doi:10.5281/zenodo.19323720.
- [3] J. Beau. Admissible Frontier Saturation and the Cascade Exponent: A Representation-Theoretic Obstruction and its Matrix-Level Refinement, 2026. Spectral admissibility sub-program "O5" paper, doi:10.5281/zenodo.19120328.
- [4] J. Beau. Admissible Non-Injective Transitions as the Primitive of Physical Description, 2026. Cosmochrony Foundation Paper "M", doi:10.5281/zenodo.20258438.
- [5] J. Beau. Asymptotic Calibration of the BFS Window and Effective Dimension of the Admissible Trajectory, 2026. O28 paper, doi:10.5281/zenodo.19767801.
- [6] J. Beau. Asymptotic Saturation of Projective Resolution: Expander Relaxation Graphs, 2026. Spectral admissibility sub-program paper, doi:10.5281/zenodo.19056975.
- [7] J. Beau. Asymptotic Stability of Exact Weil-Block Capacity on Heisenberg Graphs: Extended Prime Range, Variance Reduction, and Requalification of the $\delta \rightarrow \beta^*$ Tension, 2026. Spectral admissibility sub-program "O13" paper, doi:10.5281/zenodo.19210831.
- [8] J. Beau. Exact Weil-Block Projective Capacity on Heisenberg Graphs: Resolving the Final Obstruction to δ Extraction, 2026. Spectral admissibility sub-program "O12" paper, doi:10.5281/zenodo.19194798.
- [9] J. Beau. Fibre Admissibility and Resolution of the Capacity Exponent Discrepancy, 2026. Spectral admissibility sub-program "O16" paper, doi:10.5281/zenodo.19298427.
- [10] J. Beau. Fibre-Level Admissibility from Conjugate Weil Blocks: Structural Derivation of the Pair Observable, 2026. Cosmochrony Spectral Admissibility sub-Program "O17" paper, doi:10.5281/zenodo.19299611.
- [11] J. Beau. Matrix-Level Dynamic Redundancy and the Structural Confinement of the Cascade Exponent β , 2026. Spectral admissibility sub-program "O6" paper, doi:10.5281/zenodo.19140624.
- [12] J. Beau. Minimal Fibre Structure of the Non-Injective Projection from Born–Infeld Indiscernability: Derivation of the Parity Involution, 2026. Spectral admissibility sub-program "O18" paper, doi:10.5281/zenodo.19321413.
- [13] J. Beau. No External Dimensional Scale at the Level of Admissibility, 2026. Cosmochrony structural note, doi:10.5281/zenodo.19744767.
- [14] J. Beau. Non-Factorisability and the Emergence of Heisenberg Structure from Admissibility Constraints, 2026. spectral admissibility subprogram paper, doi:10.5281/zenodo.19635395.
- [15] J. Beau. Non-Injectivity as a Structural Necessity of Genuine Emergence: A No-Go Theorem for Faithful Emergent Descriptions, 2026. Cosmochrony companion paper "ENI" or L, doi:10.5281/zenodo.20236950.
- [16] J. Beau. Observable-Class Derivation Mismatch and the Exact Growth Equation: From Proxy Capacity to Block-Level Dynamics, 2026. O15 spectral admissibility subprogram paper, doi:10.5281/zenodo.19281420.

- [17] J. Beau. Observable-Class Mismatch and the Corrected $\delta \mapsto \beta^*$ Relation on Heisenberg Graphs: Block Normalisation, Central-Phase Contribution, 2026. O14 spectral admissibility subprogram paper, doi:10.5281/zenodo.19226272.
- [18] J. Beau. Pairwise Character Geometry and the Classification of Neutral Generators in $SU(2)$, 2026. Spectral admissibility sub-program "rigidity" paper, doi:10.5281/zenodo.18969867.
- [19] J. Beau. Polynomial-Growth Capacity Dynamics at Large q : Volumetric Growth Confirmation, Capacity Decay, and the Weil Decomposition Roadmap, 2026. Spectral admissibility sub-program "O10" paper, doi:10.5281/zenodo.19161074.
- [20] J. Beau. Projective Capacity and the Continuum Limit of Admissible Redundancy, 2026. Spectral admissibility sub-program "O7" paper, doi:10.5281/zenodo.19149773.
- [21] J. Beau. Projective Capacity Beyond Expanders: Polynomial-Growth Relaxation Graphs and the Capacity Exponent, 2026. Spectral admissibility sub-program "O9" paper, doi:10.5281/zenodo.19154517.
- [22] J. Beau. Projective Dynamics and Mass Hierarchy: Hierarchical Amplification via Growing Relational Valence, 2026. Spectral admissibility sub-program "O3" paper, doi:10.5281/zenodo.19076244.
- [23] J. Beau. Projective Persistence and the Physical Sub-Spectrum: A Dynamical Selection Criterion for the Capacity Exponent, 2026. Spectral admissibility sub-program "O20" paper, doi:10.5281/zenodo.19339888.
- [24] J. Beau. Projective Resolution Dynamics on Variable-Valence Relaxation Graphs, 2026. O1 spectral admissibility subprogram "O1" paper, doi:10.5281/zenodo.19076319.
- [25] J. Beau. Quadratic Completion of Admissible Spectral Pairs via Binary-Icosahedral Representation, 2026. O26 paper, doi:10.5281/zenodo.19488845.
- [26] J. Beau. Quaternionic Rigidity of Admissible Morphisms: Every Admissible $\Phi_{q,\rho}$ Necessarily Factors through $\mathfrak{su}(2)$, 2026. O27 paper, doi:10.5281/zenodo.19489296.
- [27] J. Beau. Shell-Alignment from Projection Locking: A Discrete Admissibility Theorem under Born-Infeld Saturation, 2026. Spectral admissibility sub-program "O22" paper, doi:10.5281/zenodo.19367375.
- [28] J. Beau. Shell-Level Saturation and the Fibre-Level Transfer $\delta_{\text{pair}} \rightarrow \beta^*$, 2026. Spectral admissibility sub-program "O21" paper, doi:10.5281/zenodo.19358961.
- [29] J. Beau. Spectral Admissibility under Bounded Flux: Non-Abelian Mode Selection on Cayley Graphs of $SU(2)$ Subgroups, 2026. Step A of Cosmochrony Spectral Admissibility sub-Program, doi:10.5281/zenodo.18944985.
- [30] J. Beau. Spectral Capacity Functional and the Binary-Polyhedral Maximality Conjecture, 2026. Step B of Cosmochrony Spectral Admissibility sub-Program, doi:10.5281/zenodo.18945273.
- [31] J. Beau. Spin- $\frac{1}{2}$ Sector Identification via the Symmetric Rank Formula: Effective Dimension of the Admissible Covariance in $\text{End}(V_\rho)$, 2026. O29 paper, doi:10.5281/zenodo.19787221.
- [32] J. Beau. Subdiffusive Valence Growth under Bounded Relational Flux: Structural Derivation of the Cascade Exponent β , 2026. Spectral admissibility sub-program "O4" paper, doi:10.5281/zenodo.19101472.

- [33] J. Beau. Systematic Pair-Level Campaign for δ_{pair} : Convergence, Inter-Pair Concentration, and Normalization Structure, 2026. O25 paper, doi:10.5281/zenodo.19462132.
- [34] J. Beau. Three Particle Generations and Mass Hierarchy from Spectral Stratigraphy, 2026. Spectral admissibility sub-program paper, doi:10.5281/zenodo.19040490.
- [35] J. Beau. Three Stable Directions from Quaternionic Minimality: Derivation of $\Sigma_c(n_3) = 3$ from Born–Infeld Fibre Admissibility, 2026. Spectral admissibility sub-program "O23" paper, doi:10.5281/zenodo.19375136.
- [36] J. Beau. Vertical Non-Injectivity and the Stability of the Observable Rank: Closing the Unconditional Transfer $c_\chi \rightarrow \delta_{\text{pair}} \rightarrow \beta^*$, 2026. O24 spectral admissibility subprogram "O24" paper, doi:10.5281/zenodo.19386581.
- [37] J. Beau. Weil-Block Projective Capacity on Heisenberg Graphs: Resolving the Representation Obstruction and Extracting the Capacity Exponent, 2026. Spectral admissibility sub-program "O11" paper, doi:10.5281/zenodo.19169987.
- [38] Jérôme Beau. Born–Infeld Parity Forces an Equatorial Admissible Fibre: Analytical Derivation of the $[1 : \frac{1}{2} : \frac{1}{2}]$ Eigenvalue Ratio in $\text{Sym}^2(V_\rho)$, 2026. "O30" paper, doi:10.5281/zenodo.20014707.
- [39] Jérôme Beau. Colour Triplet Co-admissibility on $\text{Heis}_3(\mathbb{Z}/q\mathbb{Z})$: Numerical Test of Hypothesis [H-color], 2026. "O32" paper, doi:10.5281/zenodo.20333304.
- [40] Jérôme Beau. $\text{SU}(3)$ from Colour Triplet Co-admissibility: Towards the Complete Standard Model Gauge Group, 2026. "O31" paper, doi:10.5281/zenodo.20333303.