

# Effects of below-ground space limitation on performance of Eucalyptus seedlings: Does photosynthesis really control growth?

Courtney E. Company<sup>1,3</sup>, Belinda E. Medlyn<sup>2</sup> and Remko A. Duursma<sup>1</sup>

<sup>1</sup>Hawkesbury Institute for the Environment, University of Western Sydney,  
Richmond, NSW 2753, Australia

<sup>2</sup>Department of Biological Sciences, Macquarie University, North Ryde, NSW 2109,  
Australia

<sup>3</sup>Corresponding author (c.company@uws.edu.au)

January 23, 2015

## Abstract

Interpreting limitations to plant growth requires understanding of the balance between carbon (C) source and sink activity in order to assess carbon allocation and biomass partitioning. This study used manipulations of soil volume to test how growth is coupled to physiology, allocation, and sink activity in Eucalyptus tereticornis seedlings. We grew seedlings in a large range of container sizes and planted containers flush to the soil alongside naturally sown seedlings (free). Reduced soil volume was expected to induce rapid negative effects on growth and physiology compared to free seedlings. It was hypothesized that soil volume effect would be largest in the smallest containers, resulting in physical constraints to growth independently of photosynthesis. Photosynthesis would then become sink-limited, resulting in the build-up of leaf nonstructural carbohydrates eventually leading to photosynthetic down regulation. We observed a negative container effect on aboveground growth soon after the experiment started. Although growth was consistently different across soil volumes mass, partitioning to leaves, stems, roots was conserved after 120 days. Photosynthetic capacity was also significantly reduced in containers, and was related to both leaf nitrogen content and starch accumulation, however starch effects were weaker. We developed a simple seedling growth model that utilized leaf photosynthesis rates to allocate daily C uptake towards mass growth of stems, leaves and roots. We then asked whether the observed reductions in photosynthesis explained the observed differences in seedling biomass. We found that photosynthesis reductions alone were not sufficient and the inclusion of additional carbon and mass allocation strategies were necessary to better predict the observed growth responses to decreasing soil volume. Overall, we found limited evidence for sink-limitation of photosynthesis by constraining seedling growth in containers, and argue that shifts in C allocation are necessary to explain the experimental results. This research highlights the need to further understand adaptive strategies of plant C allocation, and confirms that photosynthesis and growth are not always directly related.

## Key Words

photosynthesis, sink regulation, growth, carbon allocation, soil volume

## 38 Introduction

39 For plants to grow they need to assimilate sufficient carbon (C) to allow the buildup of struc-  
40 tural biomass and to fuel the metabolism associated with biomass maintenance (Sala et al., 2012).  
41 Roots uptake the nutrients, such as nitrogen (N), required for leaf growth and leaves fix the C  
42 required for growth and maintenance of the entire plant. This growth is driven by several simulta-  
43 neous processes, including photosynthesis (A), C investment among organs, resources acquisition  
44 and metabolic costs (Körner, 2006; Fourcaud et al., 2008). Additionally a major component of dry  
45 matter investment is a storage pool of carbohydrates used to fuel growth and metabolism when  
46 photosynthesis does not meet demand (Chapin et al., 1990). The concentration of storage car-  
47 bohydrates within plant tissues depends on the passive balance between C supply, via A, and C  
48 demands through growth and respiration (Mitchell et al., 2013), or as an actively regulated process  
49 to maintain hydraulic transport across tissues (Sala et al., 2012). Thus, understanding plant growth  
50 across spatial and temporal scales requires integration of the uptake of C and N in coalescence,  
51 as well as patterns of allocation across plant tissues

52 The distribution of assimilated C is a primary determinant of plant growth (Friedlingstein et al.,  
53 1999), yet our knowledge of the mechanisms by which allocation is regulated is poor (Poorter  
54 et al., 2012b). The allocation of C involves the shifting of the products of A between respiration and  
55 biomass production, ephemeral and long-lived tissues, and above and belowground components  
56 (Litton et al., 2007). With non-limiting availability of nutrients, light, and water a plant should al-  
57 locate resources to leaves in order to maximize C acquisition and height growth. For balanced  
58 growth between leaves and roots it is generally assumed that biomass is preferentially allocated  
59 to obtain the most limiting resource, however, it is unlikely that plants ever reach a dynamic equi-  
60 librium with allocation as supply rates of light and soil resources fluctuate continuously (Shipley  
61 and Meziane, 2002). Alternatively, acclimation of plant traits, via phenotypic plasticity of leaf and  
62 root morphology, can improve the capacity of resource capture without resulting shifts in biomass  
63 allocation (Waisel et al., 1996), especially in young plants (Reich et al., 2002). As a result, empiri-  
64 cal evidence is needed to understand the mechanisms driving patterns of C investment, biomass  
65 allocation, and trait acclimation when resource availability is not optimal.

66 Due to the multitude of mechanisms driving both A and C allocation there exists ongoing con-  
67 troversy over the hierarchy of processes controlling plant growth. Generally, greater rates of A  
68 are assumed to produce more growth but this common view ignores both the rules of sto-  
69 ichiometry and neglects processes such as C transport, respiration and storage (Körner, 2013).  
70 Furthermore, focus on C allocation and storage is often first placed on the fate of carbon that  
71 has been recently assimilated by A (Atkin, 2015). As a result, the inhibition of A by sink limitation  
72 related to water and nutrient availability or a reduction in whole plant carbon demand with build  
73 of non-structural carbohydrates (TNC) may be incorrectly ignored. As A is linked to whole plant  
74 physiology by reciprocal controls, these top down views can ignore how source-sink interactions  
75 can override direct control of photosynthesis by light and CO<sub>2</sub> (Paul and Foyer, 2001). When testing  
76 how environmental controls affect plant growth it is therefore essential to understand how sink  
77 strength of component tissues is affected in combination with A in order to understand strategies  
78 of C allocation within the whole plant.

79 Plants undergo many physiological and morphological changes in response to rooting volume,  
80 potentially affecting root/shoot growth, biomass partitioning, net photosynthetic rate, water re-  
81 lations, nutrient uptake and respiration (NeSmith and Duval, 1998, and references therein). Thus  
82 exposing plants to differential soil volumes, via container size, can be used to manipulate seedling  
83 growth and subsequently test the mechanisms that affect the balance between C sink and source  
84 activity. Recently, a meta-analysis on the effects on rooting volume reported that reduction in  
85 growth is mainly caused by a reduction in A per unit leaf area (Poorter et al., 2012a), yet the causal-  
86 ity of this reduction is unclear. The volume of containers in which plants are grown is thought to  
87 decrease the sink strength for C by reducing root growth and plants grown in small containers are  
88 more likely to exhibit photosynthetic down-regulation (Arp, 1991; McConnaughay and Bazzaz, 1991;  
89 Gunderson and Wullschlegel, 1994; Sage, 1994; Maina et al., 2002; Ronchi et al., 2006). Smaller con-  
90 tainers may also reduce N uptake, either from physical root restriction or decreased supply, and N

deficiencies can affect growth, Rubisco limitation, sugar metabolism, and carbohydrate partitioning between source and sink tissues (Stitt, 1991; Hermans et al., 2006). Alternatively, plants in small soil volumes may alter root morphology under root restriction (Krizek et al., 1985) or leaf morphology if nitrogen uptake is altered (Reich et al., 1998). By utilizing what is known about the biological constraints of small containers we can empirically test how growth strategies are adapted when seedlings reach differential thresholds of soil volume to exploit.

This study tests how *Eucalyptus tereticornis* seedlings adjust biomass allocation and physiology across a range of limiting soil volumes and compares these responses to field grown seedlings without root restriction. Seedlings were maintained under well watered conditions and the plasticity of seedlings to maintain optimal photosynthetic capacity and the duration of this response was concomitantly tracked with patterns of biomass allocation and leaf traits to understand processes that regulate seedling growth. Reducing soil volume was expected to induce rapid negative effects on growth and physiology compared to free seedlings. It was hypothesized that soil volume effect would be largest in the smallest containers, resulting in physical constraints to growth independently of photosynthesis. Photosynthesis would then become sink-limited, resulting in the build-up of leaf nonstructural carbohydrates leading to photosynthetic down regulation through time as a function of available soil volume.

## Methods

### Experimental Design

This experiment was located on the Hawkesbury Forest Experiment site in Richmond, NSW, Australia. Plots were located in open cover with a site history that consists of a paddock that was converted from native pasture grasses. Top soils at this site, used for the study, are an alluvial formation of low-fertility sandy loam soils (380 and 108 mg kg<sup>-1</sup> total nitrogen and phosphorus respectively) with low organic matter (0.7 %) and low water holding capacity. At this site a soil hard layer exists at ~1.0 m with a transition to heavy clay soils. The climate for the region is classified as sub-humid temperate.

*Eucalyptus tereticornis* seedlings, 20 weeks old and approximately 40 cm tall in tube stock, were chosen from a single local Cumberland plain cohort. Previous experiments have confirmed that species with tap roots (similar to *E. tereticornis*) use the center of the container as the medium for thick roots leaving the periphery of the soil as the most active sites for fine root proliferation (Biran and Eliassaf, 1980a,b). This is generally hypothesized to be a different response than seedlings with no taproot. By using a species with tap root growth and manipulations of container length rather than width, it is believed that a more realistic test of inhibition of growth through constrained soil volume would be achieved. Six seedlings were harvested before planting to measure initial leaf area and dry mass of leaves, stems and roots.

Six container volumes were used ranging from 5 l to 3 l, with a 22.5 cm diameter, and lengths ranging from 15 to 100 cm. Containers were constructed of PVC pipe and were filled with local top soil (described above). Soil in each container was packed to achieve a target soil bulk density of 1.7 g m<sup>-3</sup>. A Imidacloprid (BAYER CropScience) insecticide tablet was planted 5 cm below the roots of each seedling. Containers were planted flush with the soil surface inside metal sleeves, designed to minimize excess air space between the container and outside soil while also allowing for container removal. This allowed for soil temperatures in containers to reflect conditions of naturally sown (free) seedlings. Each experimental block (n=7) contained a complete replicate set of container volumes as well as one free seedling, with 1 m<sup>2</sup> spacing. For each free seedling, a 1 m<sup>2</sup> subplot was excavated to 0.5 m and replaced with the same soil used in each container. For free seedlings a border of root exclusion material was buried 0.25 m deep and extended 0.25 m above the ground surface around each subplot to exclude local vegetation.

Plants were watered bi-weekly or when needed, accounting for natural precipitation, to maintain soil moisture at field capacity (13-15 %). To achieve field capacity all soil filled containers were

weighed and soil moisture was measured (Time Domain Reflectometer at 10 cm depth) before watering. Derived equations based on container weight at planting (when soil moisture was 3 %) and at field capacity was used to calculate watering requirements for each individual plant. Drain systems were built into each pot to prevent pooling of water in containers before root expansion, from reduced root uptake, or from large rainfall events. These conditions could lead to an anaerobic environment around the root that could hinder the uptake of water through reduced root conductance (Poorter et al., 2009), an undesired experimental artifact. A collection compartment in the bottom of containers, containing gravel covered by root exclusion mesh, was used to collect excess water for 20, 25, and 35 l containers. Plastic tubing (6 mm diameter) was inset into the gravel layer and extended through the top of the container. A lysimeter pump was then used to suction excess water, through the tubing, as needed. As small containers (5, 10, and 15 l) have a larger irradiation effect a simple bottom plug was used to drain excess water from the gravel compartment.

### ***Growth and morphology metrics***

Seedlings were planted on January 21<sup>st</sup> 2013 and stem height, diameter at 15 cm and leaf count were measured weekly thereafter. Once the growth rate of individual plants had significantly declined a full biomass harvest was completed (May 21<sup>st</sup> 2013). Dry mass of leaves, stems, roots and cumulative leaf area (LI-3100C Area Meter; LI-COR, Lincoln, NE, USA) was measured for each seedling. Mean individual leaf area for each harvested seedling was calculated by dividing cumulative leaf area by total leaf count of only fully expanded leaves. This value was then used to interpolate cumulative leaf area through time with weekly leaf counts. Root mass was collected by passing soil from each container through a 1 mm sieve, washing, separating into fine and coarse roots (<2 mm and >2 mm diameter, respectively) and then drying to a constant mass. Roots from the free seedlings were collected by excavating each 1 m<sup>2</sup> subplot to 0.5 m depth. 25 g fresh weight subsamples of washed fine roots were analyzed, using WhinoRhizo software (Regent Instruments Inc.), for specific root length (SRL, cm m<sup>-1</sup>).

### ***Photosynthetic parameters***

Leaf gas exchange measurements were performed bi-weekly at saturating light ( $A_{\text{sat}}$ ) and saturating light and  $[\text{CO}_2]$  ( $A_{\text{max}}$ ) on new fully expanded leaves. Measurements were initiated only after sufficient new leaf growth occurred (March 17<sup>th</sup>, 2013), approximately 6 weeks following planting, and continued until the biomass harvest. Leaf level gas exchange was measured with a standard leaf chamber equipped with blue-red light emitting diodes using a portable gas exchange system (LI-6400, LI-COR, Lincoln, NE, USA).  $A_{\text{sat}}$  measurements were made at PPFD of 1800  $\mu\text{mol m}^{-2} \text{s}^{-1}$  and  $[\text{CO}_2]$  of 400  $\mu\text{l l}^{-1}$  and  $A_{\text{max}}$  with  $[\text{CO}_2]$  of 1600  $\mu\text{l l}^{-1}$  and PPFD of 1800  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ . This choice of light level to achieve light saturation is consistent with other studies on *Eucalyptus* species (Kallarackal and Somen, 1997; Pinkard et al., 1998; Crous et al., 2013; Drake et al., 2014). These measurements were conducted during midday (10:00-14:00 h) with leaf temperature maintained at 25°C. After leaves acclimated to the chamber environment, net  $\text{CO}_2$  assimilation rate and stomatal conductance ( $g_s$ ) were logged 5 times for both  $A_{\text{sat}}$  and  $A_{\text{max}}$ . Photosynthetic  $\text{CO}_2$  response ( $\text{AC}_i$ ) curves were also developed at 25°C on a random subset of each container size ( $n=3$ ) after new leaves were first produced and immediately prior to the final harvest (May 23<sup>rd</sup> 2013). Each  $\text{AC}_i$  curve began at the reference  $[\text{CO}_2]$  of 400  $\mu\text{l l}^{-1}$  and then consisted of 12 additional steps from  $[\text{CO}_2]$  of 50 to 1800  $\mu\text{l l}^{-1}$  at 25°C and saturating light (above). From these curves the photosynthetic parameters,  $J_{\text{max}}$  and  $V_{\text{cmax}}$ , can be quantified using biochemical model of Farquhar et al. (1980). Leaf dark respiration rates ( $R_d$ ) was measured on each seedling during the same dates as  $\text{AC}_i$  curves using detached leaves inside a conifer chamber attached to the Licor 6400 at least 1 hour after sundown. Measurements were taken at a reference  $[\text{CO}_2]$  of 400  $\mu\text{l l}^{-1}$  while leaf temperature was maintained at current ambient conditions. Specific leaf area (SLA,  $\text{m}^2 \text{kg}^{-1}$ ) was calculated by measuring leaf area and dry mass for individual leaves sampled during gas exchange campaigns.

## 190 **Leaf water potential**

191 Predawn ( $\Psi_{pd}$ ) and midday ( $\Psi_l$ ) leaf water potentials were measured for each seedling using a  
192 PMS 1505D pressure chamber (PMS Instruments, Albany, OR, USA) on fully expanded leaves dur-  
193 ing the same time period as  $AC_i$  and  $R_d$ . Leaves were collected, immediately stored inside foil  
194 covered bags and then transported to the laboratory for measurements of water potential.  $\Psi_{pd}$   
195 was measured before sunrise and  $\Psi_l$  at midday 13:00-14:30 h. These measurements were used  
196 as a measure of static water stress on the seedlings (Sellin, 1999), and to ensure that the bulk  
197 soil water availability was high enough for plants as they became larger and roots filled the soil  
198 volume.

## 199 **Leaf, root and soil chemistry**

200 Leaves used in each gas exchange measurements and subsamples of harvested roots were dried  
201 to a constant mass and milled for analysis of  $N$  content. Pre-planting soil samples ( $n=6$ ) and sub-  
202 samples of soil from each container following harvest were sieved to remove organic material,  
203 air dried and milled for analysis.  $N$  concentrations of milled samples were determined using a  
204 Carlo Erba CE1110 elemental analyzer with thermal conductivity and mass spectrometric detection  
205 (of  $N_2$  and  $CO_2$ ). The percentage of  $N$  in the sample was calculated by comparison with known  
206 standards. Leaf TNC concentration was analyzed using a total starch assay kit (Megazyme Inter-  
207 national 303 Ireland Ltd., Wicklow, Ireland) on gas exchange leaves (above) and represents the  
208 sum of starch ( $mg\ g^{-1}$ ) and soluble sugar ( $mg\ g^{-1}$ ) concentrations. Starch was quantified using a  
209 thermostable  $\alpha$ -amylase and amyloglucosidase assay (McCleary et al., 1997) and soluble sugars  
210 were determined following the anthrone method (Ebell, 1969). Full methods of the TNC assay are  
211 described in (Mitchell et al., 2013).

## 212 **Seedling growth model**

213 We developed a simple seedling growth model that utilized leaf photosynthesis to allocate daily  
214 Cassimilate towards mass production of stems, leaves and roots. The model begins with the  
215 mean initial component mass ( $leaf_i$ ,  $stem_i$  and  $root_i$ ) and a starting leaf area ( $LA_i$ ) measured prior  
216 to planting. Biomass production for each day of the experiment was estimated by multiplying the  
217 whole tree seedling carbon uptake gain per day ( $C_{day}$ ) by a  $C$  use efficiency parameter ( $Y_c$ , Mäkelä  
218 (1997)), allocating new carbon to the previous days biomass and then accounting for carbon loss  
219 via tissue respiration.  $C_{day}$  was calculated for each seedling by fitting a coupled photosynthe-  
220 sis - stomatal conductance model (Farquhar et al., 1980; Medlyn et al., 2011) in the 'plantecophys'  
221 package in R (Duursma, 2014) to the mean photosynthetic parameters ( $R_d$ ,  $J_{max}$ ,  $V_{cmax}$ , and  $g_i$ ) for  
222 each treatment and meteorological data from an onsite weather station. Examples of the photo-  
223 synthesis model are described in Medlyn et al. (2002) and the approach of the coupled leaf gas  
224 exchange model are described in Duursma et al. (2014). The  $g_i$  parameter was generated by fit-  
225 ting observed stomatal conductance values into the optimal stomatal conductance model from  
226 (Medlyn et al., 2012). Combined with the meteorological parameters PPFD, air temperature, and  
227 relative humidity, at 15 m intervals, photosynthesis rates ( $\mu mol\ CO_2\ m^{-2}\ s^{-1}$ ) were then predicted  
228 for each soil volume treatment. Rates were assumed to be representative of the entire 15 min  
229 meteorological interval.  $C_{day}$  was calculated by converting predicted rates to mass  $C$  gain over  
230 15 min ( $g\ m^{-2}$ ) and then summed for 24 h. Additionally, it is necessary to calculate a self-shading  
231 parameter ( $M$ ) when scaling flat leaf photosynthesis to seedling cumulative leaf area. This was ac-  
232 complished by utilizing 61 previously digitized Eucalyptus seedlings, covering 5 total species which  
233 include *E. tereticornis*, from Duursma et al. (2012b) to run in 'YplantQMC' package in R (Duursma  
234 et al., 2012a) to build a 3d plant structure based on digitized metrics of plant allometry and crown  
235 structure. Inputting the same physiological parameters listed above, 'YplantQMC' outputs total  
236 photosynthesis, using total leaf area, for seedlings assuming self-shading as well as for a full sun  
237 large horizontal leaf. The ratio of self-shading to flat leaf total photosynthesis was then used to

calculate  $M$ , which was then applied to modeled rates of photosynthesis when scaling to seedling cumulative leaf area. All default parameters used in model simulations are reported in Table. 3.

We then utilized this model to investigate whether the effects of soil volume on photosynthesis were enough to accurately predict overall seedling production after 120 days or whether changes in C allocation or mass partitioning were also necessary to predict biomass. First, the model was simulated using only changes in photosynthesis rates across treatments combined with mean values of mass partitioning and either published or local data of stem and root respiration rates. Next, different mass partitioning and C allocation scenarios were simulated to investigate sources of missing C from the initial model simulation. This included testing model sensitivity by altering leaf C allocation, root respiration rates, and fine root C allocation. For all cases, mass production was compared between model output and harvested seedlings to investigate what C allocation strategies beyond reduction in photosynthesis aid in prediction of seedling growth responses induced by limiting soil volume.

## Data Analysis

Differences in experimental parameters with soil volume were analysed by one-way analysis of variance (ANOVA) in R with individual containers as random effects and soil volume as a categorical fixed effect. Tukeys post-hoc tests were performed in conjunction with ANOVA to determine which specific paired comparisons among soil volumes were different. Mixed model ANOVAs of  $A_{\max}$  and leaf chemistry were performed using the 'nlme' package (Pinheiro et al., 2014) in R and  $r^2$  values of mixed models were computed as in Nakagawa and Schielzeth (2013). Tests of allometric relationships between biomass components were implemented using major axis regression in the 'smatr' package (Warton et al., 2012). Results were considered significant at  $P \leq 0.05$ .

## Results

### Growth and morphology metrics

In this open field study, colder temperatures and reductions in cumulative PPFD per day (Fig. 1) most likely lead to the reduced growth in the free seedlings in the final weeks of the experiment (Fig. 2). Combined with severe growth reductions in the smallest container volumes the experiment was chosen to be harvested after 120 days. Over the 120 days height, diameter, and leaf area diverged between container volumes (Fig. 2). First, seedling leaf area significantly diverged between soil volumes ( $P < 0.026$ ) during the 5<sup>th</sup> week of the experiment. Following this period both height (8<sup>th</sup> week) and then diameter (9<sup>th</sup> week) significantly deviated across soil volumes ( $P < 0.002$  &  $0.001$ , respectively). Negative growth effects then manifested as severely reduced height gain and declining leaf area through time with small soil volumes across the final two months of the experiment. Seedlings maintained diameter growth throughout the experiment, although marginal with smaller soil volumes in the final month. Final seedling height significantly increased with increasing soil volume ( $P < 0.001$ ). Increases in both final stem diameter ( $P < 0.001$ ) and cumulative leaf area (both  $P < 0.001$ ) were found with increasing soil volume and these differences were driven mainly by the largest container and the free seedling.

Total plant biomass at harvest was significantly different across container volumes ( $P < 0.001$ ) and with free seedlings ( $P < 0.001$ , Table. 1). We analyzed the relationship between biomass growth with each fold increase in soil volume and found an increase of 34 % with a doubling of pot size, consistent with the meta-analysis of Poorter et al. (2012a). Additionally, plant biomass was highly correlated with total leaf area across all treatments ( $r^2 = .97$ ,  $P < 0.001$ ). Differences in biomass partitioning to leaves, stems, and roots were not different across soil volumes when variation in seedling biomass within treatments was factored in the analysis. Across all treatments, the final harvested root:shoot was conserved in these seedlings, with a slightly higher shoot than root mass on average ( $\bar{x} = 0.904$ ).

SRL of harvested fine roots was not different across soil volumes (Table. 1). Over the duration of the experiment SLA was higher in free seedlings but was not different across containers sizes (Table. 1,  $P < 0.001$ ) and this pattern was evident in the first gas exchange measurement campaign ( $P < 0.001$ ).

### **Leaf and root chemistry**

Leaf  $N$  % was significantly higher in free seedlings and the largest container volume at the onset of gas exchange measurements (6th week,  $P < 0.001$ ). Over the remaining duration of the experiment the smallest container volume had a significant reduction in leaf  $N$  % compared to other soil volumes, while the free seedling maintained a greater leaf  $N$  % (Table. 1,  $P < 0.001$ ). Additionally, mean leaf starch content in the smallest container was double that of free seedlings ( $P = 0.042$ ), while leaf soluble sugars did not differ across treatments throughout the experiment (Table. 1). Differences in leaf starch between the free seedling and the smallest container were also evident during the first gas exchange campaign ( $P = 0.0013$ ).

### **Gas exchange and photosynthetic parameters**

$A_{sat}$  (Fig. 3) and  $A_{max}$  (Table. 2) were both significantly higher in the largest container volume and the free seedling at the first measurement campaign (both  $P < 0.001$ ). Across measurement campaigns  $A_{sat}$  was consistently higher in free seedlings than in containers (Figure 3,  $P < 0.001$ ). The interaction between photosynthetic capacity, leaf starch, and leaf  $N$ , on a mass basis, was marginally significant ( $P = 0.0584$ ) but  $A_{max}$  was highly correlated to both leaf  $N$  content and leaf starch (both  $P < 0.001$ ). Across all measurement campaign  $A_{max}$  was higher when foliar  $N$  was also higher, usually associated with low levels of leaf starch (Fig. 3a).  $A_{max}$  was also lower when leaf starch was high as higher leaf  $N$  often did not coincide with high leaf starch (Fig. 3b).

The photosynthetic parameters  $J_{max}$  and  $V_{c_{max}}$  were not different across measurement campaigns, therefore the parameter means per volume are reported here (Table. 2). Overall, both  $J_{max}$  and  $V_{c_{max}}$  were significantly higher in free seedlings with little variation between container volumes ( $P = 0.0012$  &  $0.0021$ , respectively). Leaf dark respiration rates were not significantly different across soil volumes (Table. 2). The  $g_1$  parameter, generated for each seedling from the Medlyn et al. (2012) optimal stomatal conductance model, was not different across soil volumes (Table. 2). Predicted values of  $g_s$ , using the  $g_1$  parameter, were highly correlated with observed values ( $r^2 = .74$ ,  $P < 0.001$ , data not shown).

Neither  $\Psi_{pd}$  nor  $\Psi_l$  were different across treatments, with mean values of  $-0.27$  and  $-1.2$  mPa across all seedlings, respectively. Although  $g_s$  in free seedlings was generally higher than those in containers (Table. 2,  $P = 0.0023$ ), the mean rates for all seedlings were high at  $0.37 \text{ mol H}_2\text{O m}^{-2} \text{ s}^{-1}$  and did not decline significantly across the experiment duration. Combined these indices provide strong evidence that water stress was not apparent on these well-watered seedlings throughout the experiment. Soil  $N$  % at harvest was not different across soil volumes ( $\bar{x} = 0.45$  %) and decreased approximately 3 % across all containers over the experiment duration. This indicates that nutrient leaching from free seedlings or from draining of containers following natural rainfall events did not differ between treatments.

### **Modelling seedling biomass**

The initial model simulation, testing only treatment specific photosynthesis rates, was unable to accurately predict the responses of actual harvested seedling biomass to reduced soil volume. When both  $C_{day}$  and biomass of modeled and harvested seedlings were scaled to the free seedling the initial model design overestimated biomass for all treatments (Fig. 6a). This suggests that reductions in photosynthesis alone could not realistically capture the reduced growth

330 response of these seedlings. Next, the model was re-run using the mean mass fractions of  
331 leaves, stems and roots from the final harvest, per treatment, to drive daily mass partitioning  
332 and seedling production. Combined with treatment specific photosynthesis rates this model still  
333 overestimated seedling production for all treatments (Fig. 6b).

334 To test whether potential sources of missing C could improve the modeled response of seedling  
335 mass production the model was run with 3 different C allocation scenarios that could not be tested  
336 within the framework of this experiment. This was accomplished by testing the sensitivity of the  
337 model to adjustments of C allocation to leaves or fine roots or root respiration rates by  $\pm 50\%$   
338 of default values. The model output of these scenarios was then compared to both the original  
339 model output and with harvested seedling biomass. Increasing C allocation to either leaves or fine  
340 roots, in coalescence with photosynthesis rates, did improve the model but neither were sufficient  
341 to accurately predict the actual biomass response (Fig. S1a,c). Altering respiration rates of fine and  
342 coarse roots did not have a noticeable deviation from the original model (Fig. S1b).

## 343 Discussion

## 344 Tables



Table 1: Responses of plant and leaf characteristics of *Eucalyptus tereticornis* seedlings to soil volume treatments. Each value reflects the mean (standard error) of each treatment.

| Volume (L) | Seedling mass (g) | SLA (m <sup>2</sup> kg <sup>-1</sup> ) | Leaf Nitrogen (%) | Leaf Sugars (%) | Leaf Starch (%) | SRL (cm m <sup>-1</sup> ) | Root Nitrogen (%) |
|------------|-------------------|--|-------------------|-----------------|-----------------|---------------------------|-------------------|
| 5          | 14.8 (1.82)       | 9.5 (0.23)                             | 1.1 (0.02)        | 6.4 (0.28)      | 12.7 (0.97)     | 39.1 (5.47)               | 0.82 (0.05)       |
| 10         | 20.0 (2.38)       | 9.8 (0.24)                             | 1.3 (0.04)        | 6.7 (0.25)      | 9.4 (0.75)      | 34.2 (5.83)               | 0.75 (0.02)       |
| 15         | 25.4 (2.49)       | 11.0 (0.47)                            | 1.4 (0.06)        | 7.2 (0.28)      | 7.3 (0.73)      | 37.6 (4.63)               | 0.71 (0.02)       |
| 20         | 23.4 (1.63)       | 9.8 (0.28)                             | 1.4 (0.05)        | 6.6 (0.26)      | 9.5 (0.88)      | 45.3 (5.50)               | 0.76 (0.04)       |
| 25         | 30.4 (5.49)       | 10.4 (0.37)                            | 1.3 (0.06)        | 6.9 (0.24)      | 9.8 (0.71)      | 47.0 (7.10)               | 0.74 (0.02)       |
| 35         | 52.2 (9.55)       | 11.3 (0.44)                            | 1.5 (0.08)        | 6.8 (0.22)      | 9.8 (0.65)      | 50.6 (11.61)              | 0.77 (0.03)       |
| Free       | 174.5 (18.02)     | 13.0 (0.44)                            | 2.4 (0.09)        | 7.4 (0.25)      | 6.8 (0.65)      | 43.7 (6.24)               | 0.87 (0.04)       |

Table 2: Responses of leaf level gas exchange parameters of *Eucalyptus tereticornis* seedlings to soil volume treatments. Each value reflects the mean(standard error) of each treatment. Units for  $A_{\max}$  and  $R_{\text{dark}}$  are  $\mu\text{mol m}^{-1} \text{s}^{-1}$  and  $g_s$  are  $\text{mol m}^{-1} \text{s}^{-1}$ .

| Volume (L) | $A_{\max}$ | $R_{\text{dark}}$ | $J_{\max}$   | $V_{C\max}$ | $g_s$       | $g_1$     |
|------------|------------|-------------------|--------------|-------------|-------------|-----------|
| 5          | 21.2 (0.9) | 2.8 (0.3)         | 103.7 (5.8)  | 62.9 (2.8)  | 0.30 (0.01) | 5.1 (0.1) |
| 10         | 22.3 (1.4) | 2.7 (0.4)         | 116.5 (5.9)  | 69.4 (2.9)  | 0.36 (0.01) | 5.4 (0.1) |
| 15         | 23.3 (1.2) | 1.4 (0.1)         | 123.1 (15.4) | 79.4 (10.0) | 0.45 (0.01) | 6.2 (0.2) |
| 20         | 26.1 (0.7) | 1.7 (0.1)         | 130.0 (14.1) | 81.4 (6.0)  | 0.38 (0.01) | 5.2 (0.2) |
| 25         | 23.9 (0.9) | 1.2 (0.1)         | 131.9 (5.9)  | 78.6 (2.6)  | 0.32 (0.01) | 4.8 (0.2) |
| 35         | 25.0 (1.0) | 1.5 (0.2)         | 121.2 (3.7)  | 78.0 (3.5)  | 0.33 (0.02) | 4.7 (0.2) |
| Free       | 33.1 (0.7) | 1.3 (0.1)         | 171.7 (18.6) | 101.2 (5.7) | 0.49 (0.02) | 5.0 (0.2) |

## Figures




Figure 1: Daily maximum and minimum temperature (a), cumulative daily PPFD (b), and daily maximum vapour pressure deficit (c) across the experiment duration in 2013.




Figure 2: Soil volume treatment means  $\pm$  SE (n=7) of height growth (a), diameter growth (b), and interpolated seedling leaf area (c) measured weekly of *Eucalyptus tereticornis* seedlings across the experiment duration in 2013.




Figure 3: Soil volume treatment means  $\pm$  SE (n=7), across all measurement dates (n=6), of light saturated rates of photosynthesis at 25°C. Different letters represent significant differences between treatments.




Figure 4: Soil volume treatment means (n=7) of mass partitioning to leaves, stems, and roots (a) and bi-variate relationships between mass allocation to leaves and stems + roots (b). Lines represent standardized major axis fitting of the log transformed allometric relationships of leaf mass fraction by treatment.




Figure 5: Photosynthetic capacity, on a leaf mass basis, as a function of accumulation of leaf starch (a) and leaf nitrogen content without TNC (b). Colors represent bins levels ( $n=5$ ) of both leaf starch and nitrogen grouped from low to high. Lines represent predictions, for each bin level, from the linear mixed effects model equation of  $A_{\text{max}}$  as a function of starch and nitrogen. The marginal  $r^2$  (fixed effects only) was 0.37 and the conditional  $r^2$  (fixed and random effects) was 0.48 for the complete model.




Figure 6: Modeled and harvested seedling biomass (g) versus the measure range of daily assimilated carbon gain ( $\text{g m}^{-2}$ ) across all treatments (a) and then with treatment specific plant component mass partitioning included (b). Values of biomass and carbon gain are scaled to the free seedling with unlimited soil volume.

## 1 Supporting Information Figures

Table 3: Seedling Growth Model Default Parameters

| Variable                 | Default Value | Units                    | Source                   |
|--------------------------|---------------|--------------------------|--------------------------|
| Leaf area <sub>i</sub>   | 0.035         | m <sup>2</sup>           | this study               |
| Leaf mass <sub>i</sub>   | 3.45          | g                        | this study               |
| Stem mass <sub>i</sub>   | 1.51          | g                        | this study               |
| Root mass <sub>i</sub>   | 0.99          | g                        | this study               |
| Y <sub>c</sub>           | .65           | g C g mass               | Mäkelä (1997)            |
| R <sub>coarse root</sub> | 0.00124       | g C g root <sup>-1</sup> | Marsden et al. (2008)    |
| R <sub>fine root</sub>   | 0.010368      | g C g root <sup>-1</sup> | Ryan et al. (2010)       |
| R <sub>stem</sub>        | 0.00187       | g C g stem <sup>-1</sup> | Drake 2014 (unpublished) |
| C <sub>day</sub>         | 4.7-6.8       | g C m <sup>2</sup>       | this study               |
| M                        | 0.80-0.87     |                          | this study               |






Figure S1: Modeled and harvested seedling biomass (g) versus daily assimilated carbon gain (g m<sup>-2</sup>) including sensitivity testing for unmeasured carbon allocation scenarios. Model parameters of leaf allocation (a), root respiration (b), and fine root allocation (c) were increased or decreased by 50%.

## References

- W J Arp. Effects of source-sink relations on photosynthetic acclimation to elevated CO<sub>2</sub>. *Plant, Cell & Environment*, 14(8):869–875, 1991.
- Owen Atkin. New phytologist and the fate of carbon in terrestrial ecosystems. *New Phytologist*, 205(1):1–3, 2015.
- I Biran and A Eliassaf. The effect of container size and aeration conditions on growth of roots and canopy of woody plants. *Scientia Horticulturae*, 12(4):385–394, 1980a.
- I Biran and A Eliassaf. The effect of container shape on the development of roots and canopy of woody plants. *Scientia Horticulturae*, 12(2):183–193, 1980b.
- F Stuart Chapin, Ernst-Detlef Schulze, and Harold A Mooney. The ecology and economics of storage in plants. *Annual review of ecology and systematics*, pages 423–447, 1990.
- Kristine Y Crous, Audrey G Quentin, Yan-Shih Lin, Belinda E Medlyn, David G Williams, Craig V M Barton, and David S Ellsworth. Photosynthesis of temperate Eucalyptus globulus trees outside their native range has limited adjustment to elevated CO<sub>2</sub> and climate warming. *Global change biology*, 19(12):3790–3807, 2013.
- John E Drake, Michael J Aspinwall, Sebastian Pfautsch, Paul D Rymer, Peter B Reich, Renee A Smith, Kristine Y Crous, David T Tissue, Oula Ghannoum, and Mark G Tjoelker. The capacity to cope with climate warming declines from temperate to tropical latitudes in two widely distributed Eucalyptus species. *Global change biology*, 2014.
- Remko Duursma. *plantecophys: Modelling and analysis of leaf gas exchange data*, 2014.
- Remko Duursma, Mik Cieslak, Robert Pearcy, and Belinda Medlyn. *YplantQMC: Plant architectural analysis with Yplant and QuasiMC.*, 2012a. URL <http://www.remkoduursma.com/yplantqmc>, <https://www.bitbucket.org/remkoduursma/yplantqmc/>.
- Remko A Duursma, Daniel S Falster, Fernando Valladares, F J Sterck, R W Pearcy, Christopher H Lusk, K M Sendall, Marisa Nordenstahl, N C Houter, B J Atwell, and Others. Light interception efficiency explained by two simple variables: a test using a diversity of small-to medium-sized woody plants. *New Phytologist*, 193(2):397–408, 2012b.
- Remko A Duursma, Craig V M Barton, Yan-Shih Lin, Belinda E Medlyn, Derek Eamus, David T Tissue, David S Ellsworth, and Ross E McMurtrie. The peaked response of transpiration rate to vapour pressure deficit in field conditions can be explained by the temperature optimum of photosynthesis. *Agricultural and Forest Meteorology*, 189:2–10, 2014.
- L F Ebell. Variation in total soluble sugars of conifer tissues with method of analysis. *Phytochemistry*, 8(1):227–233, 1969.
- G D Farquhar, S von von Caemmerer, and J A Berry. A biochemical model of photosynthetic CO<sub>2</sub> assimilation in leaves of C<sub>3</sub> species. *Planta*, 149(1):78–90, 1980.
- Thierry Fourcaud, Xiaopeng Zhang, Alexia Stokes, Hans Lambers, and Christian Körner. Plant growth modelling and applications: the increasing importance of plant architecture in growth models. *Annals of Botany*, 101(8):1053–1063, 2008.
- Pierre Friedlingstein, G Joel, C B Field, and I Y Fung. Toward an allocation scheme for global terrestrial carbon models. *Global Change Biology*, 5(7):755–770, 1999.
- Carla A Gunderson and Stan D Wullschleger. Photosynthetic acclimation in trees to rising atmospheric CO<sub>2</sub>: a broader perspective. *Photosynthesis research*, 39(3):369–388, 1994.
- Christian Hermans, John P Hammond, Philip J White, and Nathalie Verbruggen. How do plants respond to nutrient shortage by biomass allocation? *Trends in plant science*, 11(12):610–617, 2006.
- J Kallarackal and C K Somen. An ecophysiological evaluation of the suitability of Eucalyptus grandis for planting in the tropics. *Forest Ecology and Management*, 95(1):53–61, 1997.

393 C Körner. Growth controls photosynthesis—mostly. *Nova Acta Leopoldina*, 114:273–283, 2013.

394 Christian Körner. Plant CO<sub>2</sub> responses: an issue of definition, time and resource supply. *New*  
395 *phytologist*, 172(3):393–411, 2006.

396 Donald T Krizek, Avner Carmi, Roman M Mirecki, FREEMAN W SNYDER, and JAMES A BUNCE. Com-  
397 parative effects of soil moisture stress and restricted root zone volume on morphogenetic and  
398 physiological responses of soybean [*Glycine max* (L.) Merr.]. *Journal of experimental botany*, 36  
399 (1):25–38, 1985.

400 Creighton M Litton, James W Raich, and Michael G Ryan. Carbon allocation in forest ecosystems.  
401 *Global Change Biology*, 13(10):2089–2109, 2007.

402 Godfrey G Maina, Joel S Brown, and Mordechai Gersani. Intra-plant versus inter-plant root com-  
403 petition in beans: avoidance, resource matching or tragedy of the commons. *Plant Ecology*, 160  
404 (2):235–247, 2002.

405 Annikki Mäkelä. A carbon balance model of growth and self-pruning in trees based on structural  
406 relationships. *Forest Science*, 43(1):7–24, 1997.

407 Claire Marsden, Yann Nouvellon, and Daniel Epron. Relating coarse root respiration to root diam-  
408 eter in clonal Eucalyptus stands in the Republic of the Congo. *Tree physiology*, 28(8):1245–1254,  
409 2008.

410 B. V. McCleary, T. S. Gibson, and D. C. Mugford. Measurement of total starch in cereal products by  
411 amyloglucosidase- $\alpha$ -amylase method: Collaborative study. *Journal of AOAC International*, 80(3):  
412 571–579, 1997. ISSN 1060-3271.

413 K D M McConnaughay and F A Bazzaz. Is physical space a soil resource? *Ecology*, pages 94–103,  
414 1991.

415 B E Medlyn, Erwin Dreyer, D Ellsworth, M Forstreuter, P C Harley, M U F Kirschbaum, Xavier Le  
416 Roux, Pierre Montpied, J Strassmeyer, A Walcroft, and Others. Temperature response of pa-  
417 rameters of a biochemically based model of photosynthesis. II. A review of experimental data.  
418 *Plant, Cell & Environment*, 25(9):1167–1179, 2002.

419 Belinda E Medlyn, Remko A Duursma, Derek Eamus, David S Ellsworth, I Colin Prentice, Craig V M  
420 Barton, Kristine Y Crous, Paolo de Angelis, Michael Freeman, and Lisa Wingate. Reconciling the  
421 optimal and empirical approaches to modelling stomatal conductance. *Global Change Biology*,  
422 17(6):2134–2144, 2011.

423 Belinda E Medlyn, Remko A Duursma, Derek Eamus, David S Ellsworth, I Colin Prentice, Craig V M  
424 Barton, Kristine Y Crous, Paolo Angelis, Michael Freeman, and Lisa Wingate. Reconciling the  
425 optimal and empirical approaches to modelling stomatal conductance. *Global Change Biology*,  
426 18(11):3476, 2012.

427 Patrick J Mitchell, Anthony P O’Grady, David T Tissue, Donald A White, Maria L Ottenschlaeger, and  
428 Elizabeth A Pinkard. Drought response strategies define the relative contributions of hydraulic  
429 dysfunction and carbohydrate depletion during tree mortality. *New Phytologist*, 197(3):862–872,  
430 2013.

431 Shinichi Nakagawa and Holger Schielzeth. A general and simple method for obtaining R<sup>2</sup> from  
432 generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 4(2):133–142, 2013.

433 D Scott NeSmith and John R Duval. The effect of container size. *HortTechnology*, 8(4):495–498,  
434 1998.

435 Matthew J Paul and Christine H Foyer. Sink regulation of photosynthesis. *Journal of experimental*  
436 *botany*, 52(360):1383–1400, 2001.

437 Jose Pinheiro, Douglas Bates, Saikat DebRoy, Deepayan Sarkar, and R Core Team. *{nlme}: Linear*  
438 *and Nonlinear Mixed Effects Models*, 2014. URL <http://cran.r-project.org/package=nlme>.

439 E A Pinkard, C L Beadle, N J Davidson, and M Battaglia. Photosynthetic responses of *Eucalyptus*  
440 *nitens* (Deane and Maiden) Maiden to green pruning. *Trees*, 12(3):119–129, 1998.

441 Hendrik Poorter, Ülo Niinemets, Lourens Poorter, Ian J Wright, and Rafael Villar. Causes and  
442 consequences of variation in leaf mass per area (LMA): a meta-analysis. *New Phytologist*, 182(3):  
443 565–588, 2009.

444 Hendrik Poorter, Jonas Bühler, Dagmar van Dusschoten, José Climent, and Johannes A Postma.  
445 Pot size matters: a meta-analysis of the effects of rooting volume on plant growth. *Functional*  
446 *Plant Biology*, 39(11):839–850, 2012a.

447 Hendrik Poorter, Karl J Niklas, Peter B Reich, Jacek Oleksyn, Pieter Poot, and Liesje Mommer.  
448 Biomass allocation to leaves, stems and roots: meta-analyses of interspecific variation and en-  
449 vironmental control. *New Phytologist*, 193(1):30–50, 2012b.

450 PB Reich, MB Walters, MG Tjoelker, D Vanderklein, and C Buschena. Photosynthesis and respira-  
451 tion rates depend on leaf and root morphology and nitrogen concentration in nine boreal tree  
452 species differing in relative growth rate. *Functional Ecology*, 12(3):395–405, 1998.

453 Peter B Reich, Y Weisel, A Eshel, and U Kafkafi. Root-shoot relations: optimality in acclimation and  
454 adaptation or the emperors new clothes. *Plant roots: the hidden half*, pages 205–220, 2002.

455 Cláudio P Ronchi, Fábio M DaMatta, Karine D Batista, Gustavo A B K Moraes, Marcelo E Loureiro,  
456 and Carlos Ducatti. Growth and photosynthetic down-regulation in *Coffea arabica* in response  
457 to restricted root volume. *Functional Plant Biology*, 33(11):1013–1023, 2006.

458 Michael G Ryan, Jose Luiz Stape, Dan Binkley, Sebastiao Fonseca, Rodolfo A Loos, Ernesto N Taka-  
459 hashi, Claudio R Silva, Sergio R Silva, Rodrigo E Hakamada, Jose Mario Ferreira, and Others.  
460 Factors controlling *Eucalyptus* productivity: How water availability and stand structure alter  
461 production and carbon allocation. *Forest ecology and management*, 259(9):1695–1703, 2010.

462 Rowan F Sage. Acclimation of photosynthesis to increasing atmospheric CO<sub>2</sub>: the gas exchange  
463 perspective. *Photosynthesis research*, 39(3):351–368, 1994.

464 Anna Sala, David R Woodruff, and Frederick C Meinzer. Carbon dynamics in trees: feast or famine?  
465 *Tree Physiology*, 32(6):764–775, 2012.

466 Arne Sellin. Does pre-dawn water potential reflect conditions of equilibrium in plant and soil water  
467 status? *Acta Oecologica*, 20(1):51–59, 1999.

468 B Shipley and D Meziane. The balanced-growth hypothesis and the allometry of leaf and root  
469 biomass allocation. *Functional Ecology*, 16(3):326–331, 2002.

470 M Stitt. Rising CO<sub>2</sub> levels and their potential significance for carbon flow in photosynthetic cells.  
471 *Plant, Cell & Environment*, 14(8):741–762, 1991.

472 Y Waisel, A Eshel, and U Kafkafi. Plant roots: the hidden half. *Marcel Dekker, Inc.: New York*, 1002:  
473 543–581, 1996.

474 David I Warton, Remko A Duursma, Daniel S Falster, and Sara Taskinen. smatr 3—an R package for  
475 estimation and inference about allometric lines. *Methods in Ecology and Evolution*, 3(2):257–259,  
476 2012.