

Effects of below-ground space limitation on performance of Eucalyptus seedlings: Does photosynthesis really control growth?

Courtney E. Company^{1,3}, Belinda E. Medlyn² and Remko A. Duursma¹

¹Hawkesbury Institute for the Environment, University of Western Sydney,
Richmond, NSW 2753, Australia

²Department of Biological Sciences, Macquarie University, North Ryde, NSW 2109,
Australia

³Corresponding author (c.company@uws.edu.au)

January 5, 2015

Abstract

Negative feedbacks to photosynthesis are commonly interpreted as the primary factor responsible for growth limitation in plants under stress. This study used manipulations of soil volume to test how growth is coupled to physiology, allocation, and sink activity in *Eucalyptus tereticornis* seedlings. We grew seedlings in a large range of container sizes and planted containers flush to the soil alongside freely rooted seedlings (free). Reduced soil volume was expected to induce rapid negative effects on growth and physiology compared to free seedlings. It was hypothesized that soil volume effect would be largest in the smallest containers, resulting in physical constraints to growth independently of photosynthesis. Photosynthesis would then become sink-limited, resulting in the build-up of leaf nonstructural carbohydrates eventually leading to photosynthetic down regulation. We observed a negative container effect on aboveground growth soon after the experiment started. Although growth was consistently different across soil volumes mass, partitioning to leaves, stems, roots was conserved after 120 days. Photosynthetic capacity was also significantly reduced in containers, and was related to both leaf nitrogen content and starch accumulation, however starch effects were weaker. We developed a simple seedling growth model that utilized leaf photosynthesis rates to allocate daily carbon (C) uptake towards mass growth of stems, leaves and roots. We then asked whether the observed reductions in photosynthesis explained the observed differences in seedling biomass. We found that photosynthesis reductions alone were not sufficient and the inclusion of additional carbon and mass allocation strategies were necessary to better predict the observed growth responses to decreasing soil volume. Overall, we found limited evidence for sink-limitation of photosynthesis by constraining seedling growth in containers, and argue that shifts in C allocation are necessary to explain the experimental results. This research highlights the need to further understand adaptive strategies of plant C allocation, and confirms that photosynthesis and growth are not always directly related.

Key Words

photosynthesis, sink regulation, growth, carbon allocation, soil volume

Introduction

Methods

Experimental Design

This experiment was located on the Hawkesbury Forest Experiment site in Richmond, NSW, Australia. Plots were located in open cover with a site history that consists of a paddock that was converted from native pasture grasses. Top soils at this site, used for the study, are an alluvial formation of low-fertility sandy loam soils (380 and 108 mg kg⁻¹ total nitrogen and phosphorus respectively) with low organic matter (0.7 %) and low water holding capacity. At this site a soil hard layer exists at ~1.0 m with a transition to heavy clay soils. The climate for the region is classified as sub-humid temperate.

Eucalyptus tereticornis seedlings, 20 weeks old and approximately 40 cm tall in tube stock, were chosen from a single local Cumberland plain cohort. Previous experiments have confirmed that species with tap roots (similar to *E. tereticornis*) use the center of the container as the medium for thick roots leaving the periphery of the soil as the most active sites for fine root proliferation (Biran and Eliassaf, 1980a,b). This is generally hypothesized to be a different response than seedlings with no taproot. By using a species with tap root growth and manipulations of container length rather than width, it is believed that a more realistic test of inhibition of growth through constrained soil volume would be achieved. Six seedlings were harvested before planting to measure initial leaf area and dry mass of leaves, stems and roots.

Six container volumes were used ranging from 5 l to 3 l, with a 22.5 cm diameter, and lengths ranging from 15 to 100 cm. Containers were constructed of PVC pipe and were filled with local top soil (described above). Soil in each container was packed to achieve a target soil bulk density of 1.7 g m⁻³. A Imidacloprid (BAYER CropScience) insecticide tablet was planted 5 cm below the roots for each seedling. Containers were planted flush with the soil surface inside metal sleeves, designed to minimize excess air space between the container and outside soil while also allowing for container removal. This allowed for soil temperatures in containers to reflect conditions of naturally sown seedlings. Each experimental block (n=7) contained a complete replicate set of container volumes as well as one free seedling, with 1 m² spacing. For each free seedling, a 1 m² subplot was excavated to 0.5 m and replaced with the same soil used in each container. Around each subplot a border of root exclusion material was buried 0.25 m deep and extended 0.25 m above the ground surface to exclude local vegetation.

Plants were watered bi-weekly or when needed, accounting for natural precipitation, to maintain soil moisture at field capacity (13-15 %). To achieve field capacity all soil filled containers were weighed and soil moisture was measured (Time Domain Reflectometer at 10 cm depth) before watering. Derived equations based on container weight at planting (when soil moisture was 3 %) and at field capacity was used to calculate watering requirements for each individual plant. Drain systems were built into each pot to prevent pooling of water in containers before root expansion, from reduced root uptake, or from large rainfall events. These conditions could lead to an anaerobic environment around the root that could hinder the uptake of water through reduced root conductance (Poorter et al., 2009), an undesired experimental artifact. A collection compartment in the bottom of containers, containing gravel covered by root exclusion mesh, was used collect water for 20, 25, and 35 l containers. Plastic tubing (6 mm diameter) was inset into the gravel layer and extended through the top of the container. A lysimeter pump was then used to suction excess water, through the tubing, as needed. As small containers (5, 10, and 15 l) have a larger irradiation effect a simple bottom plug was used to drain water from the gravel compartment.

Growth and morphology metrics

Seedlings were planted on January 21st 2013 and stem height, diameter at 15 cm and leaf count were measured weekly thereafter. Once the growth rate of individual plants had significantly declined a full biomass harvest was completed (May 5th 2013). Dry mass of leaves, stems, roots and cumulative leaf area (LI-3100C Area Meter; LI-COR, Lincoln, NE, USA) was measured for each seedling. Mean individual leaf area for each harvested seedling was calculated by dividing cumulative leaf area by total leaf count of only fully expanded leaves. This value was then used to interpolate cumulative leaf area through time with weekly leaf counts. Root mass was collected by passing soil from each container through a 1 mm sieve, washing, separating into fine and coarse roots (<2 mm and >2 mm diameter, respectively) and then drying to a constant mass. Roots from the free seedlings were collected by excavating each 1 m² subplot to 0.5 m depth. 25 g fresh weight subsamples of washed fine roots were analyzed, using WhinoRhizo software (Regent Instruments Inc.), for specific root length (SRL, cm m⁻¹).

Photosynthetic parameters

Leaf gas exchange measurements were performed bi-weekly at saturating light (A_{sat}) and saturating light and $[\text{CO}_2]$ (A_{max}) on new fully expanded leaves. Measurements were initiated only after new leaf growth occurred (March 17th, 2013), approximately 6 weeks following planting, and continued until the biomass harvest. Leaf level gas exchange was measured with a standard leaf chamber equipped with blue-red light emitting diodes using a portable gas exchange system (LI-6400, LI-COR, Lincoln, NE, USA). A_{sat} measurements were made at PPFD of 1800 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and $[\text{CO}_2]$ of 400 $\mu\text{l l}^{-1}$ and A_{max} with $[\text{CO}_2]$ of 1600 $\mu\text{l l}^{-1}$ and PPFD of 1800 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. This choice of light level to achieve light saturation is consistent with other studies on *Eucalyptus* species (Kallarackal and Somen, 1997; Pinkard et al., 1998; Crous et al., 2013; Drake et al., 2014). These measurements were conducted during midday (10:00-14:00 h) with leaf temperature maintained at 25°C. After leaves acclimated to the chamber environment, net CO_2 assimilation rate and stomatal conductance (g_s) were logged 5 times for both A_{sat} and A_{max} . Photosynthetic CO_2 response (AC_i) curves were also developed at 25°C on a random subset of each container size ($n=3$) after new leaves were first produced and immediately prior to the final harvest (May 23rd 2013). Each AC_i curve began at the reference $[\text{CO}_2]$ of 400 $\mu\text{l l}^{-1}$ and then consisted of 12 additional steps from $[\text{CO}_2]$ of 50 to 1800 $\mu\text{l l}^{-1}$ at 25°C and saturating light (above). From these curves the photosynthetic parameters, J_{max} and V_{cmax} , can be quantified using biochemical model of Farquhar et al. (1980). Leaf dark respiration rates (R_d) was measured on each seedling during the same dates as AC_i curves using detached leaves inside a conifer chamber attached to the Licor 6400 at least 1 hour after sundown. Measurements were taken at a reference $[\text{CO}_2]$ of 400 $\mu\text{l l}^{-1}$ while leaf temperature was maintained at current ambient conditions. Specific leaf area (SLA, m² kg⁻¹) was calculated by measuring leaf area and dry mass for individual leaves sampled during gas exchange campaigns.

Leaf water potential

Predawn (Ψ_{pd}) and midday (Ψ_l) leaf water potentials were measured for each seedling using a PMS 1505D pressure chamber (PMS Instruments, Albany, OR, USA) on fully expanded leaves during the same time period as AC_i and R_d . Leaves were collected, immediately stored inside foil covered bags and then transported to the laboratory for measurements of water potential. Ψ_{pd} was measured before sunrise and Ψ_l at midday 13:00-14:30 h. These measurements were used as a measure of static water stress on the seedlings (Sellin, 1999), and to ensure that the bulk soil water availability was high enough for plants as they became larger and roots filled the soil volume.

Leaf, root and soil chemistry

Leaves used in each gas exchange measurements and subsamples of harvested roots were dried to a constant mass and milled for analysis of *N* content. Pre-planting soil samples (*n*=6) and subsamples of soil from each container following harvest were sieved to remove organic material, air dried and milled for analysis. *N* concentrations of milled samples were determined using a Carlo Erba CE1110 elemental analyzer with thermal conductivity and mass spectrometric detection (of *N*₂ and CO₂). The percentage of *N* in the sample was calculated by comparison with known standards. Leaf total non-structural carbohydrates concentration was analyzed using a total starch assay kit (Megazyme International 303 Ireland Ltd., Wicklow, Ireland) on gas exchange leaves (above) and represents the sum of starch (mg g⁻¹) and soluble sugar (mg g⁻¹) concentrations. Starch was quantified using a thermostable α -amylase and amyloglucosidase assay (McCleary et al., 1997) and soluble sugars were determined following the anthrone method (Ebell, 1969). Full methods of the TNC assay are described in (Mitchell et al., 2013).

Seedling growth model

We developed a simple seedling growth model that utilized leaf photosynthesis to allocate daily carbon assimilate towards mass production of stems, leaves and roots. The model begins with the mean initial component mass (*leaf*_{*i*}, *stem*_{*i*} and *root*_{*i*}) and a starting leaf area (*LA*_{*i*}) measured prior to planting. Biomass production for each day of the experiment was estimated by multiplying the whole tree seedling carbon uptake gain per day (*C*_{day}) by a biomass conversion efficiency parameter (ξ_c , (Zhu et al., 2008)), allocating new carbon to the previous days biomass and then accounting for carbon loss via tissue respiration. *C*_{day} was calculated for each seedling by fitting a coupled photosynthesis - stomatal conductance model (Farquhar et al., 1980; Medlyn et al., 2011) in the 'plantecophys' package in R (Duursma, 2014) to the mean photosynthetic parameters (*R*_d, *J*_{max}, *V*_{cmax}, and *g*_i) for each treatment and meteorological data from an onsite weather station. Examples of the photosynthesis model are described in Medlyn et al. (2002) and the approach of the coupled leaf gas exchange model are described in Duursma et al. (2014). The *g*_i parameter was generated by fitting observed stomatal conductance values into the optimal stomatal conductance model from (Medlyn et al., 2012). Combined with the meteorological parameters PPFD, air temperature, and relative humidity, at 15 m intervals, photosynthesis rates ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) were then predicted for each soil volume treatment. Rates were assumed to be representative of the entire 15 min meteorological interval. *C*_{day} was then calculated by converting predicted rates to mass *C* gain over 15 min (g m⁻²) and then summed for 24 h. Additionally, it is necessary to calculate a self-shading parameter (*M*) when scaling flat leaf photosynthesis to seedling cumulative leaf area. This was accomplished by utilizing 61 previously digitized Eucalyptus seedlings, covering 5 total species which include *E. tereticornis*, from Duursma et al. (2012b) to run in 'YplantQMC' package in R (Duursma et al., 2012a) to build a 3d plant structure based on digitized metrics of plant allometry and crown structure. Inputting the same physiological parameters listed above, 'YplantQMC' outputs total photosynthesis, using total leaf area, for seedlings assuming self-shading as well as for a full sun large horizontal leaf. The ratio of self-shading to flat leaf total photosynthesis was then used to calculate *M*, which was then applied to modeled rates of photosynthesis when scaling to seedling cumulative leaf area. All default parameters used in model simulations are reported in Table. 3.

We then utilized this model to investigate whether the effects of soil volume on photosynthesis were enough to accurately predict overall seedling production after 120 days or whether changes in *C* allocation or mass partitioning were also necessary to predict biomass. First, the model was simulated using only changes in photosynthesis rates across treatments combined with mean values of mass partitioning and either published or local data of stem and root respiration rates. Next, different mass partitioning and *C* allocation scenarios were simulated to investigate sources of missing *C* from the initial model simulation. This included testing model sensitivity by altering leaf *C* allocation, root respiration rates, and fine root *C* allocation. For all cases, mass production was compared between model output and harvested seedlings to investigate what *C* allocation

strategies beyond reduction in photosynthesis aid in prediction of seedling growth responses induced by limiting soil volume.

Data Analysis

Differences in experimental parameters with soil volume were analysed by one-way analysis of variance (ANOVA) in R with individual containers as random effects and soil volume as a categorical fixed effect. Tukeys post-hoc tests were performed in conjunction with ANOVA to determine which specific paired comparisons among soil volumes were different. Mixed model ANOVAs of A_{\max} and leaf chemistry were performed using the 'nlme' package (Pinheiro et al., 2014) in R and r^2 values of mixed models were computed as in Nakagawa and Schielzeth (2013). Tests of allometric relationships between biomass components were implemented using major axis regression in the 'smatr' package (Warton et al., 2012).

Results

Growth and morphology metrics

In this open field study, colder temperatures and reductions in cumulative PPFD per day (Fig. 1) most likely lead to the reduced growth in the free seedlings in the final weeks of the experiment (Fig. 2). Combined with severe growth reductions in the smallest container volumes the experiment was chosen to be harvested after 120 days. Over the 120 days height, diameter, and leaf area diverged between container volumes (Fig. 2). First, seedling leaf area significantly diverged between soil volumes ($P < 0.026$) during the 5th week of the experiment. Following this period both height (8th week) and then diameter (9th week) significantly deviated across soil volumes ($P < 0.002$ & 0.001 , respectively). Negative growth effects then manifested as severely reduced height gain and declining leaf area through time with small container volumes across the final two months of the experiment. Seedlings maintained diameter growth throughout the experiment, although marginal with smaller soil volumes in the final month. Final seedling height significantly increased with increasing soil volume ($P < 0.001$). Increases in both final stem diameter ($P < 0.001$) and cumulative leaf area (both $P < 0.001$) were found with increasing soil volume and these differences were driven mainly by the largest container and the free seedling.

Total plant biomass at harvest was significantly different across container volumes ($P < 0.001$) and with free seedlings ($P < 0.001$, Table. 1). We analyzed the relationship between biomass growth with each fold increase in soil volume and found an increase of 34 % with a doubling of pot size, consistent with the meta-analysis of Poorter et al. (2012). Additionally, plant biomass was highly correlated with total leaf area across all treatments ($r^2 = .97$, $P < 0.001$). Differences in biomass partitioning to leaves, stems, and roots were not different across soil volumes when variation in seedling biomass within treatments was factored in the analysis. Across all treatments, the final harvested root:shoot was conserved in these seedlings, with a slightly higher shoot than root mass on average ($\bar{x} = 0.904$).

SRL of harvested fine roots was not different across soil volumes (Table. 1). Over the duration of the experiment SLA was higher in free seedlings but was not different across containers sizes (Table. 1, $P < 0.001$) and this pattern was evident in the first gas exchange measurement campaign (6th week, $P < 0.001$).

Leaf and root chemistry

Leaf N % was significantly higher in free seedlings and the largest container volume at the onset of gas exchange measurements (6th week, $P < 0.001$). Over the remaining duration of the experiment

the smallest container volume had a significant reduction in leaf N % compared to other soil volumes, while the free seedling maintained a greater leaf N % (Table. 1, $P < 0.001$). Additionally, mean leaf starch content in the smallest container was double that of free seedlings ($P = 0.042$), while leaf soluble sugars did not differ across treatments throughout the experiment (Table. 1). Differences in leaf starch between the free seedling and the smallest container were also evident during the first gas exchange campaign ($P = 0.0013$).

Gas exchange and photosynthetic parameters

A_{sat} (Fig. 3) and A_{max} (Table. 2) were both significantly higher in the largest container volume and the free seedling at the first measurement campaign (both $P < 0.001$). Across measurement campaigns A_{sat} was consistently higher in free seedlings than in containers (Figure 3, $P < 0.001$). The interaction between photosynthetic capacity on a mass basis, leaf starch, and foliar N was marginally significant ($P = 0.0584$) but A_{max} was highly correlated to both leaf N content and leaf starch (both $P < 0.001$). Across all measurement campaign A_{max} was higher when foliar N was also higher, usually associated with low levels of leaf starch (Fig. 3a). A_{max} was also lower when leaf starch was high as higher leaf N often did not coincide with high leaf starch (Fig. 3b)

The photosynthetic parameters J_{max} and V_{Cmax} were not different across measurement campaigns, therefore the parameter means per volume are reported here (Table. 2). Overall, both J_{max} and V_{Cmax} were significantly higher in free seedlings with little variation between container volumes ($P = 0.0012$ & 0.0021 , respectively). Leaf dark respiration rates were not significantly different across soil volumes (Table. 2). The g_1 parameter, generated for each seedling from the Medlyn et al. (2012) optimal stomatal conductance model, was not different across soil volumes (Table. 2). Predicted values of g_s , using the g_1 parameter, were highly correlated with observed values ($r^2 = .74$, $P < 0.001$, data not shown).

Neither Ψ_{pd} nor Ψ_l were different across treatments, with mean values of -0.27 and -1.2 mPa across all seedlings, respectively. Although stomatal conductance in free seedlings was generally higher than those in containers (Table. 2, $P = 0.0023$), the mean rates for all seedlings were high at $0.37 \text{ mol H}_2\text{O m}^{-2} \text{ s}^{-1}$ and did not decline significantly across the experiment duration. Combined these indices provide strong evidence that water stress was not apparent on these well-watered seedlings throughout the experiment. Soil N % at harvest was not different across soil volumes ($\bar{x} = 0.45$ %) and decreased approximately 3 % across all containers over the experiment duration. This indicates that nutrient leaching from free seedlings or from draining of containers following natural rainfall events did not differ between treatments.

Modelling seedling biomass

The initial model simulation, testing only treatment specific photosynthesis rates, was unable to accurately predict the responses of actual harvested seedling biomass to reduced soil volume. When both C_{day} and biomass of modeled and harvested seedlings were scaled to to the 'free' seedling the initial model design overestimated biomass for all treatments (Fig. 6a). This suggests that reductions in photosynthesis alone could not realistically capture the reduced growth response of these seedlings. Next, the model was re-run using the mean mass fractions of leaves, stems and roots from the final harvest, per treatment, to drive daily mass partitioning and seedling production. Combined with treatment specific photosynthesis rates this model still overestimated seedling production for all treatments (Fig. 6b).

To test whether potential sources of missing C could improve the modeled response of seedling mass production the model was run with 3 different C allocation scenarios that could not be tested within the framework of this experiment. This was accomplished by testing the sensitivity of the model to adjustments of C allocation to leaves or fine roots or root respiration rates by ± 50 % of default values. The model output of these scenarios was then compared to both the original model output and with harvested seedling biomass. Increasing C allocation to either leaves or fine

roots, in coalescence with photosynthesis rates, did improve the model but neither were sufficient to accurately predict the actual biomass response (Fig. S1a,c). Altering respiration rates of fine and coarse roots did not have a noticeable deviation from the original model (Fig. S1b).

Discussion

Tables

Table 1: Responses of plant and leaf characteristics of *Eucalyptus tereticornis* seedlings to soil volume treatments. Each value reflects the mean(standard error) of each treatment.

Volume (L)	Seedling mass (g)	SLA (m ² kg ⁻¹)	Leaf Nitrogen (%)	Leaf Sugars (%)	Leaf Starch (%)	SRL (cm m ⁻¹)	Root Nitrogen (%)
5	14.8 (1.82)	9.5 (0.23)	1.1 (0.02)	6.4 (0.28)	12.7 (0.97)	39.1 (5.47)	0.82 (0.05)
10	20.0 (2.38)	9.8 (0.24)	1.3 (0.04)	6.7 (0.25)	9.4 (0.75)	34.2 (5.83)	0.75 (0.02)
15	25.4 (2.49)	11.0 (0.47)	1.4 (0.06)	7.2 (0.28)	7.3 (0.73)	37.6 (4.63)	0.71 (0.02)
20	23.4 (1.63)	9.8 (0.28)	1.4 (0.05)	6.6 (0.26)	9.5 (0.88)	45.3 (5.50)	0.76 (0.04)
25	30.4 (5.49)	10.4 (0.37)	1.3 (0.06)	6.9 (0.24)	9.8 (0.71)	47.0 (7.10)	0.74 (0.02)
35	52.2 (9.55)	11.3 (0.44)	1.5 (0.08)	6.8 (0.22)	9.8 (0.65)	50.6 (11.61)	0.77 (0.03)
Free	174.5 (18.02)	13.0 (0.44)	2.4 (0.09)	7.4 (0.25)	6.8 (0.65)	43.7 (6.24)	0.87 (0.04)

Table 2: Responses of leaf level gas exchange parameters of *Eucalyptus tereticornis* seedlings to soil volume treatments. Each value reflects the mean(standard error) of each treatment. Units for A_{\max} and R_{dark} are $\mu\text{mol m}^{-1} \text{s}^{-1}$ and g_s are $\text{mol m}^{-1} \text{s}^{-1}$.

Volume (L)	A_{\max}	R_{dark}	J_{\max}	V_{Cmax}	g_s	g_1
5	21.2 (0.9)	2.8 (0.3)	103.7 (5.8)	62.9 (2.8)	0.30 (0.01)	5.1 (0.1)
10	22.3 (1.4)	2.7 (0.4)	116.5 (5.9)	69.4 (2.9)	0.36 (0.01)	5.4 (0.1)
15	23.3 (1.2)	1.4 (0.1)	123.1 (15.4)	79.4 (10.0)	0.45 (0.01)	6.2 (0.2)
20	26.1 (0.7)	1.7 (0.1)	130.0 (14.1)	81.4 (6.0)	0.38 (0.01)	5.2 (0.2)
25	23.9 (0.9)	1.2 (0.1)	131.9 (5.9)	78.6 (2.6)	0.32 (0.01)	4.8 (0.2)
35	25.0 (1.0)	1.5 (0.2)	121.2 (3.7)	78.0 (3.5)	0.33 (0.02)	4.7 (0.2)
Free	33.1 (0.7)	1.3 (0.1)	171.7 (18.6)	101.2 (5.7)	0.49 (0.02)	5.0 (0.2)

Figures




Figure 1: Daily maximum and minimum temperature (a), cumulative daily PPFD (b), and daily maximum vapour pressure deficit (c) across the experiment duration.




Figure 2: Soil volume treatment means \pm SE (n=7) of height growth (a), diameter growth (b), and interpolated seedling leaf area (c) measured weekly of *Eucalyptus tereticornis* seedlings over 120 days.




Figure 3: Soil volume treatment means \pm SE (n=7), across all measurement dates (n=6), of light saturated rates of photosynthesis at 25°C




Figure 4: Soil volume treatment means (n=7) of mass partitioning to leaves, stems, and roots (a) and bi-variate relationships between mass allocation to leaves and stems + roots (b). Lines represent standardized major axis fitting of the log transformed allometric relationships of leaf mass fraction by treatment.




Figure 5: Photosynthetic capacity, on a leaf mass basis, as a function of accumulation of leaf starch (a) and leaf nitrogen content without TNC (b). Colors represent bins levels ($n=5$) of both leaf starch and nitrogen grouped from low to high. Lines represent predictions, for each bin level, from the linear mixed effects model equation of A_{max} as a function of starch and nitrogen. The marginal r^2 (fixed effects only) was 0.37 and the conditional r^2 (fixed and random effects) was 0.48 for the complete model.




Figure 6: Modeled and harvested seedling biomass (g) versus daily assimilated carbon gain (g m^{-2}). Values of biomass and carbon gain are scaled to the free seedling with unlimited soil volume.

1 Supporting Information Figures

Table 3: Seedling Growth Model Default Parameters

Variable	Default Value	Units	Source
Leaf area _i	0.035	m ²	this study
Leaf mass _i	3.45	g	this study
Stem mass _i	1.51	g	this study
Root mass _i	0.99	g	this study
ξ _c	.65		Zhu et al. (2008)
R _{coarse root}	0.00124	g C g root ⁻¹	Marsden et al. (2008)
R _{fine root}	0.010368	g C g root ⁻¹	Ryan et al. (2010)
R _{stem}	0.00187	g C g stem ⁻¹	Drake 2014 (unpublished)
C _{day}	4.7-6.8	g C m ²	this study
M	0.80-0.87		this study




Figure S1: Modeled and harvested seedling biomass (g) versus daily assimilated carbon gain (g m²) including sensitivity testing for unmeasured carbon allocation scenarios. Model parameters of leaf allocation (a), root respiration (b), and fine root allocation (c) were increased or decreased by 50%.

References

- I Biran and A Eliassaf. The effect of container size and aeration conditions on growth of roots and canopy of woody plants. *Scientia Horticulturae*, 12(4):385–394, 1980a.
- I Biran and A Eliassaf. The effect of container shape on the development of roots and canopy of woody plants. *Scientia Horticulturae*, 12(2):183–193, 1980b.
- Kristine Y Crous, Audrey G Quentin, Yan-Shih Lin, Belinda E Medlyn, David G Williams, Craig V M Barton, and David S Ellsworth. Photosynthesis of temperate *Eucalyptus globulus* trees outside their native range has limited adjustment to elevated CO₂ and climate warming. *Global change biology*, 19(12):3790–3807, 2013.
- John E Drake, Michael J Aspinwall, Sebastian Pfautsch, Paul D Rymer, Peter B Reich, Renee A Smith, Kristine Y Crous, David T Tissue, Oula Ghannoum, and Mark G Tjoelker. The capacity to cope with climate warming declines from temperate to tropical latitudes in two widely distributed *Eucalyptus* species. *Global change biology*, 2014.
- Remko Duursma. *plantecophys: Modelling and analysis of leaf gas exchange data*, 2014.
- Remko Duursma, Mik Cieslak, Robert Pearcy, and Belinda Medlyn. *YplantQMC: Plant architectural analysis with Yplant and QuasiMC.*, 2012a. URL <http://www.remkoduursma.com/yplantqmc>, <https://www.bitbucket.org/remkoduursma/yplantqmc/>.
- Remko A Duursma, Daniel S Falster, Fernando Valladares, F J Sterck, R W Pearcy, Christopher H Lusk, K M Sendall, Marisa Nordenstahl, N C Houter, B J Atwell, and Others. Light interception efficiency explained by two simple variables: a test using a diversity of small-to medium-sized woody plants. *New Phytologist*, 193(2):397–408, 2012b.
- Remko A Duursma, Craig V M Barton, Yan-Shih Lin, Belinda E Medlyn, Derek Eamus, David T Tissue, David S Ellsworth, and Ross E McMurtrie. The peaked response of transpiration rate to vapour pressure deficit in field conditions can be explained by the temperature optimum of photosynthesis. *Agricultural and Forest Meteorology*, 189:2–10, 2014.
- L F Ebell. Variation in total soluble sugars of conifer tissues with method of analysis. *Phytochemistry*, 8(1):227–233, 1969.
- G D Farquhar, S von von Caemmerer, and J A Berry. A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta*, 149(1):78–90, 1980.
- J Kallarackal and C K Somen. An ecophysiological evaluation of the suitability of *Eucalyptus grandis* for planting in the tropics. *Forest Ecology and Management*, 95(1):53–61, 1997.
- Claire Marsden, Yann Nouvellon, and Daniel Epron. Relating coarse root respiration to root diameter in clonal *Eucalyptus* stands in the Republic of the Congo. *Tree physiology*, 28(8):1245–1254, 2008.
- B. V. McCleary, T. S. Gibson, and D. C. Mugford. Measurement of total starch in cereal products by amyloglucosidase- α -amylase method: Collaborative study. *Journal of AOAC International*, 80(3): 571–579, 1997. ISSN 1060-3271.
- B E Medlyn, Erwin Dreyer, D Ellsworth, M Forstreuter, P C Harley, M U F Kirschbaum, Xavier Le Roux, Pierre Montpied, J Strassemeyer, A Walcroft, and Others. Temperature response of parameters of a biochemically based model of photosynthesis. II. A review of experimental data. *Plant, Cell & Environment*, 25(9):1167–1179, 2002.
- Belinda E Medlyn, Remko A Duursma, Derek Eamus, David S Ellsworth, I Colin Prentice, Craig V M Barton, Kristine Y Crous, Paolo de Angelis, Michael Freeman, and Lisa Wingate. Reconciling the optimal and empirical approaches to modelling stomatal conductance. *Global Change Biology*, 17(6):2134–2144, 2011.
- Belinda E Medlyn, Remko A Duursma, Derek Eamus, David S Ellsworth, I Colin Prentice, Craig V M Barton, Kristine Y Crous, Paolo Angelis, Michael Freeman, and Lisa Wingate. Reconciling the

- optimal and empirical approaches to modelling stomatal conductance. *Global Change Biology*, 18(11):3476, 2012.
- Patrick J Mitchell, Anthony P O'Grady, David T Tissue, Donald A White, Maria L Ottenschlaeger, and Elizabeth A Pinkard. Drought response strategies define the relative contributions of hydraulic dysfunction and carbohydrate depletion during tree mortality. *New Phytologist*, 197(3):862–872, 2013.
- Shinichi Nakagawa and Holger Schielzeth. A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 4(2):133–142, 2013.
- Jose Pinheiro, Douglas Bates, Saikat DebRoy, Deepayan Sarkar, and R Core Team. *{nlme}: Linear and Nonlinear Mixed Effects Models*, 2014. URL <http://cran.r-project.org/package=nlme>.
- E A Pinkard, C L Beadle, N J Davidson, and M Battaglia. Photosynthetic responses of Eucalyptus nitens (Deane and Maiden) Maiden to green pruning. *Trees*, 12(3):119–129, 1998.
- Hendrik Poorter, Ülo Niinemets, Lourens Poorter, Ian J Wright, and Rafael Villar. Causes and consequences of variation in leaf mass per area (LMA): a meta-analysis. *New Phytologist*, 182(3): 565–588, 2009.
- Hendrik Poorter, Jonas Bühler, Dagmar van Dusschoten, José Climent, and Johannes A Postma. Pot size matters: a meta-analysis of the effects of rooting volume on plant growth. *Functional Plant Biology*, 39(11):839–850, 2012.
- Michael G Ryan, Jose Luiz Stape, Dan Binkley, Sebastiao Fonseca, Rodolfo A Loos, Ernesto N Takahashi, Claudio R Silva, Sergio R Silva, Rodrigo E Hakamada, Jose Mario Ferreira, and Others. Factors controlling Eucalyptus productivity: How water availability and stand structure alter production and carbon allocation. *Forest ecology and management*, 259(9):1695–1703, 2010.
- Arne Sellin. Does pre-dawn water potential reflect conditions of equilibrium in plant and soil water status? *Acta Oecologica*, 20(1):51–59, 1999.
- David I Warton, Remko A Duursma, Daniel S Falster, and Sara Taskinen. smatr 3—an R package for estimation and inference about allometric lines. *Methods in Ecology and Evolution*, 3(2):257–259, 2012.
- Xin-Guang Zhu, Stephen P Long, and Donald R Ort. What is the maximum efficiency with which photosynthesis can convert solar energy into biomass? *Current opinion in biotechnology*, 19(2): 153–159, 2008.