

Automatic construction and description of nonparametric models

James Robert Lloyd¹, David Duvenaud¹, Roger Grosse², Joshua B. Tenenbaum², Zoubin Ghahramani¹

Massachusetts **Institute of Technology**



1: Department of Engineering, University of Cambridge, UK 2: Massachusetts Institute of Technology, USA

This analysis was automatically generated





Figure 1: Raw data (left) and model posterior with extrapolation (right)

The Automatic Bayesian Covariance Discovery system has discovered four additive components in


- A linearly increasing function.
- An approximately periodic function with a period of 1.0 years and with linearly increasing amplitude.
- A smooth function.
- Uncorrelated noise with linearly increasing standard deviation.

Component 1: A linearly increasing function This component is linearly increasing.




Component 2: An approximately periodic function with a period of 1.0 years and with linearly increasing amplitude

This component is approximately periodic with a period of 1.0 years and varying amplitude. Across periods the shape of this function varies very smoothly. The amplitude of the function increases linearly. The shape of this function within each period has a typical lengthscale of 6.0 weeks.




2.3 Component 3 : A smooth function




Modelling structure through Gaussian process kernels


• The kernel specifies which structures are likely under the GP prior - which determines the generalisation properties of the model.



Composite kernels can express many types of structure



We build models by a greedy search



Automatically describing model properties


Kernels can be distributed into a sum of products

$$SE \times (LIN + PER + SE)$$

becomes (after simplification)

$$(SE \times LIN) + (SE \times PER) + (SE).$$

Sums of kernels correspond to sums of functions



If $f_1(x) \sim \text{GP}(0, k_1)$ and $f_2(x) \sim \text{GP}(0, k_2)$ then $f_1(x) + f_2(x) \sim \text{GP}(0, k_1 + k_2)$. Therefore, a sum of kernels can be described as a sum of functions.

The compositional structure of products of kernels maps onto compositionally constructed sentences

Kernel	Noun phrase	Postmodifier phrase
WN	uncorrelated noise	n/a
\mathbf{C}	constant	n/a
SE	smooth function	whose shape changes smoothly
Per	periodic function	modulated by a periodic function
LIN	linear function	with linearly varying amplitude
$\prod_k \operatorname{Lin}^{(k)}$	polynomial	with polynomially varying amplitude
$\prod_{k}^{\infty} oldsymbol{\sigma}^{(k)}$	n/a	which applies until / from [changepoint]

Example description

periodic function with linearly growing amplitude which applies until [date]

 $ext{Per}$ has been chosen to act as the noun while $ext{Lin}$ and $oldsymbol{\sigma}$ modify the description

Visit the website - try the (simple) demo

www.automaticstatistician.com