

# All of Statistics

My proposed solutions to the book:

**All of Statistics - A Concise Course in Statistical Inference.**

I really needed a proper refresher on statistics.

Let me know if you find any mistakes! I am sure there are plenty. :(

## Contents

<b>1</b>	<b>Chapter 1 - Probability</b>	<b>2</b>
<b>2</b>	<b>Chapter 2 - Random Variables</b>	<b>25</b>
<b>3</b>	<b>Chapter 3 - Expectation</b>	<b>49</b>

# 1 Chapter 1 - Probability

## Exercises

### 1.1

Proving the Continuity of Probabilities.

**1.8 Theorem** (Continuity of Probabilities). If  $A_n \rightarrow A$  then

$$\mathbb{P}(A_n) \rightarrow \mathbb{P}(A)$$

as  $n \rightarrow \infty$ .

PROOF. Suppose that  $A_n$  is monotone increasing so that  $A_1 \subset A_2 \subset \dots$ . Let  $A = \lim_{n \rightarrow \infty} A_n = \bigcup_{i=1}^{\infty} A_i$ . Define:

$$\begin{aligned} B_1 &= A_1 \\ B_2 &= \{\omega \in \Omega : \omega \in A_2, \omega \notin A_1\} \\ B_3 &= \{\omega \in \Omega : \omega \in A_3, \omega \notin A_2, \omega \notin A_1\} \\ &\vdots \end{aligned}$$

(1) Showing that  $B_i$  are disjoint sets.

We have  $B_1 = A_1$ . Since  $B_2 = \{\omega \in \Omega : \omega \in A_2, \omega \notin A_1\}$ , we can rewrite this as  $B_2 = A_2 - A_1$  by definition of set difference.

Since  $B_3 = \{\omega \in \Omega : \omega \in A_3, \omega \notin A_2, \omega \notin A_1\}$  and since  $A_1 \subset A_2$ , we can rewrite this as  $B_3 = A_3 - (A_1 \cup A_2)$ . In general;  $B_k = A_k - (\bigcup_{i=1}^{k-1} A_i)$ .

Assuming some random sets  $B_m, B_p$  for some  $m, p \in \mathbb{N}$ . Without loss of generality, we assume  $m > p$ . Then:

$$\begin{aligned} B_m \cap B_p &= \left( A_m - \left( \bigcup_{i=1}^{m-1} A_i \right) \right) \cap \left( A_p - \left( \bigcup_{i=1}^{p-1} A_i \right) \right) \\ &= \left( A_m \cap \left( \bigcup_{i=1}^{m-1} A_i \right)^c \right) \cap \left( A_p \cap \left( \bigcup_{i=1}^{p-1} A_i \right)^c \right) \end{aligned}$$

DeMorgan's law

$$= \left( A_m \cap A_1^c \cap A_2^c \cap \dots \cap A_p^c \cap \dots \cap A_{m-1}^c \right) \cap \left( A_p \cap A_1^c \cap A_2^c \cap \dots \cap A_{p-1}^c \right)$$

Reshuffling terms and repeated use of  $C \cap C = C$

$$\begin{aligned} &= A_m \cap A_1^c \cap A_2^c \cap \dots \cap \underbrace{A_p^c \cap A_p}_{=\emptyset} \cap \dots \cap A_{m-1}^c \\ &= \emptyset \end{aligned}$$

Since  $B_m \cap B_p = \emptyset$ , they are disjoint. (*Quite certain this is a correct argument, but could have made it a bit easier by going directly to the  $A_3 \cap A_2^c \cap A_1^c$  version of the sets.*)

(2) Showing that

$$A_n = \bigcup_{i=1}^n A_i = \bigcup_{i=1}^n B_i$$

for each  $n$ . Showing this with an induction argument. Note that  $A_1 = B_1$ , and  $A_2 \cup A_1 = A_2$  since  $A_1 \subset A_2$ . Note also that:

$$\begin{aligned} B_2 \cup B_1 &= (A_2 \cap A_1^c) \cup A_1 \\ &= (A_2 \cup A_1) \cap (A_1^c \cup A_1) \\ &= (A_2 \cup A_1) \cap (\Omega) \\ &= A_2 \cup A_1 \\ &= A_2 \end{aligned}$$

So this is true for  $n = 2$ . Assume that  $A_k = \bigcup_{i=1}^k A_i = \bigcup_{i=1}^k B_i$  for  $k \in \mathbb{N}$ . Showing that this applies to  $k + 1$ .

$$\begin{aligned} \bigcup_{i=1}^{k+1} A_i &= A_{k+1} \cup \left( \bigcup_{i=1}^k A_i \right) = A_{k+1} \cup A_k = A_{k+1} \\ \bigcup_{i=1}^{k+1} B_i &= B_{k+1} \cup \left( \bigcup_{i=1}^k B_i \right) \\ &= B_{k+1} \cup A_k \\ &= \left[ A_{k+1} \cap \left( A_k^c \cap \dots \cap A_1^c \right) \right] \cup A_k \\ &= \left[ A_{k+1} \cap \left( A_k \cup \dots \cup A_1 \right)^c \right] \cup A_k \\ &= \left[ A_{k+1} \cap A_k^c \right] \cup A_k \\ &= (A_{k+1} \cup A_k) \cap (A_k^c \cup A_k) \\ &= (A_{k+1} \cup A_k) \cap \Omega \\ &= A_{k+1} \cup A_k \\ &= A_{k+1} \end{aligned}$$

Result verified by induction argument.

(3) Showing that

$$\bigcup_{i=1}^{\infty} B_i = \bigcup_{i=1}^{\infty} A_i. \quad (\diamond)$$

By definition:

$$A = \lim_{n \rightarrow \infty} A_n = \bigcup_{i=1}^{\infty} A_i.$$

By step (2):

$$A = \lim_{n \rightarrow \infty} A_n = \lim_{n \rightarrow \infty} \bigcup_{i=1}^n A_i = \lim_{n \rightarrow \infty} \bigcup_{i=1}^n B_i = \bigcup_{i=1}^{\infty} B_i.$$

Hence,  $(\diamond)$  is satisfied.

## 1.2

Proving some well known results by using the axioms:

$$\mathbb{P}(A) \geq 0, \forall A \subset \Omega \quad (\text{A.1})$$

$$\mathbb{P}(\Omega) = 1, \quad (\text{A.2})$$

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i) \quad (\text{A.3})$$

where  $A_1, A_2, \dots$  are disjoint in (A.3).

• **Claim:**  $\mathbb{P}(\emptyset) = 0$ .

PROOF. Since  $\Omega \cap \emptyset = \emptyset$  and the empty set is disjoint with itself, we can make set  $E_1 = \Omega$  and  $E_k = \emptyset$  for all  $k \geq 2$ . Now assume for contradiction that  $\mathbb{P}(\emptyset) = a > 0$  for some  $a \in \mathbb{R}$ . Then by (A.3):

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} E_i\right) \stackrel{\text{A.3}}{=} \sum_{i=1}^{\infty} \mathbb{P}(E_i) = \mathbb{P}(\Omega) + \sum_{i=2}^{\infty} \mathbb{P}(\emptyset) = 1 + \sum_{i=2}^{\infty} a = \infty$$

Now instead of using (A.3) we use that the infinite set of  $E_i$  becomes  $\Omega$ :

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} E_i\right) = \mathbb{P}(\Omega) \stackrel{\text{A.2}}{=} 1$$

We have reached a contradiction. This shows that  $\mathbb{P}(\emptyset) = 0$ . □

• **Claim:**  $A \cap B = \emptyset \implies \mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$ .

PROOF. Set  $E_1 = A$  and  $E_2 = B$  and  $E_k = \emptyset$  for all  $k \geq 3$ . Then:

$$\mathbb{P}(A \cup B) = \mathbb{P}\left(\bigcup_{i=1}^{\infty} E_i\right) \stackrel{\text{A.3}}{=} \sum_{i=1}^{\infty} \mathbb{P}(E_i) = \mathbb{P}(A) + \mathbb{P}(B) + \sum_{k=3}^{\infty} \mathbb{P}(\emptyset) = \mathbb{P}(A) + \mathbb{P}(B) + 0 = \mathbb{P}(A) + \mathbb{P}(B)$$
□

With this result, we can apply (A.3) indirectly to any finite sum of disjoint sets. All other sets are set to  $\emptyset$  and then (A.3) is applied. Not giving a formal argument, but if  $A, B, C$  are mutually disjoint we can set  $E_k = \emptyset$  for all  $k \geq 4$ .

$$\mathbb{P}(A \cup B \cup C) = \mathbb{P}\left(\bigcup_{i=1}^{\infty} E_i\right) \stackrel{\text{A.3}}{=} \sum_{i=1}^{\infty} \mathbb{P}(E_i) = \mathbb{P}(A) + \mathbb{P}(B) + \mathbb{P}(C) + 0 = \mathbb{P}(A) + \mathbb{P}(B) + \mathbb{P}(C)$$

• **Claim:**  $A \subset B \implies \mathbb{P}(A) \leq \mathbb{P}(B)$ .

PROOF. Since  $A \subset B$  we can split  $B$  into two disjoint parts:  $B = A \cup B - A$ . By axiom (A.3):

$$\mathbb{P}(B) = \mathbb{P}(A \cup B - A) \stackrel{\text{A.3}}{=} \mathbb{P}(A) + \mathbb{P}(B - A) \stackrel{\text{A.1}}{\geq} \mathbb{P}(A) \implies \mathbb{P}(A) \leq \mathbb{P}(B) \quad \square$$

• **Claim:**  $0 \leq \mathbb{P}(A) \leq 1$ .

PROOF. Since  $A \subset \Omega$  and by the previous proof:  $\mathbb{P}(A) \leq \mathbb{P}(\Omega) = 1$  Combining this with axiom (A.1):

$$0 \stackrel{\text{A.1}}{\leq} \mathbb{P}(A) \leq \mathbb{P}(\Omega) = 1 \implies 0 \leq \mathbb{P}(A) \leq 1 \quad \square$$

• **Claim:**  $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$ .

PROOF. Since  $A \cup A^c$  are disjoint, we get by finite version of (A.3):

$$\mathbb{P}(A \cup A^c) = \mathbb{P}(A) + \mathbb{P}(A^c)$$

Since  $\Omega = A \cup A^c$  we also get by (A.2):

$$\mathbb{P}(A \cup A^c) = \mathbb{P}(\Omega) = 1$$

Putting them together:

$$\mathbb{P}(A) + \mathbb{P}(A^c) = 1 \implies \mathbb{P}(A^c) = 1 - \mathbb{P}(A)$$

□

### 1.3

$\Omega$  is a sample space and  $A_1, A_2, \dots$  are events. We define:

$$B_n = \bigcup_{i=n}^{\infty} A_i, \quad C_n = \bigcap_{i=n}^{\infty} A_i$$

(a) **Claim:**  $B_1 \supset B_2 \supset \dots$

PROOF. This follows directly from the definition of  $B_n$ . First we see that:

$$B_1 = \bigcup_{i=1}^{\infty} A_i = A_1 \cup \left( \bigcup_{i=2}^{\infty} A_i \right) \supset \bigcup_{i=2}^{\infty} A_i = B_2, \implies B_1 \supset B_2$$

The same argument can be used for some  $k \in \mathbb{N}$ .

$$B_k = \bigcup_{i=k}^{\infty} A_i = A_k \cup \left( \bigcup_{i=k+1}^{\infty} A_i \right) \supset \bigcup_{i=k+1}^{\infty} A_i = B_{k+1}, \implies B_k \supset B_{k+1}$$

From this, it follows that  $B_1 \supset B_2 \supset \dots$

□

**Claim:**  $C_1 \subset C_2 \subset \dots$

PROOF. This also follows directly from the definition. First note that:

$$C_1 = \bigcap_{i=1}^{\infty} A_i = A_1 \cap \left( \bigcap_{i=2}^{\infty} A_i \right) \subset \bigcap_{i=2}^{\infty} A_i = C_2 \implies C_1 \subset C_2$$

And, in general, for some  $k \in \mathbb{N}$ .

$$C_k = \bigcap_{i=k}^{\infty} A_i = A_k \cap \left( \bigcap_{i=k+1}^{\infty} A_i \right) \subset \bigcap_{i=k+1}^{\infty} A_i = C_{k+1} \implies C_k \subset C_{k+1}$$

This shows that  $C_1 \subset C_2 \subset \dots$

□

(b) **Claim:**

$$\omega \in \bigcap_{n=1}^{\infty} B_n \iff \omega \in A_1, A_2, \dots$$

PROOF.

$\Rightarrow$ ) Assume  $\omega \in \bigcap_{n=1}^{\infty} B_n$ , which means  $\omega \in B_k, \forall k \in \mathbb{N}$ . Expanding the intersection:

$$\omega \in B_1 \cap B_2 \cap \dots \cap B_k \cap B_{k+1} \cap \dots$$

Now, from the definition of  $B_1$ , like we used above:

$$B_1 = \bigcup_{i=1}^{\infty} A_i = A_1 \cup \bigcup_{i=2}^{\infty} A_i = A_1 \cup B_2$$

So, we can write:  $B_1 \cap B_2 = (A_1 \cup B_2) \cap B_2 = (A_1 \cap B_2) \cup (B_2 \cap B_2) = (A_1 \cap B_2) \cup B_2$ . And in general:

$$B_k = \bigcup_{i=k}^{\infty} A_i = A_k \cup \bigcup_{i=k+1}^{\infty} A_i = A_k \cup B_{k+1},$$

so  $B_k \cap B_{k+1} = (A_k \cup B_{k+1}) \cap B_{k+1} = (A_k \cap B_{k+1}) \cup B_{k+1}$ . So, each  $B_i$  can be decomposed into  $(A_i \cap B_{i+1}) \cup B_{i+1}$  and the  $B_{i+1}$  is rewritten as its own intersection. Ultimately, we get:

$$\begin{aligned} \omega \in B_1 \cap B_2 \cap B_3 \cap \dots \cap B_k \cap B_{k+1} \cap \dots &\implies \\ \omega \in [(A_1 \cap B_2) \cup B_2] \cap B_3 \cap \dots \cap B_k \cap B_{k+1} \cap \dots &\implies \\ \omega \in [(A_1 \cap B_2) \cup (A_2 \cap B_3) \cup B_3] \cap \dots \cap B_k \cap B_{k+1} \cap \dots &\implies \\ \omega \in (A_1 \cap B_2) \cup (A_2 \cap B_3) \cup \dots \cup (A_k \cap B_{k+1}) \cup (A_{k+1} \cap B_{k+2}) \cup \dots &\implies \\ \omega \in (A_k \cap B_{k+1}) \forall k \in \mathbb{N} \text{ by assumption that } \omega \in B_k \forall k \in \mathbb{N} &\implies \\ \omega \in A_k, \forall k \in \mathbb{N} \end{aligned}$$

which means  $\omega \in A_1, A_2, \dots, A_k, A_{k+1}, \dots$  proving the first implication.

$\Leftarrow$ ) Now assume  $\omega \in A_1, A_2, \dots, A_k, \dots$  which means that  $\omega \in A_k$  for any  $k \in \mathbb{N}$ . Then,

$$\omega \in A_k \implies \omega \in A_k \cup \left( \bigcup_{i=k+1}^{\infty} A_i \right) = \bigcup_{i=k}^{\infty} A_i = B_k \implies \omega \in B_k,$$

So  $\omega$  is included in all  $B_k$  for  $k \in \mathbb{N}$  which means  $\omega$  is also in the intersection of all  $B_i$ .

$$\omega \in B_1, B_2, \dots, B_k, \dots \implies \omega \in \bigcap_{n=1}^{\infty} B_n$$

By implication both ways, we have proved the equivalency. □

(c) *Skipped.*

#### 1.4

Proving DeMorgan's laws. First in the case when  $i = \{1, 2, \dots, n\}$ .

**Claim:**

$$\left( \bigcup_{i=1}^n A_i \right)^c = \bigcap_{i=1}^n A_i^c$$

PROOF. Starting with  $x \in (\bigcup_{i=1}^n A_i)^c$ , we have the following equivalencies:

$$\begin{aligned} x \in \left( \bigcup_{i=1}^n A_i \right)^c &\iff x \notin \bigcup_{i=1}^n A_i \\ &\iff x \notin A_1 \text{ and } x \notin A_2 \text{ and } \dots \text{ and } x \notin A_n \\ &\iff x \in A_1^c \text{ and } x \in A_2^c \text{ and } \dots \text{ and } x \in A_n^c \\ &\iff x \in \bigcap_{i=1}^n A_i^c \end{aligned} \quad \square$$

**Claim:**

$$\left( \bigcap_{i=1}^n A_i \right)^c = \bigcup_{i=1}^n A_i^c$$

PROOF. Starting with  $x \in (\bigcap_{i=1}^n A_i)^c$ , we have the following equivalencies:

$$\begin{aligned} x \in \left( \bigcap_{i=1}^n A_i \right)^c &\iff x \notin \bigcap_{i=1}^n A_i \\ &\iff \exists p \in \{1, \dots, n\} \ x \notin A_p \\ &\iff \exists p \in \{1, \dots, n\} \ x \in A_p^c \\ &\iff x \in \bigcup_{i=1}^n A_i^c \end{aligned} \quad \square$$

Proving these results for a random index set is essentially the same, except finding some way of numerating the index set with  $i_1, \dots, i_n$  (since it has to be countably infinite) which I won't bother doing.

#### 1.5

Tossing a fair coin until we get exactly two heads. Description of the sample space:

$$\Omega = \{(\omega_1, \omega_2, \dots, \omega_{k-1}, \omega_k) : \omega_i \in \{H, T\}, \omega_{k-1} = \omega_k = H, \omega_{p-1} = H \Rightarrow \omega_p = T, p < k\}$$

Now, to calculate the probability that there will be exactly  $k$  tosses. Let us look at a few specific examples:

$$\begin{aligned} k = 2 : \{H, H\} &\implies \mathbb{P}(2) = \frac{1}{2^2} = \frac{1}{4} = 0.25 \\ k = 3 : \{T, H, H\} &\implies \mathbb{P}(3) = \frac{1}{2^3} = \frac{1}{8} = 0.125 \end{aligned}$$

For all tosses after  $k > 3$ , the last three tosses have to be  $\{T, H, H\}$ .

$$\begin{aligned}
k=4: & \left. \begin{array}{c} H \\ T \end{array} \right|, T, H, H = 2 \cdot \frac{1}{2^4} = \frac{1}{8} = 0.125 \\
k=5: & \left. \begin{array}{c} TT \\ TH \\ HT \end{array} \right|, T, H, H = 3 \cdot \frac{1}{2^5} = \frac{3}{32} = 0.09375 \\
k=6: & \left. \begin{array}{c} TTT \\ TTH \\ THT \\ HTT \\ HTH \end{array} \right|, T, H, H = 5 \cdot \frac{1}{2^6} = \frac{5}{64} = 0.078125 \\
k=7: & \left. \begin{array}{c} TTTT \\ TTTH \\ TTHT \\ THTT \\ HTTT \\ THTH \\ HTHT \\ HTTH \end{array} \right|, T, H, H = 8 \cdot \frac{1}{2^7} = \frac{1}{2^4} = \frac{1}{16} = 0.0625
\end{aligned}$$

We can see a pattern emerging. The last three tosses are fixed, and for the first  $k - 3$  tosses, we have to count all the combinations that do not contain 2 simultaneous H. The details are quite interesting, and it turns out that the number in the  $k - 3$  first tosses follows the Fibonacci sequence (as we can see above). Defining:

$$F(1) = 2, F(2) = 3, F(3) = 5, \dots$$

In general, we will then get:

$$\mathbb{P}(K = k) = \begin{cases} \frac{1}{2^k} & k = 2, 3 \\ \frac{F(k-2)}{2^k} & k > 3 \end{cases}$$

## 1.6

Let the sample space be  $\Omega = \{0, 1, 2, \dots\}$  (which is  $\{0\} \cup \mathbb{N}$ ). We are going to prove that there is no uniform distribution on this sample space. That is, if  $\mathbb{P}(A) = \mathbb{P}(B)$  whenever  $|A| = |B|$  (the cardinality of the sets) then  $\mathbb{P}$  cannot satisfy the axioms of probability.

The first two properties are satisfied. If we take some  $k \in \mathbb{N}$  and define  $A_1 = \{1, \dots, k\}$  in such a way that  $P(A_1) > 0$ , and generally define  $A_m = \{mk+1, mk+2, \dots, mk+k\}$ , then  $A_n \cap A_m = \emptyset$ . (This can be checked by setting  $k = 100$  and  $n = 5$  and  $m = 6$  for instance). Also,  $|A_n| = |A_m|$  and  $\mathbb{P}(A_n) = \mathbb{P}(A_m) > 0$ . This will cause problems in the third axiom:

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i) = \infty,$$

since we are summing an infinite number of positive values. The axioms of probability are not satisfied, so we cannot have a uniform distribution on  $\{0, 1, 2, \dots\}$ .



### 1.7

For some events  $A_1, A_2, \dots$ , we are going to show that:

$$\mathbb{P}\left(\bigcup_{n=1}^{\infty} A_n\right) \leq \sum_{n=1}^{\infty} \mathbb{P}(A_n).$$

PROOF. Following the hint, we define:

$$B_n = A_n - \bigcup_{i=1}^{n-1} A_i.$$

Next, we show that any  $B_m$  and  $B_n$  are disjoint for  $n, m \in \mathbb{N}$  and  $m > n$ .

$$\begin{aligned} B_m \cap B_n &= \left(A_m - \bigcup_{i=1}^{m-1} A_i\right) \cap \left(A_n - \bigcup_{i=1}^{n-1} A_i\right) \\ &= \left(A_m \cap \left(\bigcup_{i=1}^{m-1} A_i\right)^c\right) \cap \left(A_n \cap \left(\bigcup_{i=1}^{n-1} A_i\right)^c\right) \\ &= \left(A_m \cap \bigcap_{i=1}^{m-1} A_i^c\right) \cap \left(A_n \cap \bigcap_{i=1}^{n-1} A_i^c\right) \\ &= \left(A_m \cap A_{m-1}^c \cap \dots \cap A_n^c \cap \dots \cap A_2^c \cap A_1^c\right) \cap \left(A_n \cap A_{n-1}^c \cap \dots \cap A_2^c \cap A_1^c\right) \\ &= A_m \cap A_{m-1}^c \cap \dots \cap A_2^c \cap A_1^c \cap A_{n-1}^c \cap \dots \cap A_2^c \cap A_1^c \underbrace{\cap A_n^c \cap A_n}_{=\emptyset} \\ &= \emptyset \end{aligned}$$

Hence, the sets  $B_i$  are mutually disjoint. Next, we show that:

$$\left(\bigcup_{n=1}^{\infty} A_n\right) = \left(\bigcup_{n=1}^{\infty} B_n\right).$$

We will use an induction argument. For the first element we have:  $B_1 = A_1$ , and:

$$\begin{aligned} \bigcup_{i=1}^2 B_i &= B_1 \cup B_2 \\ &= A_1 \cup (A_2 - A_1) \\ &= A_1 \cup (A_2 \cap A_1^c) \\ &= (A_1 \cup A_2) \cap (A_1 \cup A_1^c) \\ &= (A_1 \cup A_2) \cap \Omega \\ &= (A_1 \cup A_2) \\ &= \bigcup_{i=1}^2 A_i \end{aligned}$$

So, we assume that for  $k \in \mathbb{N}$ , we have:

$$\left(\bigcup_{n=1}^k A_n\right) = \left(\bigcup_{n=1}^k B_n\right),$$

and show that this applies to  $k+1$ .

$$\begin{aligned}
\bigcup_{i=1}^{k+1} B_i &= B_{k+1} \cup \left( \bigcup_{i=1}^k B_i \right) \\
&= B_{k+1} \cup \left( \bigcup_{i=1}^k A_i \right) \\
&= \left( A_{k+1} - \bigcup_{i=1}^k A_i \right) \cup \left( \bigcup_{i=1}^k A_i \right) \\
&= \left( A_{k+1} \cap \left( \bigcup_{i=1}^k A_i \right)^c \right) \cup \left( \bigcup_{i=1}^k A_i \right) \\
&= \left( A_{k+1} \cup \bigcup_{i=1}^k A_i \right) \cap \left[ \left( \bigcup_{i=1}^k A_i \right)^c \cup \left( \bigcup_{i=1}^k A_i \right) \right] \\
&= \left( \bigcup_{i=1}^{k+1} A_i \right) \cap \Omega \\
&= \bigcup_{i=1}^{k+1} A_i
\end{aligned}$$

So, by induction, we have verified the equality for  $k + 1$ , so it follows that the sets are equal for all indexes.

By definition of  $B_n$ , we have  $B_n \subset A_n$  which means  $\mathbb{P}(B_n) \leq \mathbb{P}(A_n)$ . Using this, we can prove the main statement:

$$\mathbb{P} \left( \bigcup_{n=1}^{\infty} A_n \right) = \mathbb{P} \left( \bigcup_{n=1}^{\infty} B_n \right) = \sum_{n=1}^{\infty} \mathbb{P}(B_n) \leq \sum_{n=1}^{\infty} \mathbb{P}(A_n). \quad \square$$

### 1.8

Suppose that  $\mathbb{P}(A_i) = 1$  for each  $A_i$ . We are going to prove that:

$$\mathbb{P} \left( \bigcap_{i=1}^{\infty} A_i \right) = 1.$$

PROOF. We will prove this by induction. For events  $A_1$  and  $A_2$ , we know that  $\mathbb{P}(A_1) = 1$  and  $\mathbb{P}(A_2) = 1$ . By the probability axioms:

$$A_1 \subset A_1 \cup A_2 \subset \Omega \implies \mathbb{P}(A_1) \leq \mathbb{P}(A_1 \cup A_2) \leq \mathbb{P}(\Omega)$$

Using that  $\mathbb{P}(A_1) = 1$  and  $\mathbb{P}(\Omega) = 1$ :

$$1 \leq \mathbb{P}(A_1 \cup A_2) \leq 1 \implies \mathbb{P}(A_1 \cup A_2) = 1$$

By Lemma 1.6:

$$\begin{aligned}
\mathbb{P}(A_1 \cup A_2) &= \mathbb{P}(A_1) + \mathbb{P}(A_2) - \mathbb{P}(A_1 \cap A_2) \\
1 &= 2 - \mathbb{P}(A_1 \cap A_2) \\
\mathbb{P}(A_1 \cap A_2) &= 1
\end{aligned}$$

Assuming this holds for  $k \in \mathbb{N}$ , i.e.:

$$\mathbb{P}\left(\bigcap_{i=1}^k A_i\right) = 1.$$

To simplify the notation, we call this intersection  $W_k$ , so  $\mathbb{P}(W_k) = 1$ . Now we will show it also applies to  $k + 1$ .

$$A_1 \cap A_2 \cap \dots \cap A_k \cap A_{k+1} = W_k \cap A_{k+1}$$

Again, from the probability axioms:

$$W_k \subset W_k \cup A_{k+1} \subset \Omega \implies \mathbb{P}(W_k) \leq \mathbb{P}(W_k \cup A_{k+1}) \leq \mathbb{P}(\Omega)$$

Using that  $\mathbb{P}(W_k) = 1$  and  $\mathbb{P}(\Omega) = 1$ :

$$1 \leq \mathbb{P}(W_k \cup A_{k+1}) \leq 1 \implies \mathbb{P}(W_k \cup A_{k+1}) = 1$$

By Lemma 1.6, and using that  $\mathbb{P}(A_i) = 1$  for each  $i$ :

$$\begin{aligned} \mathbb{P}(W_k \cup A_{k+1}) &= \mathbb{P}(W_k) + \mathbb{P}(A_{k+1}) - \mathbb{P}(W_k \cap A_{k+1}) \\ 1 &= 2 - \mathbb{P}(W_k \cap A_{k+1}) \\ \mathbb{P}(W_k \cap A_{k+1}) &= 1 \\ \mathbb{P}\left(\bigcap_{i=1}^k A_i \cap A_{k+1}\right) &= 1 \\ \mathbb{P}\left(\bigcap_{i=1}^{k+1} A_i\right) &= 1 \end{aligned}$$

Which verifies that equality holds for  $k + 1$ . By induction, we can conclude that:

$$\mathbb{P}\left(\bigcap_{i=1}^{\infty} A_i\right) = 1. \quad \square$$

## 1.9

For a fixed  $B$  such that  $\mathbb{P}(B) > 0$ , we will show that  $\mathbb{P}(\cdot|B)$  satisfies the axioms of probability. Recalling the definition of conditional probability:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$$

• **Axiom 1.**  $\mathbb{P}(A|B) \geq 0$  for all  $A$ .

PROOF. The set  $A \cap B$  is an event wrt.  $\mathbb{P}(\cdot)$ , so  $\mathbb{P}(A \cap B) = a \geq 0$ .

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} = \frac{a}{\mathbb{P}(B)} \geq 0,$$

since  $\mathbb{P}(B) > 0$ . □

- **Axiom 2.**  $\mathbb{P}(\Omega|B) = 1$ .

PROOF.

$$\mathbb{P}(\Omega|B) = \frac{\mathbb{P}(\Omega \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B)}{\mathbb{P}(B)} = 1$$

□

- **Axiom 3.** If  $A_1, A_2, \dots$  are disjoint, then:

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i \middle| B\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i|B)$$

PROOF. First a supporting argument. If we define  $C_i := A_i \cap B$  for each  $i$ , then  $C_1, C_2, \dots$  are mutually disjoint sets and by (A.3):

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} C_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(C_i).$$

This means:

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i \middle| B\right) = \frac{\mathbb{P}(\bigcup_{i=1}^{\infty} A_i \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(\bigcup_{i=1}^{\infty} C_i)}{\mathbb{P}(B)} = \frac{\sum_{i=1}^{\infty} \mathbb{P}(C_i)}{\mathbb{P}(B)} = \sum_{i=1}^{\infty} \frac{\mathbb{P}(A_i \cap B)}{\mathbb{P}(B)} = \sum_{i=1}^{\infty} \mathbb{P}(A_i|B)$$

which proves the final axiom for the conditional probability.

□

### 1.10 Monty Hall

There are three doors, 1, 2, 3, and there is a prize between one of them. We don't know which door contains the prize, so the probability of selecting the correct door is evenly spread out:

$$P(1) = \frac{1}{3}, \quad P(2) = \frac{1}{3}, \quad P(3) = \frac{1}{3}$$

To simplify the calculations, we assume we pick door 1, and that the prize is not behind door 3. Now we will decide if it's better to switch or not. What is the probability that the host will open door III conditioned on where the prize is?

$$P(\text{III}|1) = \frac{1}{2} \quad (\text{Correct door is selected, door 3 is opened randomly})$$

$$P(\text{III}|2) = 1 \quad (\text{Host never reveals the prize})$$

$$P(\text{III}|3) = 0 \quad (\text{Host never reveals the prize})$$

Now, what is the probability that the prize is behind door number 2, given that door III was opened? This can be calculated by Bayes Law.

$$P(A_k|B) = \frac{P(B|A_k)P(A_k)}{\sum_{i=1}^n P(B|A_i)P(A_i)}$$

In our case, probability of door 2 containing the prize, given that door III was opened:

$$\begin{aligned}
P(2|III) &= \frac{\mathbb{P}(III|2)\mathbb{P}(2)}{\mathbb{P}(III|1)\mathbb{P}(1) + \mathbb{P}(III|2)\mathbb{P}(2) + \mathbb{P}(III|3)\mathbb{P}(3)} \\
&= \frac{(1)(\frac{1}{3})}{(\frac{1}{2})(\frac{1}{3}) + (1)(\frac{1}{3}) + (0)(\frac{1}{3})} \\
&= \frac{\frac{1}{3}}{\frac{1}{6} + \frac{1}{3} + 0} = \frac{1}{3} / \frac{1}{2} \\
&= \frac{2}{3}
\end{aligned}$$

Probability that door 1 contains the prize, given that door III was opened:

$$\begin{aligned}
P(1|III) &= \frac{\mathbb{P}(III|1)\mathbb{P}(1)}{\mathbb{P}(III|1)\mathbb{P}(1) + \mathbb{P}(III|2)\mathbb{P}(2) + \mathbb{P}(III|3)\mathbb{P}(3)} \\
&= \frac{(\frac{1}{2})(\frac{1}{3})}{(\frac{1}{2})(\frac{1}{3}) + (1)(\frac{1}{3}) + (0)(\frac{1}{3})} \\
&= \frac{\frac{1}{6}}{\frac{1}{6} + \frac{1}{3} + 0} = \frac{1}{6} / \frac{1}{2} \\
&= \frac{1}{3}
\end{aligned}$$

This shows that we are better off switching doors then remaining in the one we initially selected.

### 1.11

**Claim.** If  $A$  and  $B$  are independent, then  $A^c$  and  $B^c$  are independent.

PROOF. By definition of independence  $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$ . By property of probabilities:  $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$  and  $\mathbb{P}(B^c) = 1 - \mathbb{P}(B)$ . Applying Lemma 1.6 to  $A^c \cup B^c$ :

$$\mathbb{P}(A^c \cup B^c) = \mathbb{P}(A^c) + \mathbb{P}(B^c) - \mathbb{P}(A^c \cap B^c) \quad (2.11.1)$$

By property of complements, and using DeMorgan's law:

$$\mathbb{P}(A \cap B) = 1 - \mathbb{P}([A \cap B]^c) = 1 - \mathbb{P}(A^c \cup B^c)$$

Which we can rewrite in the following way, and then apply the independence of  $A$  and  $B$ :

$$\mathbb{P}(A^c \cup B^c) = 1 - \mathbb{P}(A \cap B) = 1 - \mathbb{P}(A)\mathbb{P}(B)$$

By property of complements, we can represent this as:

$$\begin{aligned}
\mathbb{P}(A^c \cup B^c) &= 1 - [1 - \mathbb{P}(A^c)][1 - \mathbb{P}(B^c)] \\
&= 1 - [1 - \mathbb{P}(B^c) - \mathbb{P}(A^c) + \mathbb{P}(A^c)\mathbb{P}(B^c)] \\
&= \mathbb{P}(A^c) + \mathbb{P}(B^c) - \mathbb{P}(A^c)\mathbb{P}(B^c)
\end{aligned} \quad (2.11.2)$$

By setting (2.11.1) equal to (2.11.2) we can deduce that  $\mathbb{P}(A^c)\mathbb{P}(B^c) = \mathbb{P}(A^c \cap B^c)$  which proves independence.  $\square$

*Alternative Proof.* Slightly more compact argument, but it still relies on Lemma 1.6.

$$\begin{aligned}
\mathbb{P}(A^c)\mathbb{P}(B^c) &= [1 - \mathbb{P}(A)][1 - \mathbb{P}(B)] \\
&= 1 - \mathbb{P}(A) - \mathbb{P}(B) + \mathbb{P}(A)\mathbb{P}(B) \\
&= 1 - [\mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A)\mathbb{P}(B)] \\
&\stackrel{IND}{=} 1 - [\mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)] \\
&\stackrel{L1.6}{=} 1 - \mathbb{P}(A \cup B) \\
&= \mathbb{P}([A \cup B]^c) \\
&= \mathbb{P}(A^c \cap B^c)
\end{aligned}$$

□

### 1.12

Initially, we have three cards that we call  $G$  (Green both sides),  $R$  (Red both sides) and  $M$  (mixed colors). Next we define  $r$  as 'Red side' and  $g$  as 'Green side'. We can define the joint distribution as follows:

	$G$	$M$	$R$	
$r$		1/6	2/6	1/2
$g$	2/6	1/6		1/2
	1/3	1/3	1/3	

From the marginal distributions, we can see that the initial probabilities for selecting each card is:

$$\mathbb{P}(G) = \frac{1}{3}, \quad \mathbb{P}(R) = \frac{1}{3}, \quad \mathbb{P}(M) = \frac{1}{3}$$

And the probabilities for randomly viewing a colored card side:

$$\mathbb{P}(g) = \frac{1}{2}, \quad \mathbb{P}(r) = \frac{1}{2}$$

Note that we do not have independence, since  $\mathbb{P}(G \cap g) = 2/6$  while  $\mathbb{P}(G)\mathbb{P}(g) = (1/3)(1/2) = 1/6$ .

We want to find the probability that we have a green card given that we observe a green side. This can be calculated by:

$$\mathbb{P}(G|g) = \frac{\mathbb{P}(G \cap g)}{\mathbb{P}(g)} = \frac{2/6}{1/2} = 4/6 = 2/3$$

### 1.13

A fair coin is thrown until we have both H and T.

(a) The sample space is:

$$\Omega = \{(\omega_1, \dots, \omega_K) : \omega_i \in \{H, T\}, 1 \leq i \leq K \text{ and } \omega_{K-1} \neq \omega_K, K \geq 2\}$$

(b) The probability of getting 3 tosses can only happen in two ways  $(H, H, T)$  and  $(T, T, H)$ . So:

$$P(K = 3) = 2 \cdot \frac{1}{2^3} = \frac{1}{4}$$

(it would be more complicated if it were e.g.  $K \leq 3$ ).

### 1.14

**Claim:** If  $\mathbb{P}(A) = 0$  or  $\mathbb{P}(A) = 1$ , then  $A$  is independent of every other event.

PROOF. First consider the case  $\mathbb{P}(A) = 0$ . Obviously, for any  $E \in \Omega$ :

$$\mathbb{P}(A)\mathbb{P}(E) = 0 \cdot \mathbb{P}(E) = 0.$$

By the axioms of probability:  $A \cap E \subset A$  so  $\mathbb{P}(A \cap E) \leq \mathbb{P}(A) = 0$  and since for any event:  $0 \leq \mathbb{P}(A \cap E)$  we can conclude that  $\mathbb{P}(A \cap E) = 0$ . So:

$$\mathbb{P}(A \cap E) = 0 = \mathbb{P}(A)\mathbb{P}(E) \implies \mathbb{P}(A \cap E) = \mathbb{P}(A)\mathbb{P}(E)$$

which proves independence.

Now consider  $\mathbb{P}(A) = 1$ . Then, for any event  $E \in \Omega$ :

$$\mathbb{P}(A)\mathbb{P}(E) = 1 \cdot \mathbb{P}(E) = \mathbb{P}(E).$$

Since  $A \subset A \cup E \subset \Omega$ , then  $\mathbb{P}(A) \leq \mathbb{P}(A \cup E) \leq \mathbb{P}(\Omega)$ . We know the probabilities of  $A$  and  $\Omega$ , so  $1 \leq \mathbb{P}(A \cup E) \leq 1$  means that  $\mathbb{P}(A \cup E) = 1$ . By Lemma 1.6:

$$\begin{aligned} \mathbb{P}(A \cup E) &= \mathbb{P}(A) + \mathbb{P}(E) - \mathbb{P}(A \cap E) \\ 1 &= 1 + \mathbb{P}(E) - \mathbb{P}(A \cap E) \\ \mathbb{P}(A \cap E) &= \mathbb{P}(E) \end{aligned}$$

In conclusion:

$$\mathbb{P}(A \cap E) = \mathbb{P}(E) = \mathbb{P}(A)\mathbb{P}(E) \implies \mathbb{P}(A \cap E) = \mathbb{P}(A)\mathbb{P}(E)$$

which proves independence. □

**Claim:** If  $A$  is independent of itself, then  $\mathbb{P}(A) = 0$  or  $\mathbb{P}(A) = 1$ .

PROOF. If  $A$  is independent of itself, we can write:

$$\begin{aligned} \mathbb{P}(A \cap A) &= \mathbb{P}(A)\mathbb{P}(A) \\ \mathbb{P}(A) &= \mathbb{P}(A)^2 \end{aligned}$$

since  $A \cap A = A$ . We can write this as an equation, by defining  $p := \mathbb{P}(A)$ .

$$\begin{aligned} p &= p^2 \\ p^2 - p &= 0 \\ p(p - 1) &= 0 \end{aligned}$$

This equation is solved when  $p = \mathbb{P}(A) = 1$  or  $p = \mathbb{P}(A) = 0$ . □

### 1.15

The probability that a child has blue eyes is  $1/4$ ,  $\mathbb{P}(B) = 1/4$ . We assume independence in the eye color of the children.

(a) Given that one child has blue eyes, what is the probability that 2 or more children have blue eyes? Since it's given that one child has blue eyes, we just need to find the probability that one, or both, of the remaining children have blue eyes. Define  $B_i$ : as the event that child  $i$  has blue eyes, and  $\neg B_i$  the event that child  $i$  does NOT have blue eyes. We only need to sum up the following three probabilities:

$$\begin{aligned}\mathbb{P}(B_1 \cap B_2) &= \mathbb{P}(B_1)\mathbb{P}(B_2) = \left(\frac{1}{4}\right)\left(\frac{1}{4}\right) = \frac{1}{16} \\ \mathbb{P}(B_1 \cap \neg B_2) &= \mathbb{P}(B_1)\mathbb{P}(\neg B_2) = \left(\frac{1}{4}\right)\left(\frac{3}{4}\right) = \frac{3}{16} \\ \mathbb{P}(\neg B_1 \cap B_2) &= \mathbb{P}(\neg B_1)\mathbb{P}(B_2) = \left(\frac{3}{4}\right)\left(\frac{1}{4}\right) = \frac{3}{16}\end{aligned}$$

By summing these up, we find that the probability of two or more children having blue eyes is  $\frac{7}{16}$ .

(b) Which child has blue eyes is irrelevant. The probability will be the same:  $\frac{7}{16}$ .

### 1.16

Proving Lemma 1.14. If  $A$  and  $B$  are independent events, then  $\mathbb{P}(A|B) = \mathbb{P}(A)$ . Also, for any pair of events  $A$  and  $B$ ,

$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B)\mathbb{P}(B) = \mathbb{P}(B|A)\mathbb{P}(A).$$

PROOF. Suppose  $A$  and  $B$  are independent. By definition of independence,  $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$ . By definition of conditional probability:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(A)\mathbb{P}(B)}{\mathbb{P}(B)} = \mathbb{P}(A),$$

which proves the first statement. Again, by definition of conditional probability:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} \implies \mathbb{P}(A \cap B) = \mathbb{P}(A|B)\mathbb{P}(B)$$

$$\mathbb{P}(B|A) = \frac{\mathbb{P}(B \cap A)}{\mathbb{P}(A)} \implies \mathbb{P}(A \cap B) = \mathbb{P}(B|A)\mathbb{P}(A),$$

since  $A \cap B = B \cap A$  which proves the remaining two statements. □



**1.17**

Show that

$$\mathbb{P}(A \cap B \cap C) = \mathbb{P}(A|B \cap C)\mathbb{P}(B|C)\mathbb{P}(C).$$

PROOF. Define  $E := B \cap C$ , so  $A \cap B \cap C = A \cap E$ . By Lemma 1.14:

$$\mathbb{P}(A \cap B \cap C) = \mathbb{P}(A \cap E) = \mathbb{P}(A|E)\mathbb{P}(E) = \mathbb{P}(A|B \cap C)\mathbb{P}(B \cap C) \quad (2.17.1)$$

By Lemma 1.14 again:

$$\mathbb{P}(B \cap C) = \mathbb{P}(B|C)\mathbb{P}(C),$$

which we can replace in equation (2.17.1) and get the desired result:

$$\mathbb{P}(A \cap B \cap C) = \mathbb{P}(A|B \cap C)\mathbb{P}(B|C)\mathbb{P}(C) \quad \square$$

**1.18**

Suppose  $k$  events form a partition of  $\Omega$ , i.e. they are disjoint and  $\bigcup_{i=1}^k A_i = \Omega$ . Assume that  $\mathbb{P}(B) > 0$ . If  $\mathbb{P}(A_1|B) < \mathbb{P}(A_1)$  then  $\mathbb{P}(A_i|B) > \mathbb{P}(A_i)$  for some  $i = 2, 3, \dots, k$ .

PROOF. By summing over all  $A_i$  with and without conditioning on  $B$ , we get the following two equalities:

$$\sum_{i=1}^k \mathbb{P}(A_i|B) = 1, \quad \sum_{i=1}^k \mathbb{P}(A_i) = 1$$

We can set these equal to each other:

$$\sum_{i=1}^k \mathbb{P}(A_i|B) = \sum_{i=1}^k \mathbb{P}(A_i)$$

Removing the terms for  $A_1$  from the sums:

$$\mathbb{P}(A_1|B) + \sum_{i=2}^k \mathbb{P}(A_i|B) = \mathbb{P}(A_1) + \sum_{i=2}^k \mathbb{P}(A_i)$$

Since  $\mathbb{P}(A_1|B) < \mathbb{P}(A_1)$ , then  $\mathbb{P}(A_1) - \mathbb{P}(A_1|B) = a > 0$  for some  $a \in \mathbb{R}$ . By subtracting  $\mathbb{P}(A_1|B)$  from both sides, we can write:

$$\begin{aligned} \sum_{i=2}^k \mathbb{P}(A_i|B) &= \mathbb{P}(A_1) - \mathbb{P}(A_1|B) + \sum_{i=2}^k \mathbb{P}(A_i) \\ \sum_{i=2}^k \mathbb{P}(A_i|B) &= a + \sum_{i=2}^k \mathbb{P}(A_i) > \sum_{i=2}^k \mathbb{P}(A_i) \end{aligned}$$

So, we get:

$$\sum_{i=2}^k \mathbb{P}(A_i|B) > \sum_{i=2}^k \mathbb{P}(A_i)$$

which means there exists some  $i \in \{2, 3, \dots, k\}$  such that  $\mathbb{P}(A_i|B) > \mathbb{P}(A_i)$ .  $\square$

### 1.19

Define  $M$  as Mac users,  $W$  as Windows users, and  $L$  as Linux users. We have:

$$\mathbb{P}(M) = 0.3, \quad \mathbb{P}(W) = 0.5, \quad \mathbb{P}(L) = 0.2$$

The computers are infected with a virus, which give the following probabilities:

$$\mathbb{P}(V|M) = 0.65, \quad \mathbb{P}(V|W) = 0.82, \quad \mathbb{P}(V|L) = 0.50$$

A person is selected at random and we learn that her computer has the virus. What is the probability that she is using Windows? Or, what is  $\mathbb{P}(W|V)$ ? We calculate this with Bayes Law.

$$\begin{aligned} \mathbb{P}(W|V) &= \frac{\mathbb{P}(V|W)\mathbb{P}(W)}{\mathbb{P}(V|M)\mathbb{P}(M) + \mathbb{P}(V|W)\mathbb{P}(W) + \mathbb{P}(V|L)\mathbb{P}(L)} \\ &= \frac{(0.82)(0.5)}{(0.65)(0.3) + (0.82)(0.5) + (0.5)(0.2)} \\ &= 0.5815 \end{aligned}$$

### 1.20

We have a box containing 5 coins,  $C_1, \dots, C_5$ . Each coin is selected at random, so  $\mathbb{P}(C_i) = 1/5$  for all coins. They have the following probabilities of getting  $H$  when flipped ( $p_i = \mathbb{P}(H|C_i)$ ):

$$p_1 = 0, \quad p_2 = 1/4, \quad p_3 = 1/2, \quad p_4 = 3/4, \quad p_5 = 1$$

(a) We flip a coin and get a head. Calculate  $\mathbb{P}(C_i|H)$  for all coins. First an intermediary calculation:

$$\begin{aligned} \sum_{i=1}^5 \mathbb{P}(H|C_i)\mathbb{P}(C_i) &= \mathbb{P}(H|C_1)\mathbb{P}(C_1) + \mathbb{P}(H|C_2)\mathbb{P}(C_2) + \mathbb{P}(H|C_3)\mathbb{P}(C_3) + \mathbb{P}(H|C_4)\mathbb{P}(C_4) + \mathbb{P}(H|C_5)\mathbb{P}(C_5) \\ &= (0) \left(\frac{1}{5}\right) + \left(\frac{1}{4}\right) \left(\frac{1}{5}\right) + \left(\frac{1}{2}\right) \left(\frac{1}{5}\right) + \left(\frac{3}{4}\right) \left(\frac{1}{5}\right) + (1) \left(\frac{1}{5}\right) \\ &= 0 + \frac{1}{20} + \frac{2}{20} + \frac{3}{20} + \frac{4}{20} \\ &= \frac{1}{2} \end{aligned}$$

Calculating the posterior for each  $C_i$ .

$$\begin{aligned} \mathbb{P}(C_1|H) &= \frac{\mathbb{P}(H|C_1)\mathbb{P}(C_1)}{\sum_{i=1}^5 \mathbb{P}(H|C_i)\mathbb{P}(C_i)} = \frac{0}{\frac{1}{2}} = 0 \\ \mathbb{P}(C_2|H) &= \frac{\mathbb{P}(H|C_2)\mathbb{P}(C_2)}{\sum_{i=1}^5 \mathbb{P}(H|C_i)\mathbb{P}(C_i)} = \frac{\frac{1}{20}}{\frac{1}{2}} = 1/10 \\ \mathbb{P}(C_3|H) &= \frac{\mathbb{P}(H|C_3)\mathbb{P}(C_3)}{\sum_{i=1}^5 \mathbb{P}(H|C_i)\mathbb{P}(C_i)} = \frac{\frac{2}{20}}{\frac{1}{2}} = 2/10 \end{aligned}$$

$$\mathbb{P}(C_4|H) = \frac{\mathbb{P}(H|C_4)\mathbb{P}(C_4)}{\sum_{i=1}^5 \mathbb{P}(H|C_i)\mathbb{P}(C_i)} = \frac{\frac{3}{20}}{\frac{1}{2}} = 3/10$$

$$\mathbb{P}(C_5|H) = \frac{\mathbb{P}(H|C_5)\mathbb{P}(C_5)}{\sum_{i=1}^5 \mathbb{P}(H|C_i)\mathbb{P}(C_i)} = \frac{\frac{4}{20}}{\frac{1}{2}} = 4/10$$

(b) We toss the coin again. Finding the probability  $\mathbb{P}(H_2|H_1)$  ( $H_i$ : getting  $H$  on toss  $i$ ). We are still using the same random coin, so we must calculate the probability for all coins together.

### 1.21

Proportion of H vs T for  $p = 0.3$ .


```
# 1.21 - Plotting proportion of H vs T
n = 1000
p = 0.3
coinTosses = sample(c("H","T"), prob = c(p, 1-p), size = n, replace = TRUE)
proportionHeads = rep(0, n)
headCount = 0
for(i in 1:n) {
  if (coinTosses[i] == "H") {
    headCount = headCount + 1
  }
  proportionHeads[i] = headCount/i
}

# PDF
pdf("~/ALLSTAT/ch1_2.21a.pdf")
plot(x = 1:n, y = proportionHeads, type = "l", ylim = c(0,1),
     main = paste0("Proportion heads for p=", p))
abline(h = p, col = "red", lty = "dashed")
dev.off()
```

R

### Result

After some initial randomness due to few samples, we can see that the simulated results stabilizes around 0.3, as expected.



Proportion of H vs T for  $p = 0.03$ .


```
# 1.21 - Plotting proportion of H vs T
n = 5000
p = 0.03
coinTosses = sample(c("H","T"), prob = c(p, 1-p), size = n, replace = TRUE)
proportionHeads = rep(0, n)
headCount = 0
for(i in 1:n) {
  if (coinTosses[i] == "H") {
    headCount = headCount + 1
  }
  proportionHeads[i] = headCount/i
}

# PDF
pdf("~/ALLSTAT/ch1_2.21b.pdf")
plot(x = 1:n, y = proportionHeads, type = "l", ylim = c(0,1),
     main = paste0("Proportion heads for p=", p))
abline(h = p, col = "red", lty = "dashed")
dev.off()
```

R

## Result

After some initial randomness due to few samples, we can see that the simulated results stabilizes around 0.03, as expected. Did a simulation of 5000 to make the 'convergence' clearer. Note that the y-axis only goes up to 0.1 in this plot.



## 1.22

Investigating some properties of Binomial random variables.

```
# 1.22 - Binomial Random Variables
REP = 10 # Number of simulations per n
p = 0.3 # Probability of H
sim10 = rep(0, REP)
sim100 = rep(0, REP)
sim1000 = rep(0, REP)
# Simulating 10
for(i in 1:REP) {
  # 1 means head
  coinTosses = sample(c(1,0), prob = c(p, 1-p), size = 10, replace = TRUE)
  sim10[i] = sum(coinTosses)
}
# Simulating 100
for(i in 1:REP) {
  # 1 means head
  coinTosses = sample(c(1,0), prob = c(p, 1-p), size = 100, replace = TRUE)
  sim100[i] = sum(coinTosses)
}
# Simulating 1000
for(i in 1:REP) {
  # 1 means head
  coinTosses = sample(c(1,0), prob = c(p, 1-p), size = 1000, replace = TRUE)
  sim1000[i] = sum(coinTosses)
}
df = data.frame(
  SIM10 = sim10,
  SIM100 = sim100,
  SIM1000 = sim1000
)
# Output
df
apply(df, 2, mean)
```

R

As seen in the results, the mean of the 10 simulations is close to  $np$  which would be 3, 30 and 300.

```
> df
  SIM10 SIM100 SIM1000
1     1     32     282
2     2     30     298
3     3     33     319
4     3     28     290
5     3     29     284
6     2     34     286
7     4     37     310
8     2     30     294
9     2     30     297
10    2     31     326
> apply(df, 2, mean)
  SIM10 SIM100 SIM1000
  2.4    31.4   298.6
```

### 1.23

Simulating conditional probabilities. First we simulate an independent experiment.

```
# 1.23 - Simulating a fair die
options(digits=8)
numberOfTosses = 10000
A = c(2, 4, 6)
B = c(1, 2, 3, 4)
AandB = intersect(A, B) # c(2, 4)

dieTosses = sample(1:6, size = numberOfTosses, replace = TRUE)

# Calculating P(A), P(B), P(A)*P(B) and P(A cap B)
PA = sum(dieTosses %in% A)/numberOfTosses
PB = sum(dieTosses %in% B)/numberOfTosses
PAandB = sum(dieTosses %in% AandB)/numberOfTosses

# Output
PA
PB
PA*PB
PAandB
```

R

Results from the calculation. As we can see, the estimates for  $\mathbb{P}(A) \approx 1/2$  and  $\mathbb{P}(B) \approx 2/3$ . Also  $\mathbb{P}(A \cap B) \approx \mathbb{P}(A)\mathbb{P}(B) \approx 1/3$ . Differences are probably due to rounding errors.

```
> # Output
> PA
[1] 0.5005
> PB
[1] 0.6695
> PA*PB
[1] 0.33508475
> PAandB
[1] 0.3344
```

Now we will construct an experiment with a conditional probability. We will use a fair coin and a die. When we get heads, this will correspond to 1 and the die is unchanged. If we get tails, this will be 2 and will double the die count; so if we get tails and we roll a 2, this will give us a 4.

	1	2	3	4	5	6	7	8	9	10	11	12
H	1/12	1/12	1/12	1/12	1/12	1/12						
T		1/12		1/12		1/12		1/12		1/12		1/12

Define the events:  $A = \{2, 3, 4, 5, 6\}$  and  $B = \{2, 4, 6, 8\}$  which will give  $A \cap B = \{2, 4, 6\}$ . The theoretical probabilities are:

$$\begin{aligned}\mathbb{P}(A) &= 8/12 = 2/3 = 0.666 \\ \mathbb{P}(B) &= 7/12 \approx 0.5833 \\ \mathbb{P}(A)\mathbb{P}(B) &= 7/18 \approx 0.3888 \\ \mathbb{P}(A \cap B) &= 1/2 = 0.5\end{aligned}$$

Simulating conditional probabilities.

```
# 1.23 - Simulating a conditional probability
options(digits=8)
numberOfTosses = 10000
A = c(2, 3, 4, 5, 6)
B = c(2, 4, 6, 8)
AandB = intersect(A, B) # c(2, 4, 6)

dieTosses = sample(1:6, size = numberOfTosses, replace = TRUE)
coinTosses = sample(1:2, size = numberOfTosses, replace = TRUE)
jointToss = dieTosses*coinTosses

# Calculating P(A), P(B), P(A)*P(B) and P(A cap B)
PA = sum(jointToss %in% A)/numberOfTosses
PB = sum(jointToss %in% B)/numberOfTosses
PAandB = sum(jointToss %in% AandB)/numberOfTosses

# Output
PA
PB
PA*PB
PAandB
```

R

```
> # Output
> PA
[1] 0.6691
> PB
[1] 0.5825
> PA*PB
[1] 0.38975075
> PAandB
[1] 0.4995
```

Repeating the theoretical values from previous page:

$$\mathbb{P}(A) = 8/12 = 2/3 = 0.666$$

$$\mathbb{P}(B) = 7/12 \approx 0.5833$$

$$\mathbb{P}(A)\mathbb{P}(B) = 7/18 \approx 0.3888$$

$$\mathbb{P}(A \cap B) = 1/2 = 0.5$$

The simulated results are very close to the theoretical calculations. We can see that we do not have independence, as expected.



## 2 Chapter 2 - Random Variables

### Exercises

#### 2.1

**Claim:**  $\mathbb{P}(X = x) = F(x^+) - F(x^-)$ . (Discrete)

PROOF. By definition of the CDF:

$$F(x^+) = \lim_{z \downarrow x} F(z) = \lim_{z \downarrow x} \mathbb{P}(X \leq z), \quad F(x^-) = \lim_{y \uparrow x} F(y) = \lim_{y \uparrow x} \mathbb{P}(X \leq y)$$

(so  $y < x$  and  $y \rightarrow x$ , and  $x < z$  and  $x \leftarrow z$ ). By the right continuous property, we can deduce that  $z > y$  and we can set  $z = x$  and  $y = x - 1$ .

$$\mathbb{P}(X \leq x^+) = \mathbb{P}(X \leq x) = \mathbb{P}(X = x) + \mathbb{P}(X \leq x - 1), \quad \mathbb{P}(X \leq x^-) = \mathbb{P}(X \leq x - 1)$$


So:

$$\begin{aligned} \mathbb{P}(X = x) &= \mathbb{P}(X = x) + \mathbb{P}(X \leq x - 1) - \mathbb{P}(X \leq x - 1) \\ &= \mathbb{P}(X \leq x) - \mathbb{P}(X \leq x - 1) \\ &= \mathbb{P}(X \leq x^+) - \mathbb{P}(X \leq x^-) \\ &= F(x^+) - F(x^-) \end{aligned}$$

□

#### 2.2

Let  $X$  be such that  $\mathbb{P}(X = 2) = \mathbb{P}(X = 3) = 1/10$  and  $\mathbb{P}(X = 5) = 8/10$ . Here is a plot of the CDF.



By reading the plot, we can see that:

$$\mathbb{P}(2 < X \leq 4.8) = F(4.8) - F(2) = 2/10 - 1/10 = 1/10$$

$$\mathbb{P}(2 \leq X \leq 4.8) = F(4.8) = 2/10$$

### 2.3

**Lemma 2.15** Let  $F$  be the CDF for a random variable  $X$ . Then:

1.  $\mathbb{P}(X = x) = F(x) - F(x^-)$
2.  $\mathbb{P}(x < X \leq y) = F(y) - F(x)$
3.  $\mathbb{P}(X > x) = 1 - F(x)$
4. If  $X$  is continuous, then

$$F(b) - F(a) = \mathbb{P}(a < X < b) = \mathbb{P}(a \leq X < b) = \mathbb{P}(a < X \leq b) = \mathbb{P}(a \leq X \leq b)$$

PROOF. We will prove each statement in turn. (1.) was proved in exercise **2.1**. Doing (3) first, since we need it to prove (2).

(3) By definition of complements of sets  $A = \{X > x\}$  means  $A^c = \{X \leq x\}$ , and it follows that:

$$\mathbb{P}(X > x) = \mathbb{P}(A) = 1 - \mathbb{P}(A^c) = 1 - \mathbb{P}(X \leq x) = 1 - F(x).$$

(2) Assume  $x < y$ . We will need that  $\{X > x\} \cup \{X \leq y\} = \Omega$ , and we will also use Lemma 1.6 (in reverse).

$$\begin{aligned} \mathbb{P}(x < X \leq y) &= \mathbb{P}(\{X > x\} \cap \{X \leq y\}) \\ &= \mathbb{P}(X > x) + \mathbb{P}(X \leq y) - \mathbb{P}(\{X > x\} \cup \{X \leq y\}) \\ &= 1 - F(x) + F(y) - 1 \\ &= F(y) - F(x) \end{aligned}$$

(4) Similar argument for all cases, so will just do one. We just need to turn the inequalities into strict inequalities. For continuous random variables, pointwise probabilities are 0. Again, we will need to use  $\{X > a\} \cup \{X < b\} = \Omega$ .

Define  $A := \{a \leq X\}$  and  $B := \{X < b\}$ . First, we make the following observation:

$$\begin{aligned} \mathbb{P}(A) &= \mathbb{P}(\{a \leq X\}) \\ &= \mathbb{P}(\{a = X\} \cup \{a < X\}) \\ &= \mathbb{P}(\{a = X\}) + \mathbb{P}(\{a < X\}) - \mathbb{P}(\{a = X\} \cap \{a < X\}) \\ &= 0 + \mathbb{P}(A') - 0 \\ &= \mathbb{P}(A') \end{aligned}$$

where  $A' = \{a < X\}$ . We get 0 for the pointwise probability, since this is continuous, and we get 0 because the sets are disjoint. We have shown that  $\mathbb{P}(A) = \mathbb{P}(A')$  and can use this to conclude the proof.

$$\begin{aligned} \mathbb{P}(a \leq X < b) &= \mathbb{P}(\{a \leq X\} \cap \{X < b\}) \\ &= \mathbb{P}(A \cap B) \\ &= \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cup B) \\ &= \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(\Omega) \\ &= \mathbb{P}(A') + \mathbb{P}(B) - \mathbb{P}(A' \cup B) \\ &= \mathbb{P}(A' \cap B) \\ &= \mathbb{P}(a < X < b) \end{aligned}$$


□

## 2.4

$X$  has the probability density (PDF):

$$f_X(x) = \begin{cases} 1/4 & 0 < x < 1 \\ 3/8 & 3 < x < 5 \\ 0 & \text{otherwise} \end{cases}$$

Plot of the PDF:



From the relatively simple structure, we can easily determine the area under the graph:

$$A = (1) \left( \frac{1}{4} \right) + (2) \left( \frac{3}{8} \right) = \frac{2}{8} + \frac{6}{8} = 1$$

(a) Finding the CDF by integrating the PDF. We will split up the integral in several parts. First for the case when  $y \in (0, 1)$ :

$$F_X(y) = \int_{-\infty}^y f_X(t) dt = \frac{1}{4} \int_0^y 1 dt = \frac{1}{4} [t]_0^y = \frac{y}{4}$$

When  $y = 1$  we have  $F_X(1) = 1/4$ . Next, we must consider the case  $y \in (1, 3)$ . Here the PDF is 0, so it doesn't increase. It remains constant at  $1/4$  (since the CDF doesn't decrease).

$$F_X(y) = \frac{1}{4}$$

Next is the case  $y \in (3, 5)$ . Consider the intermediary integral:

$$I_1 = \int_3^y \frac{3}{8} dt = \frac{3}{8} [t]_3^y = \frac{3y - 9}{8}$$

For values  $y \in (3, 5)$  we start on  $1/4$ , so the CDF in this region becomes:

$$F_X(y) = \frac{3y - 9}{8} + \frac{1}{4}$$

So, the full expression for the CDF becomes:

$$F_X(y) = \begin{cases} y/4 & y \in (0, 1) \\ 1/4 & y \in (1, 3) \\ \frac{3y-9}{8} + \frac{1}{4} & y \in (3, 5) \\ 1 & y \geq 5 \end{cases}$$

Note that when  $y = 5$  we get:

$$F_X(5) = \frac{3(5)-9}{8} + \frac{1}{4} = \frac{6}{8} + \frac{2}{8} = 1$$

Plot of the CDF:



(b) Defining  $Y = 1/X$  and finding the PDF of  $Y$ . Following the hint we are given, we will consider the following three sets:

$$A_1 = \frac{1}{5} \leq y \leq \frac{1}{3}, \quad A_2 = \frac{1}{3} \leq y \leq 1, \quad A_3 = y \geq 1$$

Where  $A_1$  corresponds to  $(3, 5)$ ,  $A_2$  to  $(1, 3)$  and  $A_3$  to  $(0, 1)$ . We can express the CDF for  $F_Y(y)$  in terms of  $F_X(x)$ :

$$\begin{aligned} F_Y(y) &= \mathbb{P}(Y \leq y) = \mathbb{P}\left(\frac{1}{X} \leq y\right) \\ &= \mathbb{P}\left(X \geq \frac{1}{y}\right) \\ &= 1 - \mathbb{P}\left(X \leq \frac{1}{y}\right) \\ &= 1 - F_X\left(\frac{1}{y}\right) \end{aligned}$$

First, we consider  $A_1 : y \in [1/5, 1/3]$ , and when we input  $1/y$  to  $F_X(\cdot)$ , it will be in  $(3, 5)$ . So:

$$\begin{aligned}
 F_Y(y) &= 1 - F_X(1/y) \\
 &= 1 - \left( \frac{3(\frac{1}{y}) - 9}{8} + \frac{1}{4} \right) \\
 &= 1 - \frac{3 - 9y}{8y} - \frac{1}{4} \\
 &= \frac{3}{4} + \frac{9y - 3}{8y} \\
 &= \frac{15y - 3}{8y}
 \end{aligned}$$

Next, we consider  $A_2 : y \in [1/3, 1]$ . The input to  $F_X(\cdot)$  will be in  $(1, 3)$ :

$$\begin{aligned}
 F_Y(y) &= 1 - F_X(1/y) \\
 &= 1 - \frac{1}{4} \\
 &= \frac{3}{4}
 \end{aligned}$$

Next, we consider  $A_3 : y \geq 1$ . The input to  $F_X(\cdot)$  will be in  $(0, 1)$ :

$$\begin{aligned}
 F_Y(y) &= 1 - F_X(1/y) \\
 &= 1 - \frac{1}{y} \\
 &= 1 - \frac{1}{4y}
 \end{aligned}$$


Also, whenever  $y < 1/5$ , then  $1/y > 5$  which means  $F_X(\cdot) = 1$ , and so:

$$F_Y(y) = 1 - F_X(1/y) = 1 - 1 = 0.$$

This gives a full description of the CDF for  $F_Y(y)$ .

$$F_Y(y) = \begin{cases} 0 & y < 1/5 \\ \frac{15y - 3}{8y} & 1/5 \leq y \leq 1/3 \\ \frac{3}{4} & 1/3 \leq y \leq 1 \\ 1 - \frac{1}{4y} & y \geq 1 \end{cases}$$

Plot of CDF:



Finally, we can find the PDF of  $Y$ . We differentiate each of the parts in the CDF. When  $y \in (1/5, 1/3)$ :


$$\frac{d}{dy} \left( \frac{15y-3}{8y} \right) = \frac{3}{8y^2}$$

When  $y \geq 1$ :

$$\frac{d}{dy} \left( 1 - \frac{1}{4y} \right) = \frac{1}{4y^2}$$

(All other parts are constant, so they become 0). This gives us the PDF and its plot:

$$f_Y(y) = \begin{cases} 0 & y < 1/5 \\ \frac{3}{8y^2} & 1/5 \leq y \leq 1/3 \\ 0 & 1/3 < y < 1 \\ \frac{1}{4y^2} & y \geq 1 \end{cases}$$



## 2.5

Let  $X$  and  $Y$  be discrete RV.  $X$  and  $Y$  are independent if and only if  $f_{X,Y}(x, y) = f_X(x)f_Y(y)$  for all  $x$  and  $y$ .

PROOF.

$\Rightarrow$ ) Assume that  $X$  and  $Y$  are independent. That means that for any  $x, y$ , we have

$$\mathbb{P}(X = x \cap Y = y) = \mathbb{P}(X = x)\mathbb{P}(Y = y)$$

Starting with the definition of the joint pdf:

$$\begin{aligned} f_{X,Y}(x, y) &= \mathbb{P}(X = x, Y = y) \\ &= \mathbb{P}(X = x \cap Y = y) \\ &= \mathbb{P}(X = x)\mathbb{P}(Y = y) \\ &= f_X(x)f_Y(y) \end{aligned}$$

Which shows that  $f_{X,Y}(x, y) = f_X(x)f_Y(y)$  for all  $x$  and  $y$ .

$\Leftarrow$ ) Assume that  $f_{X,Y}(x, y) = f_X(x)f_Y(y)$  for all  $x$  and  $y$ . By definition:

$$\begin{aligned} f_{X,Y}(x, y) &= \mathbb{P}(X = x, Y = y) \\ &= \mathbb{P}(X = x \cap Y = y) \end{aligned}$$

And,

$$f_X(x)f_Y(y) = \mathbb{P}(X = x)\mathbb{P}(Y = y)$$

From our assumption, these are equal, so  $\mathbb{P}(X = x \cap Y = y) = \mathbb{P}(X = x)\mathbb{P}(Y = y)$  which shows that  $X$  and  $Y$  are independent.

By implication both ways, the statement is proved.  $\square$

## 2.6

Let  $X$  have distribution  $F$  and density  $f$ , and let  $A$  be a subset of the real line, e.g.  $A = (a, b)$  for some  $a, b \in \mathbb{R}$  and  $a < b$ . We have the indicator function

$$I_A(x) = \begin{cases} 1 & x \in A \\ 0 & x \notin A \end{cases}$$

We will set  $Y = I_A(X)$  and find the PDF and CDF of  $Y$ .


The exercise asks for a probability mass function, but that cannot be correct. Since  $X$  has a density  $f$ , it is a continuous RV. If  $X \sim U(0, 1)$  and  $A = (0, 1)$ , then  $Y = X$  and it will be a uniform variable with a continuous distribution, so not necessarily discrete.

And if it can be a continuous distribution, what happens if we define  $A = \mathbb{Q} \subset \mathbb{R}$ ? There will be an infinite number of points in any interval with measure 0. Then we cannot define a PDF at all... Poorly formulated exercise in my opinion! Need to fill up with extra assumptions?

Skipping for now.

## 2.7

Let  $X$  and  $Y$  be independent and suppose that  $X, Y \sim U(0, 1)$ . For  $Z = \min(X, Y)$  we will find the density  $f_Z(z)$  for  $Z$ . Following the hint, we will first find  $\mathbb{P}(Z > z)$ . Since any observations of  $x, y \in (0, 1)$ , then we can immediately see that  $\mathbb{P}(Z > 0) = 1$  and  $\mathbb{P}(Z > 1) = 0$ . But what happens for other values? Best way to find out is with some illustrations. Here are plots of the cases  $\mathbb{P}(Z > 1/2)$ ,  $\mathbb{P}(Z > 1/4)$  and  $\mathbb{P}(Z > 3/4)$ .



If we simulate lots of  $X$  and  $Y$  values, we see that about 1/4th of them will have both  $X$  and  $Y$  values larger than 1/2, so  $\mathbb{P}(Z > 1/2) = 1/4$ . Similarly, we get  $\mathbb{P}(Z > 1/4) = 9/16$  and  $\mathbb{P}(Z > 3/4) = 1/16$ . Confirming this with a simulation.

```
# 2.7 - Simulating U(0,1)
N = 100000; X = runif(N); Y = runif(N)

Z = pmin(X, Y) # This is: Z = min{X, Y}

# Comparing simulated vs. theoretical results
sum(Z > 0.5)/N
1/4
sum(Z > 0.25)/N
9/16
sum(Z > 0.75)/N
1/16
```

```
> # Comparing simulated vs. theoretical results
> sum(Z > 0.5)/N
[1] 0.24888
> 1/4
[1] 0.25

> sum(Z > 0.25)/N
[1] 0.56156
> 9/16
[1] 0.5625

> sum(Z > 0.75)/N
[1] 0.06205
> 1/16
[1] 0.0625
```



By inspecting the images on the previous page, we can determine the 'shape' of the probabilities. For  $Z > 1/4$  we remove the union of  $X \leq 1/4$  and  $Y \leq 1/4$ . We define  $A = \{X \leq z\}$  and  $B = \{Y \leq z\}$ , and can write the general case as:

$$\begin{aligned}\mathbb{P}(Z > z) &= 1 - \mathbb{P}(\{X \leq z\} \cup \{Y \leq z\}) \\ &= 1 - \mathbb{P}(A \cup B) \\ &= 1 - [\mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)] \\ &= 1 - [\mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A)\mathbb{P}(B)]\end{aligned}$$

Where we used Lemma 1.6, and the fact that  $X$  and  $Y$  are independent. By using the probability law of complements, we can find the expression for  $\mathbb{P}(Z \leq z)$ .

$$\mathbb{P}(Z \leq z) = \mathbb{P}(X \leq z) + \mathbb{P}(Y \leq z) - \mathbb{P}(X \leq z)\mathbb{P}(Y \leq z)$$

The CDF for a uniform distribution on  $U(a, b)$  is:

$$F(z) = \frac{z - a}{b - a} \implies F_X(z) = F_Y(z) = \frac{z - 0}{1 - 0} = z$$

Which means:



$$F_Z(z) = F_X(z) + F_Y(z) - F_X(z)F_Y(z) = 2z - z^2$$

We can confirm our illustrations and simulated examples again by noting that:

$$\begin{aligned}F_Z(1/2) &= \frac{1}{2} + \frac{1}{2} - \left(\frac{1}{2} \cdot \frac{1}{2}\right) = 1 - \frac{1}{4} = \frac{3}{4} \\ F_Z(1/4) &= \frac{1}{4} + \frac{1}{4} - \left(\frac{1}{4} \cdot \frac{1}{4}\right) = \frac{8}{16} - \frac{1}{16} = \frac{7}{16} \\ F_Z(3/4) &= \frac{3}{4} + \frac{3}{4} - \left(\frac{3}{4} \cdot \frac{3}{4}\right) = \frac{24}{16} - \frac{9}{16} = \frac{15}{16}\end{aligned}$$

which gives us the opposite results as expected (since we simulated and illustrated  $\mathbb{P}(Z > z)$ ). The PDF  $f_Z(z)$  is the derivative of  $F_Z(z)$ . Including PDF-plot and histogram of the simulated  $Z$ s.

$$f_Z(z) = \frac{d}{dz}(F_Z(z)) = 2 - 2z.$$



## 2.8

The RV  $X$  has CDF  $F$ . Finding the CDF of  $X^+ = \max\{0, X\}$ .

From the definition of CDF:

$$\begin{aligned} F_{X^+}(u) &= \mathbb{P}(\max(0, X) \leq u) \\ &= \mathbb{P}(\{\omega \in \Omega : X(\omega) \leq u \text{ and } u \geq 0\}) \end{aligned}$$

We must consider two cases. When  $u < 0$ :

$$\begin{aligned} F_{X^+}(u) &= \mathbb{P}(\{\omega \in \Omega : X(\omega) \leq u \text{ and } u \geq 0\}) \\ &= \mathbb{P}(\emptyset) \\ &= 0 \end{aligned}$$

When  $u \geq 0$ :

$$\begin{aligned} F_{X^+}(u) &= \mathbb{P}(\{\omega \in \Omega : X(\omega) \leq u \text{ and } u \geq 0\}) \\ &= \mathbb{P}(\{\omega \in \Omega : X(\omega) \leq u\}) \\ &= \mathbb{P}(X \leq u) \\ &= F_X(u) \end{aligned}$$

So in summary:

$$F_{X^+}(u) = \begin{cases} 0 & u < 0 \\ F_X(u) & u \geq 0 \end{cases}$$

## 2.9

We have  $X \sim \text{Exp}(\beta)$ . The PDF is given by:

$$f(x) = \frac{1}{\beta} e^{-x/\beta}, \quad x > 0$$

Finding the CDF by integrating the PDF.

$$\begin{aligned} F(y) &= \int_{-\infty}^y f(x) dx \\ &= \frac{1}{\beta} \int_0^y e^{-x/\beta} dx \\ &= \frac{1}{\beta} [-\beta e^{-x/\beta}]_0^y \\ &= [-e^{-x/\beta}]_0^y \\ &= 1 - e^{-y/\beta} \end{aligned}$$

To find the inverse  $F^{-1}(q)$  we set  $q = F(y)$  and solve for  $y$ .

$$q = 1 - e^{-y/\beta} \implies y = -\beta \log(1 - q) \implies F^{-1}(q) = -\beta \log(1 - q)$$

### 2.10

If  $X$  and  $Y$  are independent, then  $g(X)$  and  $h(Y)$  are independent for some functions  $g$  and  $h$ .

PROOF. Let  $X$  and  $Y$  be some arbitrary random variables, and let  $x$  and  $y$  be values in the range of  $g$  and  $h$  such that  $g(X) = x$  and  $h(Y) = y$ . Then:

$$\mathbb{P}(g(X) = x, h(Y) = y) = \mathbb{P}(X = g^{-1}(x), Y = h^{-1}(y))$$

By independence of  $X$  and  $Y$ .

$$\begin{aligned} &= \mathbb{P}(X = g^{-1}(x))\mathbb{P}(Y = h^{-1}(y)) \\ &= \mathbb{P}(g(X) = x)\mathbb{P}(h(Y) = y) \end{aligned}$$

which shows that  $g$  and  $h$  satisfies the condition for independence.  $\square$

### 2.11

Tossing a coin which has probability  $p$  of getting H. We let  $X$  denote the number of heads and  $Y$  the number of tails.

(a) Showing that  $X$  and  $Y$  are dependent. First it will be helpful to consider a simplified case where we have  $N = 1$  and  $N = 2$  coin tosses, and assuming  $p = 1/2$ .

$N = 1$  Tosses

	$Y = 0$	$Y = 1$	
$X = 0$	0	1/2	1/2
$X = 1$	1/2	0	1/2
	1/2	1/2	1

$N = 2$  Tosses

	$Y = 0$	$Y = 1$	$Y = 2$	
$X = 0$	0	0	1/4	1/4
$X = 1$	0	1/2	0	1/2
$X = 2$	1/4	0	0	1/4
	1/4	1/2	1/4	1

In the case of  $N = 2$ , we see that  $f(0, 2) = 1/4$  while  $f_X(0)f_Y(2) = 1/16$ . This will be the inspiration for how we show it in the general case with  $N$  tosses and probability  $p$  of heads.

We only need to show one specific case where  $\mathbb{P}(X = x, Y = y) \neq \mathbb{P}(X = x)\mathbb{P}(Y = y)$  to show that these values are dependent. The total number of tosses will be  $N = X + Y$  and we compare the cases where we get  $N$  heads. In that case, using the Multinomial distribution:

$$\mathbb{P}(X = N, Y = 0) = \binom{N}{N, 0} = p^N(1 - p)^0 = p^N,$$

but with the two Binomial distributions:

$$\mathbb{P}(X = N)\mathbb{P}(Y = 0) = \binom{N}{N}p^N(1 - p)^0 \times \binom{N}{0}p^0(1 - p)^N = p^N(1 - p)^N.$$

These are not equal, showing that  $X$  and  $Y$  are dependent variables.

(b) Now we have  $N \sim \text{Poisson}(\lambda)$ , where  $N = X + Y$  for  $X$  heads and  $Y$  tails. Show that these values are now independent.

*Current solution, but not sure it's correct...*

Assuming  $X = x$  and  $Y = y$ . Then:

$$\begin{aligned}
 \mathbb{P}(X = x \cap Y = y) &= \mathbb{P}(X = x \cap Y = y \cap N = x + y) \\
 &= \mathbb{P}(X = x \cap Y = y \mid N = x + y) \cdot \mathbb{P}(N = x + y) \\
 &= \binom{x+y}{x} p^x (1-p)^y \cdot e^{-\lambda} \frac{\lambda^{x+y}}{(x+y)!} \\
 &= \frac{(x+y)!}{x!y!} p^x (1-p)^y \cdot e^{-\lambda} \frac{\lambda^x \lambda^y}{(x+y)!} \\
 &= e^{-\lambda} \frac{p^x \lambda^x}{x!} \cdot \frac{(1-p)^y \lambda^y}{y!} \\
 &= e^{-\lambda p} \frac{p^x \lambda^x}{x!} \cdot e^{-\lambda(1-p)} \frac{(1-p)^y \lambda^y}{y!} \\
 &= \mathbb{P}(X = x \mid \lambda p) \cdot \mathbb{P}(Y = y \mid \lambda(1-p)) \\
 &= \mathbb{P}(X = x) \mathbb{P}(Y = y)
 \end{aligned}$$

Which shows we have independence. We used:

$$e^{-\lambda} = e^{-\lambda(p-p+1)} = e^{-\lambda p + \lambda p - \lambda} = e^{-\lambda p} e^{\lambda p - \lambda} = e^{-\lambda p} e^{-\lambda(1-p)}.$$

Also, we made the assumption that  $\mathbb{P}(X = x \cap Y = y \mid N = x + y)$  is Binomial. But I think this is only true when we can assume that  $X$  and  $Y$  are independent... which is what we are trying to show. Will review this later... hopefully!

## 2.12 THEOREM 2.33

Suppose that the range of  $X$  and  $Y$  is a (possibly infinite) rectangle. If  $f(x, y) = g(x)h(y)$  for some functions  $g$  and  $h$  (not necessarily probability density functions) then  $X$  and  $Y$  are independent.

PROOF. From the joint PDF, we can find the marginal distributions:

$$f_X(x) = \int f(x, y) dy, \quad f_Y(y) = \int f(x, y) dx$$

By applying these integrals to both sides of the equality:

$$\begin{aligned}
 \int f(x, y) dy &= g(x) \int h(y) dy \implies f_X(x) = g(x) \\
 \int f(x, y) dx &= h(y) \int g(x) dx \implies f_Y(y) = h(y)
 \end{aligned}$$

We get  $g(x)$  and  $h(y)$  since they are equal to  $f(x, y)$  over the entire 'rectangle' and must therefore integrate to 1. This leaves us with:  $f(x, y) = g(x)h(y) = f_X(x)f_Y(y)$ , and by the results in exercise 2.5, this means that  $X$  and  $Y$  are independent.  $\square$

### 2.13


(a) Finding the PDF of  $Y = e^X$  when  $X \sim N(0, 1)$ .

$$\begin{aligned} F_Y(y) &= \mathbb{P}(Y \leq y) = \mathbb{P}(e^X \leq y) \\ &= \mathbb{P}(X \leq \log(y)) = F_X(\log(y)) \end{aligned}$$


We simply get the standard normal distribution with  $\log(y)$  as input. Differentiating to get the PDF:

$$f_Y(y) = \frac{d}{dy} F_X(\log(y)) = f_X(\log(y)) \cdot \frac{1}{y} = \frac{1}{y\sqrt{2\pi}} \exp\left(-\frac{1}{2}(\log(y))^2\right)$$

Plotting the function:



(b) Plotting histogram of simulated results. Comparable to the plot above.



```
x = rnorm(10000)
y = exp(x)
pdf("~/AllStatistics/files/ch2_2.13b.pdf",
     width = 4.7747, height = 4)
d = density(y)
hist(y, breaks = 200, xlim = c(0, 8),
     prob=TRUE)
lines(density(y), col="red", xlim=c(0,8))
dev.off()
```

## 2.14

We let  $(X, Y)$  be uniformly distributed on the unit disk:  $\{(x, y) : x^2 + y^2 \leq 1\}$ . Find the CDF and PDF of  $R = \sqrt{X^2 + Y^2}$ .

First, let's write some code in order to simulate the data, which are random points contained within the unit circle. And then use that to simulate  $R$  and view the histogram. Plots and code can be found on the next page. By inspection of the histogram it is clear that  $f_R(r) = 2r$  which means  $F_R(r) = r^2$ .

Following the general recipe for transformation of multiple random variables. We will define the function  $r(X, Y) = \sqrt{X^2 + Y^2}$ .

$$F_R(r) = \mathbb{P}(R \leq r) = \mathbb{P}(r(X, Y) \leq r) = \mathbb{P}(X^2 + Y^2 \leq r^2) = \int \int_{A_r} f_{X,Y}(x, y) dx dy$$

Finding the set  $A_r$  which in this case is still the unit circle, since  $\sqrt{x^2 + y^2} \leq 1 \implies x^2 + y^2 \leq 1$ .

Since the area of the unit circle is  $\pi$ , and since a uniform distribution means the probability is equal everywhere, it follows that:

$$f_{X,Y}(x, y) = \frac{1}{\pi}.$$

As we saw, we must integrate this function over the circle:  $\{(x, y) : x^2 + y^2 < r\}$  for some  $0 \leq r \leq 1$ . This is easiest when calculating the integral in polar coordinates, so we introduce  $0 \leq \theta \leq 2\pi$  and  $0 \leq \rho \leq r$  and change  $dx dy = \rho d\rho d\theta$ . So:

$$\begin{aligned} F_R(r) &= \frac{1}{\pi} \int_0^{2\pi} \int_0^r \rho d\rho d\theta \\ &= \frac{1}{\pi} \int_0^{2\pi} \left[ \frac{\rho^2}{2} \right]_0^r d\theta \\ &= \frac{1}{\pi} \int_0^{2\pi} \frac{r^2}{2} d\theta \\ &= \frac{r^2}{2\pi} \int_0^{2\pi} (1) d\theta \\ &= \frac{r^2}{2\pi} [\theta]_0^{2\pi} \\ &= \frac{r^2}{2\pi} \cdot 2\pi \\ &= r^2 \end{aligned}$$

The CDF is  $F_R(r) = r^2$  for  $0 \leq r \leq 1$ . By differentiating this, we get the PDF.


$$f_R(r) = \frac{d}{dr} F_R(r) = 2r,$$

also in the interval  $0 \leq r \leq 1$ . This confirms the results we found by simulating which are found on the next page.

## Plots and Code for 2.14


```
simulateUnitCircle = function(N) {  
  TR = runif(N)  
  UR = runif(N); VR = runif(N)  
  t = 2*pi*TR  
  u = UR+VR  
  u[u > 1] = 2 - u[u > 1]  
  r = u  
  X = r*cos(t)  
  Y = r*sin(t)  
  retVal = list()  
  retVal$X = X  
  retVal$Y = Y  
  return(retVal)  
}  
simVal = simulateUnitCircle(1000)  
X = simVal$X  
Y = simVal$Y  
pdf("~/AllStatistics/files/ch2_2.14.pdf",  
     width = 4, height = 4)  
plot(X, Y)  
dev.off()
```

Result from simulation:



```
simVal = simulateUnitCircle(50000)  
R = sqrt(simVal$X^2 + simVal$Y^2)  
pdf("~/AllStatistics/files/ch2_2.14b.pdf",  
     width = 4, height = 4)  
hist(R, breaks = 40, prob = TRUE)  
plot(X, Y)  
dev.off()
```

**Histogram of R**



## 2.15

Let  $X$  have a continuous, strictly increasing CDF  $F$ , and set  $Y = F(X)$ . Find the density of  $Y$ .

By definition of the CDF,  $F : \mathbb{R} \rightarrow [0, 1]$  which will be the domain of  $Y$ . Since  $F$  is strictly increasing, then  $F(X_1) < F(X_2)$  if  $X_1 < X_2$ , but the important property is that  $F$  must be *invertible*. For any  $y \in [0, 1]$  we get the following:

$$\begin{aligned} F_Y(y) &= \mathbb{P}(Y \leq y) = \mathbb{P}(F(X) \leq y) \\ &= \mathbb{P}(X \leq F^{-1}(y)) = F(F^{-1}(y)) = y \end{aligned}$$

Since  $F_Y(y) = y$ , then the density is  $f_Y(y) = 1$ , the derivative. From these observations, we can conclude that  $Y \sim U(0, 1)$ .

Next, if  $U \sim U(0, 1)$  and  $X = F^{-1}(U)$ , then  $X$  has  $F$  as its CDF.



$$\mathbb{P}(F^{-1}(U) \leq x) = \mathbb{P}(X \leq x) = F(x) \implies X \sim F$$

Writing a program that generates  $\text{Exp}(\beta)$  values. Using the result found in exercise 2.9.

```
simExp = function(beta, N){
  U = runif(N)
  return(-beta*log(1 - U))
}
# Simulating
expRND = simExp(beta = 2/3, N = 50000)
# Making plot
hist(expRND, breaks=80,
     xlim=c(0, 5), prob=TRUE)
lines(density(expRND), col="red", xlim=c(
  0,5), lwd=2)
```

Plot of the exponential distribution for  $\beta = 2/3$ .

$$f_X(x) = 1.5 \exp(-1.5x)$$





## 2.16

If  $X \sim \text{Poisson}(\lambda_1)$  and  $Y \sim \text{Poisson}(\lambda_2)$ , and  $X$  and  $Y$  are independent, then the distribution of  $X$  given that  $X + Y = n$  is  $\text{Binomial}(n, \rho)$  where  $\rho = \lambda_1/(\lambda_1 + \lambda_2)$ . (Changed some variable names).

PROOF. By definition of conditional probability, by applying hint 1, by independence and by using the fact that  $X + Y$  is a Poisson distribution with  $\lambda_1 + \lambda_2$ .

$$\begin{aligned}\mathbb{P}(X = x | X + Y = n) &= \frac{\mathbb{P}(X = x \cap X + Y = n)}{\mathbb{P}(X + Y = n)} \\ &= \frac{\mathbb{P}(X = x \cap Y = n - x)}{\mathbb{P}(X + Y = n)} && \text{(Hint)} \\ &= \frac{\mathbb{P}(X = x)\mathbb{P}(Y = n - x)}{\mathbb{P}(X + Y = n)} && \text{(Independence)}\end{aligned}$$

Each of these terms are known:

$$\begin{aligned}\mathbb{P}(X = x) &= e^{-\lambda_1} \cdot \frac{\lambda_1^x}{x!} \\ \mathbb{P}(Y = n - x) &= e^{-\lambda_2} \cdot \frac{\lambda_2^{(n-x)}}{(n-x)!} \\ \mathbb{P}(X + Y = n) &= e^{-(\lambda_1 + \lambda_2)} \cdot \frac{(\lambda_1 + \lambda_2)^n}{n!}\end{aligned}$$

We can invert the last equation:

$$\frac{1}{\mathbb{P}(X + Y = n)} = e^{\lambda_1 + \lambda_2} \cdot \frac{n!}{(\lambda_1 + \lambda_2)^n}$$

Putting it all together, we can complete the justification.

$$\begin{aligned}\mathbb{P}(X = x | X + Y = n) &= \frac{\mathbb{P}(X = x)\mathbb{P}(Y = n - x)}{\mathbb{P}(X + Y = n)} \\ &= \left( e^{-\lambda_1} \cdot \frac{\lambda_1^x}{x!} \right) \left( e^{-\lambda_2} \cdot \frac{\lambda_2^{(n-x)}}{(n-x)!} \right) \left( e^{\lambda_1 + \lambda_2} \cdot \frac{n!}{(\lambda_1 + \lambda_2)^n} \right) \\ &= \left( \frac{\lambda_1^x}{x!} \right) \left( \frac{\lambda_2^{(n-x)}}{(n-x)!} \right) \left( \frac{n!}{(\lambda_1 + \lambda_2)^x (\lambda_1 + \lambda_2)^{(n-x)}} \right) \\ &= \left( \frac{n!}{x!(n-x)!} \right) \left( \frac{\lambda_1}{\lambda_1 + \lambda_2} \right)^x \left( \frac{\lambda_2}{\lambda_1 + \lambda_2} \right)^{(n-x)} \\ &= \binom{n}{x} \left( \frac{\lambda_1}{\lambda_1 + \lambda_2} \right)^x \left( \frac{\lambda_2}{\lambda_1 + \lambda_2} \right)^{(n-x)} \\ &= \binom{n}{x} \left( \frac{\lambda_1}{\lambda_1 + \lambda_2} \right)^x \left( 1 - \frac{\lambda_1}{\lambda_1 + \lambda_2} \right)^{(n-x)} \\ &= \binom{n}{x} \rho^x (1 - \rho)^{(n-x)} \sim \text{Binomial}(n, \rho)\end{aligned}$$

for  $\rho = \lambda_1/(\lambda_1 + \lambda_2)$ . □

**2.17**

The joint probability density is:

$$f_{X,Y}(x,y) = \begin{cases} c(x+y^2) & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Calculating  $\mathbb{P}(X < 1/2 | Y = 1/2)$ .

We begin by calculating the marginal distribution for  $f_Y(y)$ .

$$f_Y(y) = \int_0^1 cx + cy^2 dx = c \int_0^1 x dx + c \int_0^1 y^2 dx = c \left[ \frac{x^2}{2} \right]_0^1 + cy^2 [x]_0^1 = c \left( \frac{1}{2} + y^2 \right)$$

Following the steps in example 2.38.

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)} = \frac{c(x+y^2)}{c(\frac{1}{2} + y^2)} = \frac{x+y^2}{\frac{1}{2} + y^2}$$

Calculating the conditional probability:

$$\begin{aligned} \mathbb{P}(X < 1/2 | Y = 1/2) &= \int_0^{1/2} f_{X|Y}(1/2|1/2) dx \\ &= \int_0^{1/2} \frac{x + \frac{1}{4}}{\frac{1}{2} + \frac{1}{4}} dx \\ &= \frac{4}{3} \int_0^{1/2} x + \frac{1}{4} dx \\ &= \frac{4}{3} \left[ \frac{x^2}{2} + \frac{x}{4} \right]_0^{1/2} = \frac{4}{3} \cdot \frac{1}{4} = \frac{1}{3} \end{aligned}$$

**2.18**

Let  $X \sim N(3, 16)$ . Solve the following using the Normal table and using a computer package.

(a)

$$\mathbb{P}(X < 7) = \mathbb{P}(3 + 4Z < 7) = \mathbb{P}\left(Z < \frac{7-3}{4}\right) = \mathbb{P}(Z < 1) = 0.8413$$

From **R**:

```
> pnorm(1)           > 1 - pnorm(-0.25)
[1] 0.84134475         [1] 0.59870633
```

(b)

$$\begin{aligned} \mathbb{P}(X > 2) &= 1 - \mathbb{P}(X \leq 2) = 1 - \mathbb{P}(3 + 4Z \leq 2) = 1 - \mathbb{P}\left(Z \leq \frac{2-3}{4}\right) \\ &= 1 - \mathbb{P}(Z < -1/4) = 1 - (1 - 0.5987) = 0.5987 \end{aligned}$$

(c) Find  $x$  such that  $\mathbb{P}(X > x) = 0.05$ . By finding 0.95 in the table, we see that it is generated by the value 1.645, so:

$$\mathbb{P}(Z \leq 1.645) = 0.95 \implies \mathbb{P}(Z > 1.645) = 1 - 0.95 = 0.05$$

$$\mathbb{P}(4Z > 6.58) = \mathbb{P}(3 + 4Z > 9.58) = \mathbb{P}(X > 9.58) = 0.05$$

So,  $x = 9.58$ . From **R**:

```
1 - pnorm(9.5794, mean = 3, sd = 4)
[1] 0.05000037
```

(d)

$$\begin{aligned} \mathbb{P}(0 \leq X < 4) &= \mathbb{P}(0 \leq 3 + 4Z < 4) = \mathbb{P}(-3 \leq 4Z < 1) \\ &= \mathbb{P}(-3/4 \leq Z < 1/4) = \mathbb{P}(Z < 1/4) - \mathbb{P}(Z < -3/4) \\ &= 0.5987 - (1 - 0.7734) = 0.5987 - 0.2266 \\ &= 0.3721 \end{aligned}$$

From **R**:

```
> pnorm(0.25) - pnorm(-0.75)
[1] 0.37207897
#### Alternatively
> CH = rnorm(10000000, mean = 3, sd = 4)
> sum(CH < 4 & CH > 0)/10000000
[1] 0.3719199
```

(e) Find  $x$  such that  $\mathbb{P}(|X| > |x|) = 0.05$ .

We can split this up in two sections if we assume  $x > 0$ . Using that the sets are disjoint:

$$\begin{aligned} \mathbb{P}(|X| > |x|) &= \mathbb{P}(X > x \cap X < -x) = \mathbb{P}(X > x) + \mathbb{P}(X < -x) \\ &= 1 - \mathbb{P}(X \leq x) + \mathbb{P}(X < -x) = 0.05 \end{aligned}$$

$$\mathbb{P}(X \leq x) - \mathbb{P}(X < -x) = 0.95$$

Note we get minus. Translating to standard normal:

$$\mathbb{P}\left(Z \leq \frac{x-3}{4}\right) - \mathbb{P}\left(Z < \frac{-x-3}{4}\right) = 0.95$$

I am not able to see how to solve this by just looking up values because of the dependence. By using R, we can find that  $x = 9.611$  which makes the lookup values 1.65275 and -3.15275. Verifying in **R**.

```
> X = rnorm(10000000, mean = 3, sd = 4)
> x = 9.611
> sum(X < -x | X > x)/10000000
[1] 0.0500769
> sum(abs(X) > abs(x))/10000000
[1] 0.0500769
```

### 2.19

Proving equation (2.12). Let  $X$  be a random variable with pdf  $f_X(x)$ , and define the random variable  $Y = r(X)$ , where  $r$  is either strictly monotone increasing or decreasing, so  $r(\cdot)$  has the inverse function  $s(\cdot) = r^{-1}(\cdot)$ . Then, we can express the pdf of  $Y$  as:

$$f_Y(y) = f_X(s(y)) \left| \frac{ds(y)}{dy} \right|.$$

PROOF. The proof will be in two parts. Assume first that  $r$  is strictly increasing, which means that  $s$  is strictly increasing and its derivative is positive. From the CDF of  $Y$ , we get:

$$\begin{aligned} F_Y(y) &= \mathbb{P}(Y \leq y) \\ &= \mathbb{P}(r(X) \leq y) \\ &= \mathbb{P}(X \leq r^{-1}(y)) \\ &= \mathbb{P}(X \leq s(y)) \\ &= F_X(s(y)) \end{aligned}$$

Differentiating the CDF to get the PDF.

$$f_Y(y) = \frac{d}{dy} F_Y(y) = \frac{d}{dy} F_X(s(y)) = f_X(s(y)) \frac{ds(y)}{dy}$$

Now assume that  $r$  is strictly decreasing, which means  $s$  is strictly decreasing and thus has a negative derivative. From the CDF of  $Y$ , we now get:

$$\begin{aligned} F_Y(y) &= \mathbb{P}(Y \leq y) \\ &= \mathbb{P}(r(X) \leq y) \\ &= \mathbb{P}(X \geq r^{-1}(y)) \\ &= 1 - \mathbb{P}(X \leq s(y)) \\ &= 1 - F_X(s(y)) \end{aligned}$$

$$f_Y(y) = \frac{d}{dy} F_Y(y) = \frac{d}{dy} [1 - F_X(s(y))] = -f_X(s(y)) \frac{ds(y)}{dy} = f_X(s(y)) \left( -\frac{ds(y)}{dy} \right)$$

Since the derivative will be negative, the negative of the derivative will be positive. Generalizing to both cases, we can write:

$$f_Y(y) = f_X(s(y)) \left| \frac{ds(y)}{dy} \right| \quad \square$$

A short note on why we get  $1 - \mathbb{P}(X \leq s(y))$  when  $r$  is strictly decreasing. As an example, say that  $X \sim U(0, 5)$  and we define  $Y = r(X)$  where  $r(x) = 2 - 2x$ , which is strictly decreasing. Then  $r^{-1}(y) = s(y) = 1 - \frac{y}{2}$ , which is also strictly decreasing. Then:

$$\mathbb{P}(r(X) \leq y) = \mathbb{P}(2 - 2X \leq y) = \mathbb{P}(2 - y \leq 2X) = \mathbb{P}(1 - \frac{y}{2} \leq X) = \mathbb{P}(X \geq s(y))$$

In order to get the expression in terms of  $F_X(x)$ , we rewrite it:

$$\mathbb{P}(X \geq s(y)) = 1 - \mathbb{P}(X \leq s(y)) = 1 - F_X(s(y)).$$

## 2.20

Let  $X, Y \sim U(0, 1)$  be independent. Find the PDF for  $X - Y$  and  $X/Y$ . The joint density of  $X, Y$  is

$$f(x, y) = 1, \quad x, y \in (0, 1)$$

and 0 elsewhere.

We define  $Z = X - Y$ . We note that  $Z$  can assume values between -1 and 1, which are the two extremes. This gives us  $A_z = (-1, 1)$ . We split this into two cases: when  $X > Y$ , then  $A_z = (0, 1)$  and when  $X < Y$ , which means  $A_z = (-1, 0)$ .

When  $X > Y$  and  $A_z = (0, 1)$ .



$$\begin{aligned} F_Z(z) &= \int_0^z \int_y^1 1 dx dy = \int_0^z [x]_y^1 dy \\ &= \int_0^z 1 - y dy = \left[ y - \frac{y^2}{2} \right]_0^z \\ &= z - \frac{z^2}{2} \end{aligned}$$

When  $X < Y$  and  $A_z = (-1, 0)$ .

$$\begin{aligned} F_Z(z) &= \int_{-1}^z \int_{-1}^y 1 dx dy = \int_{-1}^z [x]_{-1}^y dy \\ &= \int_{-1}^z y + 1 dy = \left[ \frac{y^2}{2} + y \right]_{-1}^z \\ &= \frac{z^2}{2} + z + \frac{1}{2} \end{aligned}$$

Summarizing the CDF, and differentiating to find the PDF. (The 1/2 carries over).

$$F_Z(z) = \begin{cases} \frac{z^2}{2} + z + \frac{1}{2} & z \in (-1, 0) \\ z - \frac{z^2}{2} + \frac{1}{2} & z \in (0, 1) \end{cases} \implies f_Z(z) = \begin{cases} z + 1 & z \in (-1, 0) \\ 1 - z & z \in (0, 1) \end{cases}$$



Now we define  $Z = X/Y$ .



$$F_Z(z) = \mathbb{P}(Z \leq z) = \mathbb{P}(X/Y \leq z) = \mathbb{P}(X \leq zY)$$

$$F_Z(z) = \int_0^1 \int_0^{\min(z,y)} 1 dx dy = \int_0^1 \min(z,y) dy = \begin{cases} \int_0^{\frac{1}{z}} zy dy + \int_{\frac{1}{z}}^1 1 dy & z \geq 1 \\ \int_0^1 zy dy & z < 1 \end{cases}$$

Evaluating the integrals to get the CDF, and differentiating to get the PDF:

$$F_Z(z) = \begin{cases} 1 - \frac{1}{2z} & z \geq 1 \\ \frac{z}{2} & z < 1 \end{cases} \implies f_Z(z) = \begin{cases} \frac{1}{2z^2} & z \geq 1 \\ \frac{1}{2} & z < 1 \end{cases}$$

Plots and simulation code.



```
# Simulating X-Y
N = 10000
X = runif(N); Y = runif(N)
d1 = X - Y
hist(d1, breaks = 50, prob = TRUE,
     main="Histogram of X-Y")

# Simulating X/Y
N = 1000000
X = runif(N); Y = runif(N)
d2 = X/Y
d3 = d2[abs(d2) < 3] # Removing very rare large vals
d3 = d3*length(d2)/length(d3) # Normalizing probability
hist(d3, breaks=60, prob=TRUE,
     main = "Histogram of X/Y")
```

R

### 2.21

Let  $X_1, \dots, X_n \sim \text{Exp}(\beta)$  be IID. Define  $Y = \max(X_1, \dots, X_n)$ . Find the PDF of  $Y$ .

Recall the PDF of the exponential distribution:

$$f_X(x) = \frac{1}{\beta} \exp(-x/\beta), \quad x \geq 0$$

Defining  $r(X_1, \dots, X_n) = \max(X_1, \dots, X_n)$ . Finding an expression for the CDF of  $Y$ .

$$F_Y(y) = \mathbb{P}(Y \leq y) = \mathbb{P}(\max(X_1, \dots, X_n) \leq y)$$


According to the hint,  $Y \leq y$  iff  $X_i \leq y$  for all  $i \in I$ , where  $I = \{1, \dots, n\}$  is the index set. We can therefore express the CDF of  $Y$  in terms of each individual CDF:

$$F_Y(y) = \mathbb{P}(X_1 \leq y) \cap \mathbb{P}(X_2 \leq y) \cap \dots \cap \mathbb{P}(X_n \leq y) = (F_X(y))^n = (1 - \exp(-y/\beta))^n$$

Differentiating wrt.  $y$  to find the PDF.

$$\begin{aligned} f_Y(y) &= \frac{d}{dy} F_Y(y) = \frac{d}{dy} (F_X(y))^n = n(F_X(y))^{n-1} f_X(y) \\ &= n(1 - \exp(-y/\beta))^{n-1} \frac{1}{\beta} \exp(-y/\beta) \\ &= \frac{n}{\beta} \exp(-y/\beta) (1 - \exp(-y/\beta))^{n-1} \end{aligned}$$

Plots and simulation code for  $n = 10$  and  $\beta = 0.2$ .



```
N = 100000
lambda = 5

exp1 = rexp(N, rate = lambda)
exp2 = rexp(N, rate = lambda)
exp3 = rexp(N, rate = lambda)
exp4 = rexp(N, rate = lambda)
exp5 = rexp(N, rate = lambda)
exp6 = rexp(N, rate = lambda)
exp7 = rexp(N, rate = lambda)
exp8 = rexp(N, rate = lambda)
exp9 = rexp(N, rate = lambda)
exp10 = rexp(N, rate = lambda)

Y = pmax(exp1, exp2, exp3, exp4, exp5,
         exp6, exp7, exp8, exp9, exp10)

hist(Y, breaks = 50, prob = TRUE)
lines(density(Y), col="red", xlim=c(0,2.5), lwd=2)
```

R



### 3 Chapter 3 - Expectation

#### Exercises

##### 3.1

Define  $X$  as the wealth after  $n$  games. The probability of winning and losing is the same for each outcome so  $p = 1/2$ .

$$\mathbb{E}[X] = \frac{1}{2} \cdot 2c + \frac{1}{2} \cdot \left(\frac{1}{2}\right) c = c + \frac{c}{4} = \frac{5}{4}c$$

We expect to have  $5/4 \cdot c$  after  $n$  games. We can also verify this result with a simulation in **R**.

```
> games = sample(c(2, 0.5), size = 1000000, replace = TRUE)
> mean(games)
[1] 1.250973
> 5/4
[1] 1.25
```

R

##### 3.2

**Claim.**  $\text{Var}(X) = 0$  if and only if  $\mathbb{P}(X = c) = 1$  for some constant  $c$ .

PROOF.

$\Rightarrow$ ) Set  $c = \mu$  and assume  $\text{Var}(X) = 0$ , which means that

$$\mathbb{E}[(X - \mu)^2] = 0 \implies \int (x - \mu)^2 dF(x) = 0.$$

This can only be 0 when  $x = \mu = c$  for the entire domain of  $X$ . Hence  $\mathbb{P}(X = c) = 1$ .

$\Leftarrow$ ) Assume  $\mathbb{P}(X = c) = 1$ . When calculating the expectation:

$$\mu = \mathbb{E}[X] = \int c dF(x) = c$$

When calculating the variance:

$$\text{Var}(X) = \int (x - \mu)^2 dF(x) = \int (c - c)^2 dF(x) = 0,$$

since  $x = c$  for all  $x$  in the domain of  $X$ . □

### 3.3

Let  $X_1, \dots, X_n \sim U(0, 1)$  and define  $Y = \max(X_1, \dots, X_n)$ . We will calculate  $\mathbb{E}[Y]$ . It is not stated in the exercise, but we will assume that the  $X_i$  are independent. Finding the CDF for  $Y$ .

$$\begin{aligned} F_Y(y) &= \mathbb{P}(Y \leq y) \\ &= \mathbb{P}(\max(X_1, \dots, X_n) \leq y) \\ &= \mathbb{P}(X_1 \leq y) \cap \dots \cap \mathbb{P}(X_n \leq y) \\ &= \mathbb{P}(X_1 \leq y) \mathbb{P}(X_2 \leq y) \cdots \mathbb{P}(X_n \leq y) && \text{(Independence)} \\ &= (F_X(y))^n \end{aligned}$$

Differentiating to get the PDF for  $Y$ .

$$f_Y(y) = \frac{d}{dy} F_Y(y) = \frac{d}{dy} (F_X(y))^n = n(F_X(y))^{n-1} f_X(y)$$

Since  $X_i$  are uniformly distributed, we know that  $F_X(y) = y$  and  $f_X(y) = 1$ , so:

$$f_Y(y) = ny^{n-1}$$

Now we can calculate the expectation of  $Y$ .

$$\mathbb{E}[Y] = \int_0^1 y \cdot ny^{n-1} dy = n \int_0^1 y^n dy = n \left[ \frac{y^{n+1}}{n+1} \right]_0^1 = \frac{n}{n+1}$$

Confirming this result with a numeric simulation in **R**.

```
> # 3.3
> N = 1000000
> U1 = runif(N)
> U2 = runif(N)
> U3 = runif(N)
> U4 = runif(N)
> U5 = runif(N)
> U6 = runif(N)
> U7 = runif(N)
> U8 = runif(N)
> U9 = runif(N)
> U10 = runif(N)
> Y = pmax(U1, U2, U3, U4, U5,
+         U6, U7, U8, U9, U10)
> mean(Y)
[1] 0.9091151
> # Theoretical Result
> 10/11
[1] 0.9090909
```

R

As we can see, the theoretical result is very close to the simulated result for  $n = 10$ .

### 3.4 - Random Walk

A particle starts in the origin and jumps left, a step of -1, with probability  $p$  and jumps right, a step of 1, with probability  $1 - p$ . The expected location will be:

$$\mathbb{E}[X] = (-1)p + (1)(1 - p) = -p + 1 - p = 1 - 2p$$

To calculate the variance, we start by finding the second moment:

$$\mathbb{E}[X^2] = (-1)^2p + (1)^2(1 - p) = p + 1 - p = 1$$

So the variance is:

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = 1 - (1 - 2p)^2 = 1 - (1 - 4p + 4p^2) = 4p - 4p^2$$

### 3.5

Tossing a fair coin until we get H. Finding the expected number of tosses. The reasoning is as follows. We get H on the first toss with probability  $1/2$ , first H on the second toss with probability  $1/2^2 = 1/4$  and so on. The pattern becomes as follows, for the first 7 cases:

Tosses	Outcome	Probability
1	{H}	1/2
2	{TH}	1/4
3	{TTH}	1/8
4	{TTTH}	1/16
5	{TTTTH}	1/32
6	{TTTTTH}	1/64
7	{TTTTTTH}	1/128

Define  $T$  to be the number of tosses to get H.

$$\begin{aligned}\mathbb{E}[T] &= (1) \left(\frac{1}{2}\right) + (2) \left(\frac{1}{4}\right) + \dots + (k) \left(\frac{1}{2^k}\right) + \dots \\ &= \sum_{k=1}^{\infty} \frac{k}{2^k} \\ &= 2\end{aligned}$$

Not delving in to the mathematics of the infinite sum, but it can be shown that this sum becomes 2 which will be the expected number of tosses to get a H. Here is a numeric approximation in **R**.

```
> sumApprox = 0
> for (k in 1:1000) {
+   sumApprox = sumApprox + k/2^k
+ }
> sumApprox
[1] 2
```

R

### 3.6 Theorem - The Rule of the Lazy Statistician

Proving the following result for the discrete case. Let  $Y = r(X)$ , then

$$\mathbb{E}[Y] = \mathbb{E}[r(X)] = \sum_x r(x) f_X(x)$$

PROOF. Set  $\mathcal{X}, \mathcal{Y}$  to be the set of all values for  $X$  and  $Y$ . We have the functions  $r : \mathcal{X} \rightarrow \mathcal{Y}$  and  $r^{-1} : \mathcal{Y} \rightarrow \mathcal{X}$ . We will define  $s := r^{-1}$  for convenience. By definition of the expectation:

$$\begin{aligned} \mathbb{E}[Y] &= \sum_{y \in \mathcal{Y}} y \cdot f_Y(y) = \sum_{y \in \mathcal{Y}} y \cdot \mathbb{P}(Y = y) \\ &= \sum_{y \in \mathcal{Y}} y \sum_{x \in s(y)} \mathbb{P}(X = x) \\ &= \sum_{y \in \mathcal{Y}} \sum_{x \in s(y)} y \mathbb{P}(X = x) \\ &= \sum_{x \in \mathcal{X}} r(x) \mathbb{P}(X = x) \\ &= \sum_{x \in \mathcal{X}} r(x) f_X(x) \end{aligned} \quad \square$$

A simplified case to make the proof easier a bit easier to understand. We define the variables  $X$  and  $Y = r(X) = X^2$ , and use the following distributions.

$x$	$\mathbb{P}(X = x)$		$y$	$\mathbb{P}(Y = y)$
-1	1/4	$Y = r(X) = X^2$	0	1/4
0	1/4		1	3/4
1	1/2			

Then the calculations above become:

$$\begin{aligned} \mathbb{E}[Y] &= \sum_{y \in \mathcal{Y}} y \cdot f_Y(y) = (0)(1/4) + (1)(3/4) \\ &= \sum_{y \in \mathcal{Y}} y \sum_{x \in s(y)} \mathbb{P}(X = x) = (0)(1/4) + (1)[1/4 + 1/2] \\ &= \sum_{y \in \mathcal{Y}} \sum_{x \in s(y)} y \mathbb{P}(X = x) = (0)(1/4) + (1)(1/4) + (1)(1/2) \\ &= \sum_{x \in \mathcal{X}} r(x) \mathbb{P}(X = x) = (0)^2(1/4) + (-1)^2(1/4) + (1)^2(1/2) \\ &= \sum_{x \in \mathcal{X}} r(x) f_X(x) \end{aligned}$$

In both cases, the expectation becomes 3/4.

### 3.7

$X \sim F$  is continuous and we suppose that  $\mathbb{P}(X > 0) = 1$  and that  $\mathbb{E}[X]$  exists. Show that  $\mathbb{E}[X] = \int_0^\infty \mathbb{P}(X > x)dx$ .

PROOF. By definition of the expectation, and since the domain of  $X$  must be  $(0, \infty)$ :

$$\mathbb{E}[X] = \int_0^\infty x \cdot f_X(x)dx$$

Using integration by parts, and setting  $u = x$  and  $v' = f_X$ , which gives  $u' = 1$  and  $v = F_X$ :

$$\int uv' = uv - \int u'v,$$

which gives us:

$$\begin{aligned} \mathbb{E}[X] &= \int_0^\infty x \cdot f_X(x)dx = xF_X(x) \Big|_0^\infty - \int_0^\infty F_X(x)dx \\ &= xF_X(x) \Big|_0^\infty - \int_0^\infty \mathbb{P}(X \leq x)dx \\ &= xF_X(x) \Big|_0^\infty - \int_0^\infty 1 - \mathbb{P}(X > x)dx \\ &= xF_X(x) \Big|_0^\infty - x \Big|_0^\infty + \int_0^\infty \mathbb{P}(X > x)dx \\ &= -\left( \lim_{x \rightarrow \infty} x(1 - F_X(x)) \right) + \int_0^\infty \mathbb{P}(X > x)dx \\ &= \int_0^\infty \mathbb{P}(X > x)dx \end{aligned}$$

which proves the result. In the second to last step, we applied the hint regarding the limit.  $\square$

### 3.8 - Theorem 3.17

Let  $X_1, \dots, X_n$  be IID and let  $\mu = \mathbb{E}[X_i]$ ,  $\sigma^2 = \text{Var}(X_i)$ . Then:

$$\mathbb{E}[\bar{X}] = \mu, \quad \text{Var}(\bar{X}) = \frac{\sigma^2}{n}, \quad \mathbb{E}[S^2] = \sigma^2.$$

PROOF. Start with the expectation of the sample mean:

$$\mathbb{E}[\bar{X}] = \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n X_i \right] = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[X_i] = \frac{1}{n} \sum_{i=1}^n \mu = \frac{n \cdot \mu}{n} = \mu.$$

Next, the variance.

$$\text{Var}[\bar{X}] = \text{Var} \left[ \frac{1}{n} \sum_{i=1}^n X_i \right] = \frac{1}{n^2} \sum_{i=1}^n \text{Var}[X_i] = \frac{1}{n^2} \sum_{i=1}^n \sigma^2 = \frac{n \cdot \sigma^2}{n^2} = \frac{\sigma^2}{n}.$$

Finally, the expectation of the sample variance. This calculation is a lot more involved.

$$\mathbb{E}[S^2] = \mathbb{E} \left[ \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 \right] \implies (n-1)\mathbb{E}[S^2] = \mathbb{E} \left[ \sum_{i=1}^n (X_i - \bar{X})^2 \right]$$

We will use that  $\bar{X} = (1/n)\sum X_i$  means that  $\sum X_i = n\bar{X}$ . Running through the calculations:

$$\begin{aligned} (n-1)\mathbb{E}[S^2] &= \mathbb{E} \left[ \sum_{i=1}^n (X_i^2 - 2X_i\bar{X} + \bar{X}^2) \right] \\ &= \mathbb{E} \left[ \sum_{i=1}^n X_i^2 - 2 \sum_{i=1}^n X_i\bar{X} + \sum_{i=1}^n \bar{X}^2 \right] \\ &= \mathbb{E} \left[ \sum_{i=1}^n X_i^2 \right] - \mathbb{E} \left[ 2 \sum_{i=1}^n X_i\bar{X} \right] + \mathbb{E} \left[ \sum_{i=1}^n \bar{X}^2 \right] \\ &= \sum_{i=1}^n \mathbb{E} [X_i^2] - \mathbb{E} \left[ 2\bar{X} \sum_{i=1}^n X_i \right] + \mathbb{E} \left[ \bar{X}^2 \sum_{i=1}^n 1 \right] \quad (\bar{X} \text{ indep. of } i) \\ &= n\mathbb{E} [X_i^2] - 2n\mathbb{E} [\bar{X}^2] + n\mathbb{E} [\bar{X}^2] \quad (\sum X_i = n\bar{X}) \\ &= n\mathbb{E} [X_i^2] - n\mathbb{E} [\bar{X}^2] \end{aligned}$$

By dividing both sides of the equality by  $n$ , we have shown:

$$\frac{n-1}{n}\mathbb{E}[S^2] = \mathbb{E} [X_i^2] - \mathbb{E} [\bar{X}^2] \quad (3.8.1)$$

We have the second moment for  $X$ . From the definition of the variance:

$$\text{Var}(X_i) = \mathbb{E}[X_i^2] - \mu^2 \implies \mathbb{E}[X_i^2] = \text{Var}(X_i) + \mu^2 = \sigma^2 + \mu^2$$

From the previous results:

$$\text{Var}(\bar{X}) = \mathbb{E}[\bar{X}^2] - \mathbb{E}[\bar{X}]^2 \implies \mathbb{E}[\bar{X}^2] = \text{Var}(\bar{X}) + \mathbb{E}[\bar{X}]^2 = \frac{\sigma^2}{n} + \mu^2$$

Replacing each of these into (3.8.1) and isolating  $\mathbb{E}[S^2]$ .

$$\begin{aligned} \mathbb{E}[S^2] &= \frac{n}{n-1} \left( \sigma^2 + \mu^2 - \left( \frac{\sigma^2}{n} + \mu^2 \right) \right) \\ &= \frac{n}{n-1} \left( \sigma^2 - \frac{\sigma^2}{n} \right) \\ &= \frac{n}{n-1} \left( \frac{n-1}{n} \right) \sigma^2 \\ &= \sigma^2 \end{aligned}$$


and we have finally shown the final result. □

### 3.9

Computer experiment; studying the effects of the mean over 10.000 simulations from the standard normal distribution and the Cauchy distribution.

As we can see the simulation of the normal distribution quickly stabilizes to a value of around 0 which is the expected value. The Cauchy distribution is a very heavy-tailed distribution and the expectation does not exist. Therefore, there is no mean it can stabilize to and it ends up behaving erratically even after 10.000 simulations. The jumps that can be seen are extreme outliers that are added to the data, which are relatively common in the Cauchy distribution.

Results:



```
# Setting the number of simulations
N = 10000

# Simulating N standard normal values
X = rnorm(N)
Xmean = cumsum(X)/1:N
plot(1:N, Xmean, type="l", ylim = c(-0.2, 0.2),
main = "Mean of N(0,1)")
abline(h = 0, col="red")

# Simulating N Cauchy values with loc = 0, scale = 1
Y = rcauchy(N)
Ymean = cumsum(Y)/1:N
plot(1:N, Ymean, type="l",
main = "Mean of Cauchy")
```

R

### 3.10

Let  $X \sim N(0, 1)$  and define  $Y = e^X$ . Find  $\mathbb{E}[Y]$  and  $\text{Var}(Y)$ .

One possible way to solve this is defining  $r(X) = e^X$  and evaluating:

$$\mathbb{E}[Y] = \mathbb{E}[r(X)] = \int_{\mathbb{R}} e^x f_X(x) dx = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \exp\left(x - \frac{1}{2}x^2\right) dx.$$

Completing the square in the exponential term:

$$x - \frac{1}{2}x^2 = -\frac{1}{2}(x^2 - 2x) = -\frac{1}{2}(x^2 - 2x + 1 - 1) = -\frac{1}{2}(x - 1)^2 + \frac{1}{2}$$

We can rewrite the integral:

$$\mathbb{E}[Y] = e^{1/2} \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \exp\left(-\frac{1}{2}(x - 1)^2\right) dx$$

We will integrate this by substitution:

$$u = x - 1 \implies du/dx = 1 \implies du = dx$$

$$\mathbb{E}[Y] = e^{1/2} \underbrace{\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \exp\left(-\frac{1}{2}u^2\right) du}_{=1} = e^{1/2} \approx 1.6487$$

Confirming with numerical simulation.

```
> N = 10000000; X = rnorm(N); Y = exp(X)
> mean(Y)
[1] 1.648564
> exp(0.5)
[1] 1.648721
> var(Y)
[1] 4.676382
> exp(2) - exp(1)
[1] 4.670774
```

We can calculate the variance by first finding the second moment:  $\mathbb{E}[Y^2]$ .

$$\mathbb{E}[Y^2] = \mathbb{E}[r(X)^2] = \int_{\mathbb{R}} e^{2x} f_X(x) dx = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \exp\left(2x - \frac{1}{2}x^2\right) dx.$$

Just as above, we complete the square and get:  $-(1/2)(x - 2)^2 + 2$  and move  $e^2$  outside the integral. Then we use integration by substitution:  $u = x - 2$  which gives  $du = dx$ . We end up with a similar integral:

$$\mathbb{E}[Y^2] = e^2 \underbrace{\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \exp\left(-\frac{1}{2}u^2\right) du}_{=1} = e^2$$

Now we can calculate the variance of  $Y$ .

$$\text{Var}(Y) = \mathbb{E}[Y^2] - \mathbb{E}[Y]^2 = e^2 - (e^{1/2})^2 = e^2 - e^1,$$

and as we can see in the simulation above, this corresponds to the simulated variance.



### 3.11

Simulating stocks. For each day  $Y_i$  can be either  $-1$  or  $1$ , each with probability  $1/2$ . We have  $X_n = \sum_{i=1}^n Y_i$  which is the cumulative value of the stock.

(a) Calculating the expectation and variance of  $X_n$ . First we calculate it for  $Y_i$ .

$$\begin{aligned}\mathbb{E}[Y_i] &= (-1)(1/2) + (1)(1/2) = -1/2 + 1/2 = 0 \\ \mathbb{E}[Y_i^2] &= (-1)^2(1/2) + (1)^2(1/2) = 1/2 + 1/2 = 1 \\ \text{Var}(Y_i) &= \mathbb{E}[Y_i^2] - \mathbb{E}[Y_i]^2 = 1 - 0 = 1\end{aligned}$$


Now for  $X_n$ .

$$\begin{aligned}\mathbb{E}[X_n] &= \mathbb{E}\left[\sum_{i=1}^n Y_i\right] = \sum_{i=1}^n \mathbb{E}[Y_i] = 0 \\ \text{Var}(X_n) &= \text{Var}\left[\sum_{i=1}^n Y_i\right] = \sum_{i=1}^n \text{Var}[Y_i] = \sum_{i=1}^n 1 = n\end{aligned}$$

(b) Simulating four stocks.

```
simStock <- function(N) {
  dailyMovements = sample(c(-1, 1), size=N, replace = TRUE)
  totalMovements = cumsum(dailyMovements)
  return(totalMovements)
}
N = 10000
s1 = simStock(N); s2 = simStock(N)
s3 = simStock(N); s4 = simStock(N)
plot(1:N, s1, type="l",
     ylim=c(-119, 212), # Adjust according to simulation results
     main = "Stock simulation")
lines(1:N, s2, type="l", col="red")
lines(1:N, s3, type="l", col="blue")
lines(1:N, s4, type="l", col="green")
```

R



The simulations have mean 0, so they should vary around the x-axis. Since the variance is  $n$ , it is dependent on the time. The longer the simulations last, the more they will vary which is what we see.

The standard deviation in this example will be the square root of  $n$ , which is about 100, and that is also what we see (and it is expected that they will not be exactly on the standard deviation).

### 3.12

Deriving the general expression for the expectation and variance for a whole bunch of probability distributions.

- **Bernoulli** with parameter  $p$ . Possible values:  $\mathcal{X} = \{0, 1\}$ .

$$f(x) = p^x(1-p)^{x-1}$$

Expectation.

$$\mathbb{E}[X] = \sum_{x \in \mathcal{X}} x \cdot f(x) = (0) + (1)p^1 = p$$

Second moment.

$$\mathbb{E}[X^2] = \sum_{x^2 \in \mathcal{X}} x^2 \cdot f(x) = (0) + (1)^2 p^1 = p$$

Variance.

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = p - p^2 = p(1-p)$$

- **Poisson** with parameter  $\lambda$ . Possible values:  $\mathcal{X} = \mathbb{N} \cup \{0\}$ .

$$f(x) = e^{-\lambda} \frac{\lambda^x}{x!}$$

Expectation. First term disappears. Cancel  $x$  against the factorial. Switch to  $k = x - 1$ .

$$\begin{aligned} \mathbb{E}[X] &= \sum_{x \in \mathcal{X}} x f(x) = \sum_{x=0}^{\infty} x \cdot e^{-\lambda} \frac{\lambda^x}{x!} \\ &= 0 + \lambda e^{-\lambda} \sum_{x=1}^{\infty} \frac{\lambda^{x-1}}{(x-1)!} \\ &= \lambda e^{-\lambda} \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} \\ &= \lambda e^{-\lambda} e^{\lambda} \\ &= \lambda \end{aligned}$$

Second moment. We will split the sum into two Poisson sums.

$$\begin{aligned} \mathbb{E}[X^2] &= \sum_{x^2 \in \mathcal{X}} x^2 \cdot f(x) = \sum_{x=0}^{\infty} x^2 \cdot e^{-\lambda} \frac{\lambda^x}{x!} \\ &= 0 + \lambda e^{-\lambda} \sum_{x=1}^{\infty} x \cdot \frac{\lambda^{x-1}}{(x-1)!} \\ &= \lambda e^{-\lambda} \sum_{x=1}^{\infty} (x-1+1) \cdot \frac{\lambda^{x-1}}{(x-1)!} \\ &= \lambda e^{-\lambda} \sum_{x=1}^{\infty} \left[ (x-1) \frac{\lambda^{x-1}}{(x-1)!} + \frac{\lambda^{x-1}}{(x-1)!} \right] \end{aligned}$$

$$\begin{aligned}
&= \lambda e^{-\lambda} \left( \sum_{x=2}^{\infty} (x-1) \frac{\lambda^{x-1}}{(x-1)!} + \sum_{x=1}^{\infty} \frac{\lambda^{x-1}}{(x-1)!} \right) \\
&= \lambda e^{-\lambda} \left( \lambda \sum_{x=2}^{\infty} \frac{\lambda^{x-2}}{(x-2)!} + \sum_{x=1}^{\infty} \frac{\lambda^{x-1}}{(x-1)!} \right) \\
&= \lambda e^{-\lambda} \left( \lambda \sum_{j=0}^{\infty} \frac{\lambda^j}{j!} + \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} \right) \\
&= \lambda e^{-\lambda} (\lambda e^{\lambda} + e^{\lambda}) \\
&= \lambda^2 + \lambda
\end{aligned}$$

The remaining calculation for the variance is easy.

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \lambda^2 + \lambda - \lambda^2 = \lambda$$

• **Uniform** on  $(a, b)$ . Possible values:  $\mathcal{X} = (a, b)$ .

$$f(x) = \frac{1}{b-a}$$

Expectation.

$$\begin{aligned}
\mathbb{E}[X] &= \int_{\mathcal{X}} x \cdot f(x) = \frac{1}{b-a} \int_a^b x dx \\
&= \frac{1}{b-a} \left[ \frac{x^2}{2} \right]_a^b = \frac{1}{b-a} \left( \frac{b^2 - a^2}{2} \right) \\
&= \frac{1}{b-a} \left( \frac{(b+a)(b-a)}{2} \right) \\
&= \frac{b+a}{2}
\end{aligned}$$

Second moment.

$$\begin{aligned}
\mathbb{E}[X^2] &= \int_{\mathcal{X}} x^2 \cdot f(x) = \frac{1}{b-a} \int_a^b x^2 dx \\
&= \frac{1}{b-a} \left[ \frac{x^3}{3} \right]_a^b = \frac{1}{b-a} \left( \frac{b^3 - a^3}{3} \right) \\
&= \frac{1}{b-a} \left( \frac{(b-a)(b^2 + ab + a^2)}{3} \right) \\
&= \frac{b^2 + ab + a^2}{3}
\end{aligned}$$

Variance.

$$\begin{aligned}
\text{Var}(X) &= \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \frac{b^2 + ab + a^2}{3} - \frac{(b+a)^2}{4} \\
&= \frac{4b^2 + 4ab + 4a^2 - 3b^2 - 6ab - 3a^2}{12} = \frac{b^2 - 2ab + a^2}{12} = \frac{(b-a)^2}{12}
\end{aligned}$$

• **Exponential** with parameter  $\beta$ . Possible values:  $\mathcal{X} = \{x \in \mathbb{R} : x \geq 0\}$ .

$$f(x) = \frac{1}{\beta} \exp(-x/\beta)$$

Expectation.

$$\mathbb{E}[X] = \int_{\mathcal{X}} x \cdot f(x) = \int_0^{\infty} x \cdot \frac{1}{\beta} e^{-x/\beta} dx$$

We will integrate this with the substitution  $u = x/\beta$ , so  $du/dx = 1/\beta$  which gives us  $dx = \beta du$ . Note that the antiderivative of  $ue^{-u}$  is  $-ue^{-u} - e^{-u}$ .

$$\begin{aligned} \mathbb{E}[X] &= \int_0^{\infty} x \cdot \frac{1}{\beta} e^{-x/\beta} dx = \beta \int_0^{\infty} ue^{-u} du \\ &= \beta \left[ -ue^{-u} - e^{-u} \right]_0^{\infty} \\ &= \beta \left( \lim_{u \rightarrow \infty} (-ue^{-u} - e^{-u}) - (-1) \right) \\ &= \beta \end{aligned}$$

Skipping the evaluation of the limit. The first term can be shown to converge to 0 by using L'Hôpital's rule. The second term in the limit obviously becomes 0.

Second moment. Will derive the expression for the expectation in the calculations.

$$\mathbb{E}[X^2] = \int_{\mathcal{X}} x^2 \cdot f(x) = \int_0^{\infty} x^2 \cdot \frac{1}{\beta} e^{-x/\beta} dx$$

Using integration by parts:  $u = x^2$  so  $u' = 2x$ . Setting  $v' = (1/\beta)e^{-x/\beta}$  which yields  $v = -e^{-x/\beta}$ . From the integration-by-parts formula:

$$\int uv' = uv - \int u'v$$

$$\begin{aligned} \mathbb{E}[X^2] &= \int_0^{\infty} x^2 \cdot \frac{1}{\beta} e^{-x/\beta} dx = \left[ -x^2 e^{-x/\beta} \right]_0^{\infty} - \int_0^{\infty} -2xe^{-x/\beta} dx \\ &= 0 + \int_0^{\infty} 2xe^{-x/\beta} dx \\ &= 2\beta \int_0^{\infty} x \cdot \frac{1}{\beta} e^{-x/\beta} dx \\ &= 2\beta \mathbb{E}[X] \\ &= 2\beta^2 \end{aligned}$$

The limit can be shown to converge to 0 by applying L'Hôpital's rule twice. We use a clever trick and substitute in the expectation and get the result. Finally, we calculate the variance.

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = 2\beta^2 - \beta^2 = \beta^2$$

- **Gamma** with parameters  $\alpha, \beta$ . Possible values:  $\mathcal{X} = \{x \in \mathbb{R} : x > 0\}$ .

$$f(x) = \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} \exp(-x/\beta)$$

We will use the definition of the Gamma function, and the 'Gamma difference' chaining property:

$$\Gamma(\alpha + 1) = \int_0^\infty t^\alpha e^{-t} dt, \quad \Gamma(\alpha + 1) = \alpha \Gamma(\alpha)$$

In step 3, we make the substitution  $t = x/\beta$  which makes  $x = \beta t$ .  $dt/dx = 1/\beta$ , so  $dx = \beta dt$ .

$$\begin{aligned} \mathbb{E}[X] &= \int_{\mathcal{X}} x \cdot f(x) = \int_0^\infty x \cdot \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} \exp(-x/\beta) dx \\ &= \int_0^\infty \frac{1}{\Gamma(\alpha)\beta^\alpha} x^\alpha \exp(-x/\beta) dx \\ &= \int_0^\infty \frac{1}{\Gamma(\alpha)\beta^\alpha} (\beta t)^\alpha \exp(-t) \beta dt \\ &= \int_0^\infty \frac{1}{\Gamma(\alpha)\beta^\alpha} \beta^\alpha t^\alpha \exp(-t) \beta dt \\ &= \int_0^\infty \frac{1}{\Gamma(\alpha)\beta^\alpha} \beta^{\alpha+1} t^\alpha \exp(-t) dt \\ &= \frac{\beta^{\alpha+1}}{\Gamma(\alpha)\beta^\alpha} \int_0^\infty t^\alpha \exp(-t) dt \\ &= \frac{\beta}{\Gamma(\alpha)} \int_0^\infty t^\alpha \exp(-t) dt \\ &= \frac{\beta \Gamma(\alpha + 1)}{\Gamma(\alpha)} \\ &= \frac{\beta \alpha \Gamma(\alpha)}{\Gamma(\alpha)} \\ &= \alpha \beta \end{aligned}$$

For the Gamma distribution, we will not calculate the second moment. Instead we will go straight to the variance calculation and calculate it directly. We will use the same substitution and use the definition of the Gamma function again.

$$\begin{aligned} \text{Var}(X) &= \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \int_0^\infty x^2 \cdot \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} \exp(-x/\beta) dx - \alpha^2 \beta^2 \\ &= \frac{1}{\Gamma(\alpha)\beta^\alpha} \int_0^\infty x^{\alpha+1} \exp(-x/\beta) dx - \alpha^2 \beta^2 \\ &= \frac{1}{\Gamma(\alpha)\beta^\alpha} \int_0^\infty (\beta t)^{\alpha+1} \exp(-t) \beta dt - \alpha^2 \beta^2 \\ &= \frac{\beta^{\alpha+2}}{\Gamma(\alpha)\beta^\alpha} \int_0^\infty t^{\alpha+1} \exp(-t) dt - \alpha^2 \beta^2 \end{aligned}$$

Using the definition of the Gamma function.

$$\begin{aligned}
&= \frac{\beta^2 \Gamma(\alpha + 2)}{\Gamma(\alpha)} - \alpha^2 \beta^2 \\
&= \frac{\beta^2 \Gamma(\alpha + 2) - \alpha^2 \beta^2 \Gamma(\alpha)}{\Gamma(\alpha)} \\
&= \frac{\beta^2 (\Gamma(\alpha + 2) - \alpha^2 \Gamma(\alpha))}{\Gamma(\alpha)} \\
&= \frac{\beta^2 ((\alpha + 1) \Gamma(\alpha + 1) - \alpha^2 \Gamma(\alpha))}{\Gamma(\alpha)} \\
&= \frac{\beta^2 ((\alpha + 1) \alpha \Gamma(\alpha) - \alpha^2 \Gamma(\alpha))}{\Gamma(\alpha)} \\
&= \frac{\beta^2 \Gamma(\alpha) ((\alpha + 1) \alpha - \alpha^2)}{\Gamma(\alpha)} \\
&= \beta^2 ((\alpha + 1) \alpha - \alpha^2) \\
&= \beta^2 (\alpha^2 \alpha - \alpha^2) \\
&= \alpha \beta^2
\end{aligned}$$

And the calculation for the Gamma distribution has been completed.

• **Beta** with parameter  $\alpha, \beta$ . Possible values:  $\mathcal{X} = (0, 1)$ .

$$f(x) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha) \Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1}$$

In the following, we will use the following definition:

$$\frac{\Gamma(\alpha) \Gamma(\beta)}{\Gamma(\alpha + \beta)} = \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} dt$$

Expectation.

$$\begin{aligned}
\mathbb{E}[X] &= \int_{\mathcal{X}} x \cdot f(x) = \int_0^1 x \cdot \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha) \Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1} dx \\
&= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha) \Gamma(\beta)} \int_0^1 x^{\alpha} (1-x)^{\beta-1} dx \\
&= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha) \Gamma(\beta)} \cdot \frac{\Gamma(\alpha + 1) \Gamma(\beta)}{\Gamma(\alpha + \beta + 1)} \\
&= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha) \Gamma(\beta)} \cdot \frac{\alpha \Gamma(\alpha) \Gamma(\beta)}{(\alpha + \beta) \Gamma(\alpha + \beta)} \\
&= \frac{\alpha}{\alpha + \beta} \cdot \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha) \Gamma(\beta)} \cdot \frac{\Gamma(\alpha) \Gamma(\beta)}{\Gamma(\alpha + \beta)} \\
&= \frac{\alpha}{\alpha + \beta}
\end{aligned}$$

Variance. Just like for the Gamma distribution, we calculate the second moment directly in the variance calculation. Will use a lot of the same tricks as when calculating the expectation.

$$\begin{aligned}
\text{Var}(X) &= \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \int_0^1 x^2 \cdot \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1} dx - \left(\frac{\alpha}{\alpha+\beta}\right)^2 \\
&= \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 x^{\alpha+1} (1-x)^{\beta-1} dx - \left(\frac{\alpha}{\alpha+\beta}\right)^2 \\
&= \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 x^{\alpha+1} (1-x)^{\beta-1} dx - \left(\frac{\alpha}{\alpha+\beta}\right)^2 \\
&= \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \cdot \frac{\Gamma(\alpha+2)\Gamma(\beta)}{\Gamma(\alpha+\beta+2)} - \left(\frac{\alpha}{\alpha+\beta}\right)^2 \\
&= \frac{\alpha(\alpha+1)}{(\alpha+\beta)(\alpha+\beta+1)} \cdot \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \cdot \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)} - \left(\frac{\alpha}{\alpha+\beta}\right)^2 \quad (3.12.1) \\
&= \frac{\alpha(\alpha+1)}{(\alpha+\beta)(\alpha+\beta+1)} - \left(\frac{\alpha}{\alpha+\beta}\right)^2 \\
&= \frac{\alpha^2 + \alpha}{(\alpha+\beta)(\alpha+\beta+1)} - \frac{\alpha^2}{(\alpha+\beta)^2} \\
&= \frac{(\alpha^2 + \alpha)(\alpha+\beta) - \alpha^2(\alpha+\beta+1)}{(\alpha+\beta)^2(\alpha+\beta+1)} \\
&= \frac{\alpha^3 + \alpha^2\beta + \alpha^2 + \alpha\beta - \alpha^3 - \alpha^2\beta - \alpha^2}{(\alpha+\beta)^2(\alpha+\beta+1)} \\
&= \frac{\cancel{\alpha^3} + \cancel{\alpha^2\beta} + \cancel{\alpha^2} + \alpha\beta - \cancel{\alpha^3} - \cancel{\alpha^2\beta} - \cancel{\alpha^2}}{(\alpha+\beta)^2(\alpha+\beta+1)} \\
&= \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}
\end{aligned}$$

In step (3.12.1) we used the difference/chaining property of the Gamma function.

$$\Gamma(\alpha+2) = (\alpha+1)\Gamma(\alpha+1) = (\alpha+1)\alpha\Gamma(\alpha)$$

and,

$$\Gamma(\alpha+\beta+2) = (\alpha+\beta+1)\Gamma(\alpha+\beta+1) = (\alpha+\beta+1)(\alpha+\beta)\Gamma(\alpha+\beta)$$

This concludes the last calculation. I found this exercise to be a little tedious, to be honest...

### 3.13

A variable  $X$  is generated by assuming a value in  $U(0,1)$  if a coin toss is 0 (H), and  $X$  is in  $U(3,4)$  if the coin toss is 1 (T). The coin is fair, so each of these has a probability of  $1/2$ .

(a) Calculating the mean of  $X$ . This will be a conditional expectation, and we define  $Y$  to be the coin toss. The marginal distributions might depend on the outcome, but when we calculate them:

$$f_{X|Y}(x|Y=0) = \frac{1}{1-0} = 1, \quad f_{X|Y}(x|Y=1) = \frac{1}{4-3} = 1,$$

we find that it will be  $f_{X|Y}(x|y) = 1$  either way.

Now we can calculate the expected value for specific outcomes.

$$\mathbb{E}[X|Y=0] = \int_{\mathcal{X}} x \cdot f_{X|Y}(x|y) = \int_0^1 x dx = \left[ \frac{x^2}{2} \right]_0^1 = \frac{1-0}{2} = \frac{1}{2}$$

$$\mathbb{E}[X|Y=1] = \int_{\mathcal{X}} x \cdot f_{X|Y}(x|y) = \int_3^4 x dx = \left[ \frac{x^2}{2} \right]_3^4 = \frac{16-9}{2} = \frac{7}{2}$$

Using that  $p_Y(y)$  is known, we can calculate the expected value of  $X$ :

$$\begin{aligned} \mathbb{E}[X] &= \mathbb{E}[\mathbb{E}[X|Y]] = \sum_{y \in \{0,1\}} \mathbb{E}[X|Y=y] \mathbb{P}(Y=y) \\ &= \left( \frac{1}{2} \right) \left( \frac{1}{2} \right) + \left( \frac{7}{2} \right) \left( \frac{1}{2} \right) \\ &= \frac{1}{4} + \frac{7}{4} = \frac{8}{4} = 2 \end{aligned}$$

(b) Calculating the second moment.

$$\mathbb{E}[X^2|Y=0] = \int_{\mathcal{X}} x^2 \cdot f_{X|Y}(x|y) = \int_0^1 x^2 dx = \left[ \frac{x^3}{3} \right]_0^1 = \frac{1-0}{3} = \frac{1}{3}$$

$$\mathbb{E}[X^2|Y=1] = \int_{\mathcal{X}} x^2 \cdot f_{X|Y}(x|y) = \int_3^4 x^2 dx = \left[ \frac{x^3}{3} \right]_3^4 = \frac{4^3-3^3}{3} = \frac{37}{3}$$

$$\begin{aligned} \mathbb{E}[X^2] &= \mathbb{E}[\mathbb{E}[X^2|Y]] = \sum_{y \in \{0,1\}} \mathbb{E}[X^2|Y=y] \mathbb{P}(Y=y) \\ &= \left( \frac{1}{3} \right) \left( \frac{1}{2} \right) + \left( \frac{37}{3} \right) \left( \frac{1}{2} \right) \\ &= \frac{1}{6} + \frac{37}{6} = \frac{38}{6} \\ &= \frac{19}{3} \end{aligned}$$

Calculating the variance:

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \frac{19}{3} - 4 = \frac{7}{3} \approx 2.333$$

And finally the standard deviation:

$$\text{SD}(X) = \sqrt{\text{Var}(X)} = \sqrt{\frac{7}{3}} \approx 1.5275$$

To verify these results, we did a numeric simulation.



```

N = 10000; Y = sample(c(0, 1), size=N, replace = TRUE); X = rep(0, N)
for (i in 1:N) {
  if (Y[i] == 1) X[i] = runif(1)
  else X[i] = runif(1, min=3, max=4)
}
> mean(X)
[1] 2.01424
> var(X)
[1] 2.332699
> sd(X)
[1] 1.527318

```

R

### 3.14

**Claim:** Let  $X_1, \dots, X_m$  and  $Y_1, \dots, Y_n$  be random variables and  $a_1, \dots, a_m$  and  $b_1, \dots, b_n$  be constants. Then:

$$\text{Cov} \left( \sum_{i=1}^m a_i X_i, \sum_{j=1}^n b_j Y_j \right) = \sum_{i=1}^m \sum_{j=1}^n a_i b_j \text{Cov}(X_i, Y_j).$$

PROOF. This can be verified with a direct proof. We will do a simplified example with  $m = n = 2$ , which is not a formal proof, but can easily be extended to one. We will repeatedly use:

$$\text{Cov}(A, B) = \mathbb{E}[AB] - \mathbb{E}[A]\mathbb{E}[B].$$

Starting with the covariance, we apply the identity above. We have, for  $m = n = 2$ :

$$\begin{aligned} \text{Cov}(a_1 X_1 + a_2 X_2, b_1 Y_1 + b_2 Y_2) &= \mathbb{E}[(a_1 X_1 + a_2 X_2)(b_1 Y_1 + b_2 Y_2)] - \mathbb{E}[a_1 X_1 + a_2 X_2] \mathbb{E}[b_1 Y_1 + b_2 Y_2] \\ &= S_1 - S_2 \end{aligned}$$

Multiplying and rewriting the first term:

$$\begin{aligned} S_1 &= \mathbb{E}[(a_1 X_1 + a_2 X_2)(b_1 Y_1 + b_2 Y_2)] = \mathbb{E}[a_1 b_1 X_1 Y_1 + a_1 b_2 X_1 Y_2 + a_2 b_1 X_2 Y_1 + a_2 b_2 X_2 Y_2] \\ &= a_1 b_1 \mathbb{E}[X_1 Y_1] + a_1 b_2 \mathbb{E}[X_1 Y_2] + a_2 b_1 \mathbb{E}[X_2 Y_1] + a_2 b_2 \mathbb{E}[X_2 Y_2] \end{aligned}$$

The second term:

$$\begin{aligned} S_2 &= \mathbb{E}[a_1 X_1 + a_2 X_2] \mathbb{E}[b_1 Y_1 + b_2 Y_2] = (a_1 \mathbb{E}[X_1] + a_2 \mathbb{E}[X_2]) (b_1 \mathbb{E}[Y_1] + b_2 \mathbb{E}[Y_2]) \\ &= a_1 b_1 \mathbb{E}[X_1] \mathbb{E}[Y_1] + a_1 b_2 \mathbb{E}[X_1] \mathbb{E}[Y_2] + a_2 b_1 \mathbb{E}[X_2] \mathbb{E}[Y_1] + a_2 b_2 \mathbb{E}[X_2] \mathbb{E}[Y_2] \end{aligned}$$

Subtracting:

$$\begin{aligned} S_1 - S_2 &= a_1 b_1 \mathbb{E}[X_1 Y_1] + a_1 b_2 \mathbb{E}[X_1 Y_2] + a_2 b_1 \mathbb{E}[X_2 Y_1] + a_2 b_2 \mathbb{E}[X_2 Y_2] \\ &\quad - a_1 b_1 \mathbb{E}[X_1] \mathbb{E}[Y_1] - a_1 b_2 \mathbb{E}[X_1] \mathbb{E}[Y_2] - a_2 b_1 \mathbb{E}[X_2] \mathbb{E}[Y_1] - a_2 b_2 \mathbb{E}[X_2] \mathbb{E}[Y_2] \\ &= a_1 b_1 (\mathbb{E}[X_1 Y_1] - \mathbb{E}[X_1] \mathbb{E}[Y_1]) + a_1 b_2 (\mathbb{E}[X_1 Y_2] - \mathbb{E}[X_1] \mathbb{E}[Y_2]) \\ &\quad + a_2 b_1 (\mathbb{E}[X_2 Y_1] - \mathbb{E}[X_2] \mathbb{E}[Y_1]) + a_2 b_2 (\mathbb{E}[X_2 Y_2] - \mathbb{E}[X_2] \mathbb{E}[Y_2]) \\ &= a_1 b_1 \text{Cov}(X_1, Y_1) + a_1 b_2 \text{Cov}(X_1, Y_2) + a_2 b_1 \text{Cov}(X_2, Y_1) + a_2 b_2 \text{Cov}(X_2, Y_2) \\ &= \sum_{i=1}^2 \sum_{j=1}^2 a_i b_j \text{Cov}(X_i, Y_j) \end{aligned} \quad \square$$

### 3.15

Defining the joint distribution:

$$f_{X,Y}(x,y) = \begin{cases} \frac{1}{3}(x+y) & 0 \leq x \leq 1, 0 \leq y \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

We will calculate  $\text{Var}(2X - 3Y + 8)$ . Constants disappear in the variance, and by Theorem 2.30:

$$\text{Var}(2X - 3Y + 8) = 4\text{Var}(X) + 9\text{Var}(Y) - 12\text{Cov}(X, Y)$$

Finding the marginal distributions:

$$f_X(x) = \int_0^2 \frac{1}{3}x + \frac{1}{3}y dy = \left[ \frac{1}{3}xy + \frac{1}{6}y^2 \right]_0^2 = \frac{2x}{3} + \frac{2}{3}$$

$$f_Y(y) = \int_0^1 \frac{1}{3}x + \frac{1}{3}y dx = \left[ \frac{1}{6}x^2 + \frac{1}{3}xy \right]_0^1 = \frac{y}{3} + \frac{1}{6}$$

Calculating the expectation of  $X$ .

$$\mathbb{E}[X] = \int_0^1 x \left( \frac{2x}{3} + \frac{2}{3} \right) dx = \int_0^1 \left( \frac{2x^2}{3} + \frac{2x}{3} \right) dx = \left[ \frac{2x^3}{9} + \frac{2x^2}{6} \right]_0^1 = \frac{5}{9}$$

Second moment:

$$\mathbb{E}[X^2] = \int_0^1 x^2 \left( \frac{2x}{3} + \frac{2}{3} \right) dx = \int_0^1 \left( \frac{2x^3}{3} + \frac{2x^2}{3} \right) dx = \left[ \frac{2x^4}{12} + \frac{2x^3}{9} \right]_0^1 = \frac{7}{18}$$

Variance of  $X$ :

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \frac{7}{18} - \frac{25}{81} = \frac{13}{162}$$

Calculating the expectation of  $Y$ .

$$\mathbb{E}[Y] = \int_0^2 y \left( \frac{y}{3} + \frac{1}{6} \right) dy = \int_0^2 \left( \frac{y^2}{3} + \frac{y}{6} \right) dy = \left[ \frac{y^3}{9} + \frac{y^2}{12} \right]_0^2 = \frac{11}{9}$$

Second moment:

$$\mathbb{E}[Y^2] = \int_0^2 y^2 \left( \frac{y}{3} + \frac{1}{6} \right) dy = \int_0^2 \left( \frac{y^3}{3} + \frac{y^2}{6} \right) dy = \left[ \frac{y^4}{12} + \frac{y^3}{18} \right]_0^2 = \frac{16}{9}$$

Variance of  $Y$ :

$$\text{Var}(Y) = \mathbb{E}[Y^2] - \mathbb{E}[Y]^2 = \frac{16}{9} - \frac{121}{81} = \frac{23}{81}$$

From the definition of the covariance - these calculations are pretty large, so skipping the details:

$$\begin{aligned}
\text{Cov}(X, Y) &= \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])] \\
&= \int_0^2 \int_0^1 \left(x - \frac{5}{9}\right) \left(y - \frac{11}{9}\right) \left(\frac{x}{3} + \frac{y}{3}\right) dx dy \\
&= \frac{1}{486} \int_0^2 -9y^2 + 20y - 11 dy \\
&= -\frac{1}{81}
\end{aligned}$$

Putting it all together:

$$\begin{aligned}
\text{Var}(2X + 3Y - 8) &= 4\text{Var}(X) + 9\text{Var}(Y) - 12\text{Cov}(X, Y) \\
&= 4 \left(\frac{13}{162}\right) + 9 \left(\frac{23}{81}\right) - 12 \left(-\frac{1}{81}\right) \\
&= \frac{52}{162} + \frac{207}{81} + \frac{12}{81} = \frac{26}{81} + \frac{207}{81} + \frac{12}{81} \\
&= \frac{245}{81}
\end{aligned}$$

### 3.16

Let  $r(x)$  and  $s(y)$  be functions of  $x$  and  $y$ . Then:

$$\mathbb{E}[r(X)s(Y)|X] = r(X)\mathbb{E}[s(Y)|X], \quad \mathbb{E}[r(X)|X] = r(X)$$

PROOF. By definition of the conditional expectation in the continuous case.

$$\begin{aligned}
\mathbb{E}[r(X)s(Y)|X = x] &= \int_{\mathcal{Y}} r(x)s(y)f_{Y|X}(y)dy \\
&= r(x) \int_{\mathcal{Y}} s(y)f_{Y|X}(y)dy \\
&= r(X)\mathbb{E}[s(Y)|X]
\end{aligned}$$

For the special case, we can set  $s(Y) = 1$ .

$$\begin{aligned}
\mathbb{E}[r(X)|X = x] &= \int_{\mathcal{Y}} r(x)(1)f_{Y|X}(y)dy \\
&= r(x) \int_{\mathcal{Y}} (1)f_{Y|X}(y)dy \\
&= r(X)\mathbb{E}[(1)|X] \\
&= r(X)
\end{aligned}$$

which proves the statement. It is similar in the discrete case. □