

# Probability Densities in Data Mining

**Andrew W. Moore**

**Professor**

**School of Computer Science**

**Carnegie Mellon University**

[www.cs.cmu.edu/~awm](http://www.cs.cmu.edu/~awm)

[awm@cs.cmu.edu](mailto:awm@cs.cmu.edu)

412-268-7599

Note to other teachers and users of these slides. Andrew would be delighted if you found this source material useful in giving your own lectures. Feel free to use these slides verbatim, or to modify them to fit your own needs. PowerPoint originals are available. If you make use of a significant portion of these slides in your own lecture, please include this message, or the following link to the source repository of Andrew's tutorials: <http://www.cs.cmu.edu/~awm/tutorials> . Comments and corrections gratefully received.

# Probability Densities in Data Mining

- Why we should care
- Notation and Fundamentals of continuous PDFs
- Multivariate continuous PDFs
- Combining continuous and discrete random variables


# Why we should care

- Real Numbers occur in at least 50% of database records
- Can't always quantize them
- So need to understand how to describe where they come from
- A great way of saying what's a reasonable range of values
- A great way of saying how multiple attributes should reasonably co-occur


# Why we should care

- Can immediately get us Bayes Classifiers that are sensible with real-valued data
- You'll need to **intimately** understand PDFs in order to do kernel methods, clustering with Mixture Models, analysis of variance, time series and many other things
- Will introduce us to linear and non-linear regression

# A PDF of American Ages in 2000



# A PDF of American Ages in 2000



Let  $X$  be a continuous random variable.


If  $p(x)$  is a Probability Density Function for  $X$  then...

$$P(a < X \leq b) = \int_{x=a}^b p(x) dx$$

$$P(30 < \text{Age} \leq 50) = \int_{\text{age}=30}^{50} p(\text{age}) d\text{age}$$

$$= 0.36$$

# Properties of PDFs



$$P(a < X \leq b) = \int_{x=a}^b p(x) dx$$

That means...

$$p(x) = \lim_{h \rightarrow 0} \frac{P\left(x - \frac{h}{2} < X \leq x + \frac{h}{2}\right)}{h}$$

$$\frac{\partial}{\partial x} P(X \leq x) = p(x)$$

# Properties of PDFs

$p(\text{age})$

0.015

0.01

$$P(a < X \leq b) = \int_a^b p(x) dx$$

Therefore...

$$\frac{\partial}{\partial x} P(X \leq x) = p(x)$$

Therefore...

$$\int_{-\infty}^{\infty} p(x) dx = 1$$

$$\forall x : p(x) \geq 0$$



# Talking to your stomach

- What's the gut-feel meaning of  $p(x)$ ?

If

$$p(5.31) = 0.06 \text{ and } p(5.92) = 0.03$$

then

when a value  $X$  is sampled from the distribution, you are 2 times as likely to find that  $X$  is "very close to" 5.31 than that  $X$  is "very close to" 5.92.

# Talking to your stomach

- What's the gut-feel meaning of  $p(x)$ ?

If

$$p(\textcolor{red}{a}) = 0.06 \text{ and } p(\textcolor{blue}{b}) = 0.03$$

then

when a value  $X$  is sampled from the distribution, you are 2 times as likely to find that  $X$  is “very close to”  $\textcolor{red}{a}$  than that  $X$  is “very close to”  $\textcolor{blue}{b}$ .

# Talking to your stomach

- What's the gut-feel meaning of  $p(x)$ ?

If

$$p(\textcolor{red}{a}) = 2\textcolor{green}{z} \text{ and } p(\textcolor{blue}{b}) = \textcolor{green}{z}$$

then

when a value  $X$  is sampled from the distribution, you are 2 times as likely to find that  $X$  is “very close to”  $\textcolor{red}{a}$  than that  $X$  is “very close to”  $\textcolor{blue}{b}$ .

# Talking to your stomach

- What's the gut-feel meaning of  $p(x)$ ?

If

$$p(\textcolor{red}{a}) = \alpha \textcolor{green}{z} \text{ and } p(\textcolor{blue}{b}) = \textcolor{green}{z}$$

then

when a value  $X$  is sampled from the distribution, you are  $\alpha$  times as likely to find that  $X$  is "very close to"  $\textcolor{red}{a}$  than that  $X$  is "very close to"  $\textcolor{blue}{b}$ .

# Talking to your stomach

- What's the gut-feel meaning of  $p(x)$ ?

If 
$$\frac{p(a)}{p(b)} = \alpha$$

then

when a value  $X$  is sampled from the distribution, you are  $\alpha$  times as likely to find that  $X$  is “very close to”  $a$  than that  $X$  is “very close to”  $b$  .

# Talking to your stomach

- What's the gut-feel meaning of  $p(x)$ ?


If

$$\frac{p(a)}{p(b)} = \alpha$$

then

$$\lim_{h \rightarrow 0} \frac{P(a - h < X < a + h)}{P(b - h < X < b + h)} = \alpha$$

# Yet another way to view a PDF



A recipe for sampling a random age.

1. Generate a random dot from the rectangle surrounding the PDF curve. Call the dot (age,d)
2. If  $d < p(\text{age})$  stop and return age
3. Else try again: go to Step 1.

# Test your understanding


- True or False:

$$\forall x : p(x) \leq 1$$

$$\forall x : P(X = x) = 0$$



# Expectations




$E[X]$  = the expected value of random variable  $X$

= the average value we'd see if we took a very large number of random samples of  $X$

$$= \int_{-\infty}^{\infty} x p(x) dx$$

# Expectations



$E[X]$  = the expected value of random variable  $X$

= the average value we'd see if we took a very large number of random samples of  $X$

$$= \int_{-\infty}^{\infty} x p(x) dx$$

= the first moment of the shape formed by the axes and the blue curve

= the best value to choose if you must guess an unknown person's age and you'll be fined the square of your error

# Expectation of a function


$\mu = E[f(X)]$  = the expected value of  $f(x)$  where  $x$  is drawn from  $X$ 's distribution.

= the average value we'd see if we took a very large number of random samples of  $f(X)$

$$\mu = \int_{-\infty}^{\infty} f(x) p(x) dx$$

Note that in general:

$$E[f(x)] \neq f(E[X])$$




# Variance

$\sigma^2 = \text{Var}[X]$  = the expected squared difference between  $x$  and  $E[X]$


$$\sigma^2 = \int_{x=-\infty}^{\infty} (x - \mu)^2 p(x) dx$$

= amount you'd expect to lose if you must guess an unknown person's age and you'll be fined the square of your error, and assuming you play optimally



# Standard Deviation

$\sigma^2 = \text{Var}[X]$  = the expected squared difference between  $x$  and  $E[X]$




$$\sigma^2 = \int_{x=-\infty}^{\infty} (x - \mu)^2 p(x) dx$$

= amount you'd expect to lose if you must guess an unknown person's age and you'll be fined the square of your error, and assuming you play optimally

$\sigma$  = Standard Deviation = "typical" deviation of  $X$  from its mean

$$\sigma = \sqrt{\text{Var}[X]}$$

# In 2 dimensions




$p(x,y)$  = probability density of  
random variables  $(X,Y)$  at  
location  $(x,y)$

# In 2 dimensions

Let  $X, Y$  be a pair of continuous random variables, and let  $R$  be some region of  $(X, Y)$  space...

$$P((X, Y) \in R) = \iint_{(x, y) \in R} p(x, y) dy dx$$



# In 2 dimensions

Let  $X, Y$  be a pair of continuous random variables, and let  $R$  be some region of  $(X, Y)$  space...

$$P((X, Y) \in R) = \iint_{(x, y) \in R} p(x, y) dy dx$$

density values:


$2.1\text{e-}005 \leq \text{density} < 3.4\text{e-}005$

$\text{density} \leq 8\text{e-}006$

$3.4\text{e-}005 < \text{density}$

$8\text{e-}006 \leq \text{density} < 2.1\text{e-}005$

weight



$P(20 < \text{mpg} < 30 \text{ and } 2500 < \text{weight} < 3000) =$

area under the 2-d surface within the red rectangle



# In 2 dimensions

Let  $X, Y$  be a pair of continuous random variables, and let  $R$  be some region of  $(X, Y)$  space...

$$P((X, Y) \in R) = \iint_{(x, y) \in R} p(x, y) dy dx$$

density values:


$2.1\text{e-}005 \leq \text{density} < 3.4\text{e-}005$

$\text{density} \leq 8\text{e-}006$

$3.4\text{e-}005 < \text{density}$

$8\text{e-}006 \leq \text{density} < 2.1\text{e-}005$

weight



$$P\left(\left[\frac{\text{mpg}-25}{10}\right]^2 + \left[\frac{\text{weight}-3300}{1500}\right]^2 < 1\right) =$$

area under the 2-d surface within the red oval

# In 2 dimensions

Let  $X, Y$  be a pair of continuous random variables, and let  $R$  be some region of  $(X, Y)$  space...

$$P((X, Y) \in R) = \iint_{(x, y) \in R} p(x, y) dy dx$$

Take the special case of region  $R = \text{"everywhere"}$ .

Remember that with probability 1,  $(X, Y)$  will be drawn from "somewhere".

So..

$$\int_{x=-\infty}^{\infty} \int_{y=-\infty}^{\infty} p(x, y) dy dx = 1$$

# In 2 dimensions

Let  $X, Y$  be a pair of continuous random variables, and let  $R$  be some region of  $(X, Y)$  space...

$$P((X, Y) \in R) = \iint_{(x, y) \in R} p(x, y) dy dx$$

$$p(x, y) = \lim_{h \rightarrow 0} \frac{P\left(x - \frac{h}{2} < X \leq x + \frac{h}{2} \quad \wedge \quad y - \frac{h}{2} < Y \leq y + \frac{h}{2}\right)}{h^2}$$

# In $m$ dimensions


Let  $(X_1, X_2, \dots, X_m)$  be an  $n$ -tuple of continuous random variables, and let  $R$  be some region of  $\mathbf{R}^m$  ...

$$P((X_1, X_2, \dots, X_m) \in R) =$$

$$\iint \dots \int_{(x_1, x_2, \dots, x_m) \in R} p(x_1, x_2, \dots, x_m) dx_m, \dots, dx_2, dx_1$$

# Independence

$$X \perp Y \text{ iff } \forall x, y : p(x, y) = p(x)p(y)$$




If X and Y are independent  
then knowing the value of X  
does not help predict the value  
of Y

mpg, weight NOT  
independent

# Independence

$$X \perp Y \text{ iff } \forall x, y : p(x, y) = p(x)p(y)$$




If X and Y are independent  
then knowing the value of X  
does not help predict the value  
of Y

the contours say that  
acceleration and weight are  
independent

# Multivariate Expectation

$$\mu_{\mathbf{X}} = E[\mathbf{X}] = \int \mathbf{x} p(\mathbf{x}) d\mathbf{x}$$



$E[\text{mpg}, \text{weight}] = (24.5, 2600)$

The centroid of the cloud

# Multivariate Expectation

$$E[f(\mathbf{X})] = \int f(\mathbf{x}) p(\mathbf{x}) d\mathbf{x}$$



# Test your understanding

Question : When (if ever) does  $E[X + Y] = E[X] + E[Y]$  ?

- All the time?
- Only when  $X$  and  $Y$  are independent?
- It can fail even if  $X$  and  $Y$  are independent?

# Bivariate Expectation

$$E[f(x, y)] = \int f(x, y) p(x, y) dy dx$$

$$\text{if } f(x, y) = x \text{ then } E[f(X, Y)] = \int x p(x, y) dy dx$$

$$\text{if } f(x, y) = y \text{ then } E[f(X, Y)] = \int y p(x, y) dy dx$$

$$\text{if } f(x, y) = x + y \text{ then } E[f(X, Y)] = \int (x + y) p(x, y) dy dx$$

$$E[X + Y] = E[X] + E[Y]$$

# Bivariate Covariance

$$\sigma_{xy} = \text{Cov}[X, Y] = E[(X - \mu_x)(Y - \mu_y)]$$

$$\sigma_{xx} = \sigma_x^2 = \text{Cov}[X, X] = \text{Var}[X] = E[(X - \mu_x)^2]$$

$$\sigma_{yy} = \sigma_y^2 = \text{Cov}[Y, Y] = \text{Var}[Y] = E[(Y - \mu_y)^2]$$

# Bivariate Covariance

$$\sigma_{xy} = \text{Cov}[X, Y] = E[(X - \mu_x)(Y - \mu_y)]$$


$$\sigma_{xx} = \sigma_x^2 = \text{Cov}[X, X] = \text{Var}[X] = E[(X - \mu_x)^2]$$

$$\sigma_{yy} = \sigma_y^2 = \text{Cov}[Y, Y] = \text{Var}[Y] = E[(Y - \mu_y)^2]$$


Write  $\mathbf{X} = \begin{pmatrix} X \\ Y \end{pmatrix}$ , then

$$\mathbf{Cov}[\mathbf{X}] = E[(\mathbf{X} - \boldsymbol{\mu}_x)(\mathbf{X} - \boldsymbol{\mu}_x)^T] = \boldsymbol{\Sigma} = \begin{pmatrix} \sigma_x^2 & \sigma_{xy} \\ \sigma_{xy} & \sigma_y^2 \end{pmatrix}$$

# Covariance Intuition



# Covariance Intuition



# Covariance Fun Facts

$$\mathbf{Cov}[\mathbf{X}] = E[(\mathbf{X} - \boldsymbol{\mu}_x)(\mathbf{X} - \boldsymbol{\mu}_x)^T] = \boldsymbol{\Sigma} = \begin{pmatrix} \sigma_x^2 & \sigma_{xy} \\ \sigma_{xy} & \sigma_y^2 \end{pmatrix}$$

- True or False: If  $\sigma_{xy} = 0$  then X and Y are independent
- True or False: If X and Y are independent then  $\sigma_{xy} = 0$
- True or False: If  $\sigma_{xy} = \sigma_x \sigma_y$  then X and Y are deterministically related
- True or False: If X and Y are deterministically related then  $\sigma_{xy} = \sigma_x \sigma_y$

How could you prove or disprove these?

# General Covariance

Let  $\mathbf{X} = (X_1, X_2, \dots, X_k)$  be a vector of  $k$  continuous random variables

$$\mathbf{Cov}[\mathbf{X}] = E[(\mathbf{X} - \boldsymbol{\mu}_x)(\mathbf{X} - \boldsymbol{\mu}_x)^T] = \boldsymbol{\Sigma}$$

$$\Sigma_{ij} = \text{Cov}[X_i, X_j] = \sigma_{x_i x_j}$$

$\Sigma$  is a  $k \times k$  symmetric non-negative definite matrix

If all distributions are linearly independent it is positive definite

If the distributions are linearly dependent it has determinant zero




# Test your understanding



Question : When (if ever) does  $\text{Var}[X + Y] = \text{Var}[X] + \text{Var}[Y]$ ?

- All the time?
- Only when  $X$  and  $Y$  are independent?
- It can fail even if  $X$  and  $Y$  are independent?

# Marginal Distributions






density values:  $2.1\text{e-}005 \leq \text{density} < 3.4\text{e-}005$   
 $\text{density} \leq 8\text{e-}006$   $3.4\text{e-}005 < \text{density}$   
 $8\text{e-}006 \leq \text{density} < 2.1\text{e-}005$




$$p(x) = \int_{y=-\infty}^{\infty} p(x, y) dy$$

# Conditional Distributions



density values:  $2.1\text{e-}005 \leq \text{density} < 3.4\text{e-}005$   
 $\text{density} \leq 8\text{e-}006$   $3.4\text{e-}005 < \text{density}$   
 $8\text{e-}006 \leq \text{density} < 2.1\text{e-}005$




$$p(x \mid y) =$$

p.d.f. of  $X$  when  $Y = y$

# Conditional Distributions


$p(\text{mpg} \mid \text{weight} = 4600)$



$$p(x \mid y) = \frac{p(x, y)}{p(y)}$$

Why?

density values:  $2.1\text{e-}005 \leq \text{density} < 3.4\text{e-}005$   
 $\text{density} \leq 8\text{e-}006$   $3.4\text{e-}005 < \text{density}$   
 $8\text{e-}006 \leq \text{density} < 2.1\text{e-}005$



$p(x \mid y) =$   
 p.d.f. of  $X$  when  $Y = y$

# Independence Revisited

$$X \perp Y \text{ iff } \forall x, y : p(x, y) = p(x)p(y)$$

It's easy to prove that these statements are equivalent...

$$\forall x, y : p(x, y) = p(x)p(y)$$

$$\Leftrightarrow$$

$$\forall x, y : p(x | y) = p(x)$$

$$\Leftrightarrow$$

$$\forall x, y : p(y | x) = p(y)$$


# $\infty$ More useful stuff

$$\int_{x=-\infty}^{\infty} p(x | y) dx = 1$$

(These can all be proved from definitions on previous slides)

$$p(x | y, z) = \frac{p(x, y | z)}{p(y | z)}$$

$$p(x | y) = \frac{p(y | x) p(x)}{p(y)}$$



# Mixing discrete and continuous variables

$$p(x, A = v) = \lim_{h \rightarrow 0} \frac{P\left(x - \frac{h}{2} < X \leq x + \frac{h}{2} \wedge A = v\right)}{h}$$

$$\sum_{v=1}^{n_A} \int_{x=-\infty}^{\infty} p(x, A = v) dx = 1$$

$$p(x | A) = \frac{P(A | x) p(x)}{P(A)}$$




Bayes  
Rule

$$P(A | x) = \frac{p(x | A) P(A)}{p(x)}$$



Bayes  
Rule


# Mixing discrete and continuous variables






# Mixing discrete and continuous variables

$P(\text{EduYears}, \text{Wealthy})$




$P(\text{Wealthy} | \text{EduYears})$





# Mixing discrete and continuous variables

$P(\text{EduYears}, \text{Wealthy})$



$P(\text{Wealthy} | \text{EduYears})$

$P(\text{EduYears} | \text{Wealthy})$



# What you should know

- You should be able to play with discrete, continuous and mixed joint distributions
- You should be happy with the difference between  $p(x)$  and  $P(A)$
- You should be intimate with expectations of continuous and discrete random variables
- You should smile when you meet a covariance matrix
- Independence and its consequences should be second nature

# Discussion

- Are PDFs the only sensible way to handle analysis of real-valued variables?
- Why is covariance an important concept?
- Suppose  $X$  and  $Y$  are independent real-valued random variables distributed between 0 and 1:
  - What is  $p[\min(X,Y)]$ ?
  - What is  $E[\min(X,Y)]$ ?
- Prove that  $E[X]$  is the value  $u$  that minimizes  $E[(X-u)^2]$
- What is the value  $u$  that minimizes  $E[|X-u|]$ ?