

Physics S1 Summary

Don't forget the Ppt PS questions, and MCQ will be the hardest section

- FBD will come

Free Fall

Free Fall :

- is the motion of an object where gravity is the only force acting upon it.
- The value of g (upward direction) = -9.8 m/s^2

Formulas :

$$\Delta x = v_i t + \frac{1}{2} a t^2$$

$$v_f = v_i + at$$

$$v_f^2 = v_i^2 + 2a\Delta x$$

- This lesson will mostly be in the MCQ
-

Friction

Friction : *it's dependent on the normal force acting on the object

- It is a force that opposes that relative motion or tendency of such motion of 2 surfaces in-contact.

3 Types of Friction :

1. Static Friction
2. Kinetic Friction
3. Rolling Friction

Factors that affect Friction :

1. **Surface roughness** :-
 - rougher surfaces tend to have higher friction due to more interlocking features.
2. **Normal Force** :-
 - the force pressing 2 surfaces together; greater normal force typically results in higher friction.
 - Equal in magnitude (to the weight).
3. **Material Properties** :-
 - different materials exhibit different coefficients of friction, which quantifies the frictional force relative to the normal.

Physics S1 Summary

Don't forget the Ppt PS questions, and MCQ will be the hardest section

- FBD will come

Static Force/Friction :

- Opposes the initiation of motion
- Magnitude : Varies up to a maximum value, where if exceeded kinetic force will take turn and the object starts moving.

Kinetic Force/Friction :

- Opposes the motion of an object already in motion.
- Magnitude : generally constant for a given situation.

Similarities :-

- Nature of Force : Both types of forces oppose relative motion.
- Dependent on Surface : Both types of friction depend on the surface.

Formula :

F_n is the normal force.

$$\bullet \quad F_f = \mu_s \cdot F_n$$

F_f is the maximum static frictional force,

$$F_f = \mu_k \cdot F_n$$

F_f is the kinetic frictional force,

Unbalanced Friction Forces

- No problem solving questions for this lesson.
- In short, it's when the forces acting on the object are unbalanced(not equal in magnitude to one another.)

Unbalanced friction forces :

- They occur when forces are not equal in magnitude, leading to an acceleration or deceleration of an object.

Formulas :

$$F_{net} = \mu_k \cdot F_N$$
$$F_{net} = ma$$

$$F_{net} = F_A - F_f$$
$$F_f = \mu_k mg$$

Physics S1 Summary

Don't forget the Ppt PS questions, and MCQ will be the hardest section


- FBD will come

Friction on Inclined Planes

- No problem solving for this lesson, only theoretical questions will be coming

Inclined Plane :

- It is a flat surface that is tilted at an angle to the horizontal.



Component Formulas : * not given

$$F_{g_x} = F_g \cdot \sin \theta \rightarrow \text{Horizontal}$$

$$F_{g_y} = F_g \cdot \cos \theta \rightarrow \text{Vertical}$$

$$F_N = mg$$

Tips/To take note of :

- F_N is perpendicular(straight) to the object -> straight to the (y-component)
- Angles should/must be the sum of $90^\circ \Rightarrow$ because of course it's a right angle

Physics S1 Summary

Don't forget the Ppt PS questions, and MCQ will be the hardest section

- FBD will come

Projectile Motion

Projectile Motion : *The object is called a projectile.


- It is a form of motion in which an object is thrown near the earth's surface, and it moves along a curved path under the action of gravity only.

Projectile Motion determining factors :

- Mass, angle of projection, maximum height, initial velocity, and range.
- Horizontal and vertical components $\Rightarrow V_x$ & V_y

The projectile motion drawing/diagram :

FORMULAS : TO COMPLETE



To take note of :

- Horizontal Component does not change (during projectile).
- Vertical component increases until it reaches 0m/s(max velocity), and then it starts decreasing(it's actually increasing in negative values)
- The path of the projectile is called a **trajectory**.
- Gravity is the only force that affects it.
- The projectile makes a parabola when in motion, and follows a parabolic path \Rightarrow **Parabolic projectile**

Analysis :

- 45° , always gives the maximum range(no matter what).
- Range does not depend on the angle of projection, only the max height depends on the angle of projection.
- Angles can have the same range but not max height. \rightarrow it's because each angle will have a different angle which results in different heights.

* When two angles are complementary (= 90)

$$60^\circ + 30^\circ = 90$$

^^^^^^^^^^ \Rightarrow Both will have the same range.

Physics S1 Summary

Don't forget the Ppt PS questions, and MCQ will be the hardest section

- FBD will come

Periodic Motion

Periodic Motion :

- a motion that repeats itself after a certain equal interval of time.
- Ex: a water wave, rocking chair, a tuning fork, a bouncing ball, a swing in motion, a rolling ball, and etc.

Simple Harmonic Motion (SHM) :

- Equilibrium position $\rightarrow x = 0$, it's when the block/object is at rest.
- if the force is always directed towards the equilibrium position, the motion is called **SHM**.
- Ex: Clock, Musical instruments, Spring, and etc.
- If the block was to be displaced to the left (from the right) \rightarrow negative position.

Hooke's Law :-

- The force "F" exerted by a spring is directly proportional to the displacement "x" from its equilibrium position provided the elastic limit is not exceeded.
- If it exceeds the limit, then this will not obey Hooke's Law (from this moment).

Formula :

$$F = -kx$$

Where,

F : Spring force

x : Displacement

k : Spring constant

Pendulum

Pendulum :

- is a body suspended from a fixed point so that it can swing back and forth under the influence of gravity.

Time Period of a Pendulum :

- the time a pendulum takes to complete 1 cycle.
- In a pendulum, the value of $g = 9.8 \text{ m/s}^2$

Formula :

$$T = 2\pi \sqrt{\frac{L}{g}}$$

T = period of simple pendulum (s)
L = length of pendulum (m)
g = gravitational acceleration (m/s²)

