



LYFGENIA (PREAUTHORIZATION REQUIRED)

X.224

X.224 LYFGENIA (PREAUTHORIZATION REQUIRED)

POLICY

- I. Target Agent(s) may be considered **medically necessary** when ALL of the following are met:
 - A. The patient is at least 12 years old **AND**
 - B. Provider has considered use of prophylaxis therapy for seizures prior to initiating myeloablative conditioning **AND**
 - C. Patient will be monitored for hematologic malignancies periodically after treatment **AND**
 - D. Must not be administered concurrently with live vaccines while immunosuppressed **AND**
 - E. Patient does not have a history of hypersensitivity to dimethyl sulfoxide (DMSO) or dextran 40 **AND**
 - F. Patient is HIV negative as confirmed by a negative HIV test prior to mobilization (Note: Patients who have received Lyfgenia are likely to test positive by polymerase chain reaction (PCR) assays for HIV due to integrated BB305 LVV proviral DNA, resulting in a possible false-positive PCR assay test result for HIV. Therefore, patients who have received Lyfgenia should not be screened for HIV infection using a PCR-based assay.) **AND**



1. Hydroxyurea for at least 2 months prior to mobilization and until all cycles of apheresis are completed (Note: If hydroxyurea is administered between mobilization and conditioning, discontinue 2 days prior to initiation of conditioning) **AND**
 2. Myelosuppressive iron chelators (e.g., deferiprone, etc.) for 7-days prior to mobilization, conditioning, and 6 months post-treatment **AND**
 3. Disease-modifying agents (e.g., L-glutamine, vexelotor, crizanlizumab) for at least 2 months prior to mobilization **AND**
 4. Prophylactic HIV anti-retroviral therapy (Note: Patients receiving prophylactic ART should stop therapy for at least one month prior to mobilization and until all cycles of apheresis are completed **AND**
 5. Mobilization of stem cells using granulocyte-colony stimulating factor (G-CSF) **AND**
 6. Erythropoietin for at least 2 months prior to mobilization **AND**
- H. Patient has not received other gene therapy [e.g., Casgevy™ (exagamglogene autotemcel)] **AND**
- I. Patient is a candidate for autologous hematopoietic stem cell transplant (HSCT) **AND**
- J. Patient has a confirmed diagnosis of sickle-cell disease (includes genotypes $\beta S/\beta S$ or $\beta S/\beta 0$ or $\beta S/\beta +$) as determined by one of the following



variant by hemoglobin assay **OR**

2. Identification of biallelic HBB pathogenic variants where at least one allele is the P.Glu6Val pathogenic variant on molecular genetic testing **AND**

- K. Patient does NOT have disease with more than two α -globin gene deletions **AND**
- L. Patient experienced at least two vaso-occlusive events/crises (VOE/VOC)* in the previous 12 months while adhering to the above therapy (Note: Patients experiencing four events/crises in the previous 24 months will also have met this requirement)

*VOE/VOC is defined as an event requiring a visit to a medical facility for evaluation which results in a diagnosis of such being documented due to one (or more) of the following: acute pain, acute chest syndrome, acute splenic sequestration, acute hepatic sequestration, priapism lasting > 2 hours AND necessitating subsequent interventions such as opioid pain management, non-steroidal anti-inflammatory drugs, RBC transfusion, etc

Length of Approval: 1 per lifetime

Dates

Original Effective

03-01-2024

Last Review

11-06-2024

Next Review

11-06-2025

CLINICAL RATIONALE

LYFGENIA (lovotibeglogene autotemcel) is a β A-T87Q-globin gene therapy consisting of autologous CD34+ cells from patients with sickle cell disease



a modified form of the β -globin gene (β A-T87Q-globin gene) into the patient's own HSCs.

LYFGENIA is prepared using the patient's own HSCs, which are collected via apheresis procedure(s). The autologous cells are enriched for CD34+ cells, then transduced ex vivo with BB305 LVV. The promoter, a regulatory element that controls the expression of the transgene selected for BB305 LVV, is a cellular (non-viral) promoter that controls gene expression specific to the erythroid lineage cells (red blood cells and their precursors). BB305 LVV encodes β A-T87Q-globin.

The efficacy of LYFGENIA was studied in a single-arm, 24-month, open-label, multicenter Phase 1/2 study (Study 1-C) and continued on a long-term follow-up study. In Study 1-C, 43 subjects underwent apheresis after mobilization with plerixafor of which 36 patients received myeloablative busulfan conditioning. Seven patients did not proceed to conditioning; 2 patients discontinued due to apheresis-related issues and 5 discontinued at patient and/or physician discretion.

The transplant population for VOE efficacy outcomes included patients with a history of at least 4 VOEs in the 24 months prior to informed consent. The efficacy outcomes were complete resolution of VOEs (VOE-CR) and severe VOEs (sVOE-CR) between 6 months and 18 months after infusion of LYFGENIA. VOEs were defined as any of the following events requiring evaluation at a medical facility:

- an episode of acute pain with no medically determined cause other than vaso-occlusion, lasting more than 2 hours
 - acute chest syndrome (ACS)
 - acute hepatic sequestration
 - acute splenic sequestration
- Severe VOE (sVOE) were defined as either of the following events:
- VOE requiring a hospitalization or multiple visits to an emergency department/urgent care over 72 hours and receiving intravenous medications at each visit
 - priapism requiring any level of medical attention

The median (min, max) duration of follow-up for the patients in Study 1-C (N = 36) is 38 (12, 61) months post LYFGENIA infusion. After the primary evaluation period to last follow-up, 4 of 32 patients who achieved VOE-CR experienced VOEs while maintaining GR. After the primary evaluation period up to 24 months, 17 of 35 (49%) patients were prescribed opioids for sickle cell and non-sickle cell-related pain.



Quick Code Search

Use this feature to find out if a procedure and diagnosis code pair will be approved, denied or held for review. Simply put in the procedure code, then the diagnosis code, then click "Add Code Pair". If the codes are listed in this policy, we will help you by showing a dropdown to help you.

Procedure

Enter at least the first 3 characters of the code

Diagnosis

Enter at least the first 3 characters of the code

CODES

+ HCPCS

REFERENCES



HGB-206 study. Am J Hematol. 2023 Jan;98(1):11-22. doi: 10.1002/ajh.26741. Epub 2022 Oct 10. PMID: 36161320; PMCID: PMC10092845.

2023

Lyfgenia [package insert]. Somerville, MA; Bluebird Bio, Inc., December 2023. Accessed December 2023.

2022

Bender MA, Carlberg K. Sick Cell Disease. 2003 Sep 15 [Updated 2022 Nov 17]. In: Adam MP, Everman DB, Mirzaa GM, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2022. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK1377/>.

2014

Yawn BP, Buchanan GR, Afenyi-Annan AN, et al. Management of sickle cell disease: summary of the 2014 evidence-based report by expert panel members. JAMA. 2014 Sep 10;312(10):1033-48

2020

Tisdale JF, Pierciey FJ, Bonner M, et al. (2020) Safety and feasibility of hematopoietic progenitor stem cell collection by mobilization with plerixafor followed by apheresis vs bone marrow harvest in patients with sickle cell disease in the multi-center HGB-206 trial. Am J Hematol E239–E242. <https://doi.org/10.1002/ajh.25867>.

2016

Palmer J, McCune JS, Perales M-A, et al. (2016) Personalizing Busulfan-Based Conditioning: Considerations from the American Society for Blood and Marrow Transplantation Practice Guidelines Committee. Biol Blood Marrow Transplant 1915–1925. <https://doi.org/10.1016/j.bbmt.2016.07.013>

2017

Brunson A, Keegan THM, Bang H, et al. (2017) Increased risk of leukemia among sickle cell disease patients in California. Blood 130:1597–1599. doi: 10.1182/blood-2017-05-783233.

2016

Seminog OO, Ogunlaja OI, Yeates D, Goldacre MJ (2016) Risk of individual malignant neoplasms in patients with sickle cell disease: English national record linkage study. J R Soc Med 109:303–309. doi: 10.1177/0141076816651037



Added new 07/01/2024 HCPCS code: J3394

HAVE AN IDEA? WE'RE HERE TO HELP YOU MANAGE YOUR WORK



[Privacy](#) [Legal](#) [Non-Discrimination and Translation](#) [Site Map](#)

© 2025 Blue Cross and Blue Shield of Nebraska.

Blue Shield of Nebraska is an independent licensee of the Blue Cross and Blue Shield Association. The Blue Cross and Blue Shield Association licenses Blue Cross and Blue Shield of Nebraska to offer certain products and services under the Blue Cross® and Blue Shield® brand names within the state of Nebraska.