



PCSK9 (REQUIRES PREAUTHORIZATION)

X.64

X.64 PCSK9 (REQUIRES PREAUTHORIZATION)

POLICY

Target Agents:

Praluent® (alirocumab) is FDA approved for:

- To reduce the risk of myocardial infarction, stroke, and unstable angina requiring hospitalization in adults with established cardiovascular disease
- As an adjunct to diet alone or in combination with other low density lipoprotein cholesterol (LDL-C)- lowering therapies, in adults with primary hyperlipidemia, including heterozygous familial hypercholesterolemia (HeFH), to reduce LDL-C
- As an adjunct to other LDL-C-lowering therapies in adult patients with homozygous familial hypercholesterolemia (HoFH) to reduce LDL-C

Repatha® (evolocumab) is FDA approved for:

- In adults with established cardiovascular disease to reduce the risk of myocardial infarction, stroke, and coronary revascularization
- As an adjunct to diet, alone or in combination with other LDL-C-lowering therapies in adults with primary hyperlipidemia, including HeFH, to reduce LDL-C
- As an adjunct to other LDL-C-lowering therapies in pediatric patients aged 10 years and older with HeFH, to reduce LDL-C
- As an adjunct to diet and other LDL-C-lowering therapies in adults and pediatric patients aged 10 years and older with HoFH, to reduce LDL-C

Initial Evaluation



.. ONE of the following:

A. BOTH of the following:

1. ONE of the following:

a. The patient has a diagnosis of heterozygous familial hypercholesterolemia (HeFH) AND ONE of the following:

i. Genetic confirmation of one mutant allele at the LDLR, Apo-B, PCSK9, or 1/LDLRAP1 gene
OR

ii. History of LDL-C greater than 190 mg/dL (greater than 4.9 mmol/L) (pretreatment) **OR**

iii. The patient has clinical manifestations of HeFH (e.g., cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthoma, or xanthelasma) **OR**

iv. The patient has "definite" or "possible" familial hypercholesterolemia as defined by the Simon Broome criteria
OR

v. The Patient has a Dutch Lipid Clinic Network Criteria score of greater than 5 **OR**

vi. The patient has a treated low-



equal to 100 mg/dL after treatment with antihyperlipidemic agents but prior to PCSK9 inhibitor therapy **OR**

- b. The patient has a diagnosis of homozygous familial hypercholesterolemia (HoFH) AND ONE of the following:

- i. Genetic confirmation of TWO mutant alleles at the LDLR, Apo-B, PCSK9, or LDLRAP1 gene **OR**


- ii. History of untreated LDL-C greater than 500 mg/dL (greater than 13 mmol/L) or treated LDL-C greater than or equal to 300 mg/dL (greater than or equal to 7.76 mmol/L) **OR**

- iii. The patient has clinical manifestations of HoFH (e.g., cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas, or xanthelasma) **OR**

- c. The patient has a diagnosis of clinical atherosclerotic cardiovascular disease (ASCVD) AND has ONE of the following:

- i. Acute coronary syndrome

- ii. History of myocardial infarction



OR

- 4/26



- a) LDL-C
greater than
or equal to
70 mg/dL
while on
maximally
tolerated
statin
therapy
AND
- b) ONE of the
following:
 - a) The
patient
has
extensive
or
active
burden
of
ASCVD
(i.e.,
polyvascular
ASCVD,
which
affects
all 3
vascular
beds
—
coronary,
cerebrovascular,
and
peripheral
arterial;
clinical
peripheral
arterial
disease
in
addition
to
coronary
and/or



clinical
ASCVD
event
with
multivesse
coronary
artery
disease
defined
as
greater
than
or
equal
to
40%
stenosis
in
greater
than
or
equal
to 2
large
vessels;
or
recurrent
myocardia
infarction
within
2
years
of the
initial
event)
in the
presence
of
adverse
or
poorly
controlled
cardiometabolic
risk
factors
OR



risk
elevations
in
cardiometabolic
factors
with
less-
extensive
ASCVD
(i.e.,
diabetes,
LDL-C
greater
than
or
equal
to 100
mg/dL,
less
than
high-
intensity
statin
therapy,
chronic
kidney
disease,
poorly
controlled
hypertension,
high-
sensitivity
C-
reactive
protein
greater
than 3
mg/L,
or
metabolic
syndrome,
usually
occurring
with
other
extremely



high-risk characteristics), usually with other adverse or poorly controlled cardiometabolic risk factors present
OR

c) Patients with ASCVD and LDL-C greater than or equal to 220 mg/dL with greater than or equal to 45% 10-year ASCVD risk despite statin therapy
OR

ii. The patient has 30-39% 10-year



- greater than
or equal to
100 mg/dL
while on
maximally
tolerated
statin
therapy
AND
 - b) Less-
extensive
clinical
ASCVD (i.e.,
no
polyvascular
ASCVD, no
clinical
peripheral
arterial
disease, a
prior ASCVD
event
greater than
or equal to
2 years
prior, and
no coronary
artery
bypass
grafting)
AND
 - c) Adverse or
poorly
controlled
cardiometabolic
risk factor(s)
including
age 65 years
or older,
current
smoking,
chronic
kidney
disease,



37 nmol/L,
high-
sensitivity C-
reactive
protein 1–3
mg/L,
metabolic
syndrome
with a
history of
myocardial
infarction,
ischemic
stroke, or
symptomatic
peripheral
arterial
disease,
usually in
the
presence of
other
adverse or
poorly
controlled
cardiometabolic
risk factors

OR

iii. The patient
has 20-29% 10-year
ASCVD risk AND
BOTH of the
following:

- a) LDL-C
greater than
or equal to
130 mg/dL
while on
maximally
tolerated
statins **AND**
- b) ONE of the
following:



less
extensive
ASCVD
and
well-
controlled
cardiometabolic
risk
factors
(i.e.,
no
diabetes,
nonsmoker,
on
high-
intensity
statin
with
LDL-C
less
than
100
mg/dL,
blood
pressure
less
than
140/90
mm
Hg,
and C-
reactive
protein
less
than 1
mg/dL)
OR

b) The
use is
for
primary
prevention
with
LDL-C
greater
than



mg/dL
AND
BOTH
of the
following:

- i.
No
clinical
ASCVD
or
CAC
less
than
100
Agatston
units
AND
- ii.
Poorly
controlled
cardiometabolic
risk
factor
AND

- 2. ONE of the following:
 - a. The patient has been
adherent to high-intensity
statin therapy (i.e.,
rosuvastatin greater than or
equal to 20 mg daily,
atorvastatin greater than or
equal to 40 mg daily) for
greater than or equal to 8
continuous weeks AND ONE
of the following:
 - i. The patient’s
LDL-C level after this
treatment regimen
remains greater than
or equal to 70 mg/dL
OR



C from baseline after this treatment regimen **OR**

iii. If the patient has ASCVD, and ONE of the following:

a) The patient's non HDL-C level after this treatment regimen remains greater than or equal to 100 mg/dL **OR**

b) The patient is at very high risk and the patient's LDL-C level after this treatment regimen remains greater than or equal to 55 mg/dL **OR**

b. The patient has been determined to be statin intolerant by meeting one of the following criteria:

i. The patient experienced statin-related rhabdomyolysis **OR**

ii. The patient experienced skeletal-



weakness] or myalgia
[muscle aches,
soreness, stiffness, or
tenderness]) and
BOTH of the
following:

- a) The
skeletal-
related
muscle
symptoms
(e.g.,
myopathy or
myalgia)
occurred
while
receiving
separate
trials of both
atorvastatin
AND
rosuvastatin
(as single-
entity or as
combination
products)
AND
- b) When
receiving
separate
trials of both
atorvastatin
and
rosuvastatin
(as single-
entity or as
combination
products)
the skeletal-
related
muscle
symptoms
(e.g.,
myopathy,
myalgia)



of each
 respective
 statin
 therapy
 (atorvastatin
 AND
 rosuvastatin)
OR

iii. The patient
 experienced
 elevations in hepatic
 transaminase while
 receiving separate
 trials of both
 atorvastatin and
 rosuvastatin (as
 single-entity or as
 combination
 products) **OR**

c. The patient has a
 hypersensitivity to
 atorvastatin AND rosuvastatin
OR

d. The patient has an FDA
 labeled contraindication to
 atorvastatin AND rosuvastatin
OR

B. The patient has another FDA approved
 indication for the requested agent and route
 of administration **OR**

C. The patient has another indication that is
 supported in compendia for the requested
 agent and route of administration **AND**

II. If the patient has an FDA labeled
 indication, ONE of the following:

A. The patient's age is within FDA labeling for
 the requested indication for the requested
 agent **OR**

B. The prescriber has provided information in
 support of using the requested agent for the



... The patient will NOT be using the requested agent in combination with another PCSK9 agent for the requested indication
AND

VI. The patient does NOT have an FDA approved contraindication to the requested agent **AND**

VII. ONE of the following:

- A. The request is for a preferred agent **OR**
- B. The patient has tried and had an inadequate response to the preferred agent **OR**
- C. The patient has an intolerance or hypersensitivity to the preferred agent **OR**
- D. The patient has an FDA labeled contraindication to ALL preferred agents

Length of Approval: 12 months

Renewal Evaluation

Renewal of Target Agent(s) may be considered **medically necessary** when

- I. The patient has been previously approved for the requested agent through the plan's Prior Authorization process **AND**
- II. The patient has had clinical benefit with the requested agent **AND**
- III. The patient has been adherent to requested agent **AND**
- IV. If the patient has cardiovascular disease OR hyperlipidemia, then ONE of the following:
 - A. The patient is currently adherent to high-intensity statin therapy (i.e., rosuvastatin greater than or equal to 20 mg daily, atorvastatin greater than or equal to 40 mg daily) **OR**



1. The patient experienced statin-related rhabdomyolysis **OR**
 2. The patient experienced skeletal-related muscle symptoms (e.g., myopathy [muscle weakness] or myalgia [muscle aches, soreness, stiffness, or tenderness]) and BOTH of the following:
 - a. The skeletal-related muscle symptoms (e.g., myopathy or myalgia) occurred while receiving separate trials of both atorvastatin AND rosuvastatin (as single-entity or as combination products) **AND**
 - b. When receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or as combination products) the skeletal-related muscle symptoms (e.g., myopathy, myalgia) resolved upon discontinuation of each respective statin therapy (atorvastatin AND rosuvastatin) **OR**
 3. The patient experienced elevations in hepatic transaminase while receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or as combination products) **OR**
- C. The patient has a hypersensitivity to atorvastatin AND rosuvastatin **OR**
- D. The patient has an FDA labeled contraindication to atorvastatin AND rosuvastatin **AND**
- V. The patient will NOT be using the requested agent in combination with another



... the patient does not have any FDA-labeled contraindications to the requested agent

Length of Approval: 12 months

Dates

Original Effective
09-02-2015
Last Review
11-06-2024
Next Review
11-12-2025

CLINICAL RATIONALE

Criteria have been developed to aid in diagnosing HeFH. These include the Simon Broome Register criteria and Dutch Lipid clinic Network criteria. (5) Definitive diagnosis of HeFH according to Simon Broome diagnostic criteria requires the patient has one of the following:(3,5)

- Total cholesterol greater than 6.7 mmol/L or low-density lipoprotein cholesterol (LDL-C) greater than 4.0 mmol/L in a child aged younger than 16 years, or greater than 7.5 mmol/L or LDL-C greater than 4.9 mmol/L in an adult (levels either pre-treatment or highest on treatment) **plus** tendon xanthomas in the patient, or in first-degree relative (parent, sibling or child), or in second-degree relative (e.g. grandparent, uncle or aunt) **OR**
- DNA-based evidence of an LDL receptor mutation, familial defective Apo B-100, or a PCSK9 mutation

Possible diagnosis of HeFH according to Simon Broome diagnostic criteria requires the patient has:(3)

- Total cholesterol greater than 6.7 mmol/L or low-density lipoprotein cholesterol (LDL-C) greater than 4.0 mmol/L in a child aged younger than 16 years, or greater than 7.5 mmol/L or LDL-C greater than 4.9 mmol/L in an adult (levels either pre-treatment or highest on treatment) **AND** at least one of the following:
- Family history of myocardial infarction: aged younger than 50 years in second-degree relative or aged younger than 60 years in first-degree relative **OR**
- Family history of raised total cholesterol: greater than 7.5 mmol/L in adult first- or second-degree relative or greater than 6.7 mmol/L in child, brother or sister aged younger than 16 years

The Dutch Lipid Clinic Network criteria assign points based on cholesterol levels, family history of hyperlipidemia or cardiovascular disease, clinical presentation, and/or presence of identified genetic mutation affecting plasma LDL-C. (5-7) A definitive diagnosis of HeFH can be made in patients with greater than 8 points. A probable diagnosis can be made in patients with 6-8 points

Dutch Lipid Clinic Network criteria for diagnosis of heterozygous familial hypercholesterolemia(8)

Group 1: Family history	Points
-------------------------	--------



percentile by age and gender for country	1
<ul style="list-style-type: none"> First-degree relative with tendon xanthoma and/or corneal arcus 	2
<ul style="list-style-type: none"> Children less than 18 years with LDL cholesterol greater than 95th percentile by age and gender for country 	2
Group 2: Clinical history	Points
<ul style="list-style-type: none"> Subject has premature (less than 55 years, men; less than 60 years, women) CHD 	2
<ul style="list-style-type: none"> Subject has premature (less than 55 years, men; less than 60 years, women) cerebral or peripheral vascular disease 	1
Group 3: Physical examination	Points
<ul style="list-style-type: none"> Tendon xanthoma 	6
<ul style="list-style-type: none"> Corneal arcus in a person less than 45 years 	4
Group 4: Biochemical results (LDL-C)	Points
<ul style="list-style-type: none"> greater than 8.5 mmol/L (greater than 325 mg/dL) 	8
<ul style="list-style-type: none"> 6.5–8.4 mmol/L (251–325 mg/dL) 	5
<ul style="list-style-type: none"> 5.0–6.4 mmol/L (191–250 mg/dL) 	3
<ul style="list-style-type: none"> 4.0–4.9 mmol/L (155–190 mg/dL) 	1
Group 5: Molecular genetic testing (DNA analysis)	Points
<ul style="list-style-type: none"> Causative mutation shown in the LDLR, APOB, or PCSK9 genes 	8
Use and Interpretation	
Assign only one score, the highest applicable, per group then add the points from each group to achieve the total score	
Definitive FH diagnosis: greater than 8 points	
Probable FH diagnosis: 6 to 8 points	
Possible FH diagnosis: 3 to 5 points	
Unlikely FH diagnosis: 0 to 2 points	

Guidelines advise that diagnosis of HoFH can be made based on genetic or clinical criteria. Genetic confirmation of the HoFH includes confirmation of two mutant alleles at the LDL-R, APOB, PCSK9, or LDLRAP1 genes.(4,6) While genetic testing may provide a definitive diagnosis of HoFH, it is recognized that in some patients, genetic confirmation remains elusive, despite exhaustive investigation; indeed, the existence of additional FH genes cannot be excluded. Historically, HoFH has been most commonly



age of 20 years, or the presence of untreated elevated LDL-C levels consistent with FH in both parents.(4,6)

According the American Heart Association (AHA), initial treatment for FH should include a high intensity statin.(9) If the LDL-C is not at goal after 3 months of therapy with the high intensity statin and the patient has been adherent, AHA recommends the addition of ezetimibe. For patients who do not respond to this two drug regimen within 3 months, AHA recommends addition of a PCSK9, a bile acid sequestrant, or niacin. Patients with HoFH who require additional therapy despite treatment with the three drug regimen, AHA recommends addition of Juxtapid and LDL apheresis.(9)

The AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline lists the following as clinical ASCVD:

- Acute coronary syndrome (ACS)
- Myocardial infarction (MI)
- Stable or unstable angina or coronary or other arterial revascularization
- Stroke
- Transient ischemic attack (TIA) or peripheral artery disease (PAD) including aortic aneurysm

The AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Blood Cholesterol states the following regarding PCSK9 therapy(9)

- Severe hypercholesterolemia (LDL-C greater than or equal to 190 mg/dL [greater than or equal to 4.9 mmol/L])
 - In patients 30-75 years of age with heterozygous FH and with an LDL-C level of 100 mg/dL (greater than or equal to 2.6 mmol/L) or higher while taking maximally tolerated statin and ezetimibe therapy, the addition of a PCSK9 inhibitor may be considered
 - In patients 40-75 years of age with a baseline LDL-C level of 220 mg/dL (greater than or equal to 5.7 mmol/L) or higher while receiving maximally tolerated statin and ezetimibe therapy, the addition of a PCSK9 inhibitor may be considered

The AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Blood Cholesterol states the following regarding PCSK9 therapy:(9)

- Secondary atherosclerotic cardiovascular disease (ASCVD) prevention
 - In patients with clinical ASCVD who are judged to be very high risk and considered for PCSK9 inhibitor therapy, maximally tolerated LDL-C lowering therapy should include maximally tolerated statin therapy and ezetimibe
 - In patients with clinical ASCVD who are judged to be very high risk and who are on maximally tolerated LDL-C lowering therapy with LDL-C 70 mg/dL (greater than or equal to 1.8 mmol/L) or higher or a non-HDL-C level of 100 mg/dL (greater than or equal to 2.6 mmol/L) or higher, it is reasonable to add PCSK9 inhibitor following a clinical-patient discussion about the net benefit, safety, and cost

The AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline categorizes the following statin intensities:(9)



Lowering	equal to 50%		
Statins	Atorvastatin 40-80 mg	Atorvastatin 10-20 mg	Simvastatin 10 mg
	Rosuvastatin 20-40 mg	Rosuvastatin 5-10 mg	Pravastatin 10-20 mg
		Simvastatin 20-40 mg (a)	Lovastatin 20 mg
		Pravastatin 40-80 mg	Fluvastatin 20-40 mg
		Lovastatin 40-80 mg	
		Fluvastatin XL 80 mg	
		Fluvastatin 40 mg twice daily	
		Pitavastatin 1-4 mg	

a - Although simvastatin 80 mg was evaluated in RCTs, initiation of simvastatin 80 mg or titration to 80 mg is not recommended by the FDA because of the risk of myopathy, including rhabdomyolysis

The National Lipid Association (NLA) 2019 consensus statement identifies the following patients, who are already on maximally tolerated statin therapy, as most likely to benefit from PCSK9 therapy.(10)

- Extreme high-risk (greater than or equal to 40% 10-year ASCVD risk on maximally tolerated statin therapy) with LDL-C greater than or equal to 70 mg/dL and either of the following:
 - Extensive or active burden of ASCVD (i.e., polyvascular ASCVD, which affects all 3 vascular beds—coronary, cerebrovascular, and peripheral arterial; clinical peripheral arterial disease in addition to coronary and/or cerebrovascular disease; a clinical ASCVD event with multivessel coronary artery disease defined as greater than or equal to 40% stenosis in greater than or equal to 2 large vessels; or recurrent myocardial infarction within 2 years of the initial event) in the presence of adverse or poorly controlled cardiometabolic risk factors
 - Extremely high-risk elevations in cardiometabolic factors with less-extensive ASCVD (i.e., HeFH, diabetes, LDL-C greater than or equal to 100 mg/dL, less than high-intensity statin therapy, chronic kidney disease, poorly controlled hypertension, high-



with ASCVD and history of severe hyperlipidemia (SH) LDL-C greater than or equal to 220 mg/dL are an additional group of extremely high-risk patients, with greater than or equal to 45% 10-year ASCVD risk despite statin therapy. Statin-treated HeFH patients with coronary artery calcium (CAC) score greater than 100 Agatston units also have about a 45% 10-year ASCVD risk despite statin therapy

- Very high-risk (30-39% 10-year ASCVD risk on maximally tolerated statin therapy) with LDL-C greater than or equal to 100 mg/dL and the following:
 - Less-extensive clinical ASCVD (i.e., no polyvascular ASCVD, no clinical peripheral arterial disease, a prior ASCVD event greater than or equal to 2 years prior, and no coronary artery bypass grafting)
 - Adverse or poorly controlled cardiometabolic risk factor(s) including age greater than or equal to 65 years, current smoking, chronic kidney disease, lipoprotein(a) greater than or equal to 37 nmol/L, high-sensitivity C-reactive protein 1–3 mg/L, metabolic syndrome with a history of myocardial infarction, ischemic stroke, or symptomatic peripheral arterial disease, usually in the presence of other adverse or poorly controlled cardiometabolic risk factors
- High-risk (20-29% 10-year ASCVD risk on maximally tolerated statin therapy) with LDL-C greater than or equal to 130 mg/dL and either of the following:
 - High-risk patients with ASCVD who have the following:
 - Less-extensive ASCVD
 - Well-controlled cardiometabolic risk factors (i.e., no diabetes, nonsmoker, on high-intensity statin with LDL-C less than 100 mg/dL, blood pressure less than 140/90 mm Hg, and C-reactive protein less than 1 mg/dL)
 - Primary prevention patients with HeFH or SH LDL-C greater than or equal to 220 mg/dL and have the following:
 - No clinical ASCVD or CAC less than 100 Agatston units
 - Poorly controlled cardiometabolic risk factor

CAC Agatston score in non-contrast CT can be used for patient risk classification for coronary heart disease:(11,12)

- 0 CAC = no CAC, very low risk,
- 1-99 CAC = mild CAC, mildly increased risk
- 100 - 299 CAC = moderate CAC, moderately increased risk
- greater than or equal to 300 CAC = moderate to severely increased risk

In their 2022 Expert Consensus Decision, the American College of Cardiology stated that in view of the favorable net clinical benefit of the addition of nonstatin therapies in patients with clinical ASCVD at very high risk on high-intensity statin therapy and lifestyle management and the very low levels of LDL-C achieved in RCTs (randomized clinical trials) of nonstatin therapies, a lower LDL-C threshold of LDL-C less than or equal to 55 mg/dL is recommended. Patients in this group have a history of multiple major ASCVD events (e.g., recent [within the past 12 months] acute coronary syndrome, history of MI other than the recent ACS event listed previously, history of ischemic stroke, and



chronic kidney disease, current smoking, persistently elevated LDL-C despite maximally tolerated statin therapy and ezetimibe, and history of congestive heart failure. (13)

Quick Code Search

Use this feature to find out if a procedure and diagnosis code pair will be approved, denied or held for review. Simply put in the procedure code, then the diagnosis code, then click "Add Code Pair". If the codes are listed in this policy, we will help you by showing a dropdown to help you.

Procedure

Please type a procedure code

Enter at least the first 3 characters of the code

Diagnosis

Please type a diagnosis code

Enter at least the first 3 characters of the code

Add

CODES

+ HCPCS

REFERENCES



2021

Repatha prescribing information. Amgen. September 2021

2024

Identification and Management of Familial Hypercholesterolemia. Simon Broome Diagnostic criteria for index individuals and relatives.:<http://www.ncbi.nlm.nih.gov/books/NBK53810/>.

2014

Homozygous familial hypercholesterolaemia: new insights and guidance for clinicians to improve detection and clinical management. A position paper from the Consensus Panel on Familial Hypercholesterolaemia of the European Atherosclerosis Society. July 22, 2014.
<https://academic.oup.com/eurheartj/article/35/32/2146/2481389>.

2008

National Collaborating Centre for Primary Care (UK). Identification and Management of Familial Hypercholesterolaemia (FH) [Internet]. London: Royal College of General Practitioners (UK); 2008 Aug. (NICE Clinical Guidelines, No. 71.) 3, Diagnosis. <http://www.ncbi.nlm.nih.gov/books/NBK53817/>

1999

World Health Organization. Familial Hypercholesterolaemia (FH): Report of a second WHO consultation. Geneva: World Health Organization; 1999

2013

Nordestgaard BG, Chapman MJ, Humphries ST, et al; for the European Atherosclerosis Society Consensus Panel. Familial hypercholesterolaemia is underdiagnosed and undertreated in the general population: guidance for clinicians to prevent coronary heart disease. *Eur Heart J*. 2013. doi.10.1093/eurheartj/eh273.

2015

Gidding S, Champagne M, Ferranti S, et al. The Agenda for Familial Hypercholesterolemia. A Scientific Statement From the American Heart Association. *Circulation*. 2015; 132:00-00

2018

2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol. Journal of the American College of Cardiology. <https://doi.org/10.1016/j.acc.2018.11.003>.

2019



Lipidology. July-August, 2019. 13(4): 525-537.

2017

Hect HS, Cronin P, Blaha M, et al. 2016 SCCT/STR guidelines for coronary artery calcium scoring of noncontrast noncardiac chest CT scans: A report of the Society of Cardiovascular Computed Tomography and Society of Thoracic Radiology. J Thorac Imaging. 2017;32(5):W54-S66

2017

Blaha MJ, Mortensen MB, Kianoush S, et al. Coronary artery calcium scoring. Is it time for a change in methodology. J Am Coll Cardiol Imag. 2017;10:923-937.

2022

2022 ACC Expert Consensus Decision Pathway on the Role of Nonstatin Therapies for LDL-Cholesterol Lowering in the Management of Atherosclerotic Cardiovascular Disease Risk. Journal of the American College of Cardiology, 80 (14) 2022, 1366-1418. https://www.sciencedirect.com/science/article/pii/S0735109722055942?ref=pdf_download&fr=RR-2&rr=7edef3f3691926bb

REVISIONS

10-01-2024

Removal of specialist criteria per trade contracts.

12-01-2023

Policy reviewed at Medical Policy Committee meeting on 11/8/2023 – no changes to policy.

06-30-2017

Adding indication of clinical atherosclerotic cardiovascular disease (ASCVD) to policy

01-09-2017

Policy criteria revised

HAVE AN IDEA? WE'RE HERE TO HELP YOU MANAGE YOUR WORK



[Privacy](#) [Legal](#) [Non-Discrimination and Translation](#) [Site Map](#)

© 2025 Blue Cross and Blue Shield of Nebraska.

Blue Shield of Nebraska is an independent licensee of the Blue Cross and Blue Shield Association. The Blue Cross and Blue Shield Association licenses Blue Cross and Blue Shield of Nebraska to offer certain products and services under the Blue Cross® and Blue Shield® brand names within the state of Nebraska.