Test

Kate Che

Contents

Introduction	1
Hypothesis	1
Analysis	1
Bibliography	2

Introduction

The data used in study is taken from https://open.canada.ca/en/open-data¹. The algorithm is taken from Antonov and Ermakov (2015).

The data set *contains* 518 measurements of 6 Canadian glacier mass balance, collected from 1960 till 2007. Namely, the file includes these glaciers:

- ## [1] "Devon Ice Cap NW Devon Island, Nunavut"
- ## [2] "Helm Glacier southern Coast Mountains (Garibaldi Provincial Park), British Columbia"
- ## [3] "Meighen Ice Cap Meighen Island, Nunavut"
- ## [4] "Peyto Glacier Rocky Mountain eastern slopes (Banff National Park), Alberta"
- ## [5] "Place Glacier southern Coast Mountains, British Columbia"
- ## [6] "White Glacier Axel Heiberg Island, Nunavut"

Hypothesis

We are interested in findong out whether there is a statistically significant cange in mass balance over the observed time period. For these purposes we use \mathbf{R} (version 3.6.1) and an appropriate statistical test called t-test:

$$t = \frac{\overline{x} - \mu_0}{s/\sqrt{n}}.$$

The workflow is as follow:


- 1. Read file
- 2. Run t-test for each glacier and collect p-values
- 3. Support the evidence with
 - a table of results;
 - a plot thet could help demonstrate the effect.

Analysis

Table 1: Descriptive statistics and confidence intervals

Name	YearsObserved	MeanChange	WorstChange	WorstYear	PValue	ConfidenceLimit
Devon Ice Cap NW	47	-91.2	-559	2001	5.81e-05	-39.0
Helm Glacier	31	-1277.3	-2850	1998	1.73e-07	-798.0
Meighen Ice Cap	48	-107.6	-970	1962	4.51e-03	-12.5
Peyto Glacier	42	-579.9	-2230	1998	3.62e-07	-339.7
Place Glacier	43	-861.4	-2486	1995	3.70e-09	-572.3
White Glacier	48	-152.4	-818	2007	$6.56\mathrm{e}\text{-}05$	-64.3

¹Here is the direct link to data download.



Bibliography

Antonov, A. A., and S. M. Ermakov. 2015. "Random Cubatures and Quasi-Monte Carlo Methods." *Monte Carlo Methods and Applications* 21 (3): 179–87. https://doi.org/10.1515/mcma-2015-0102.