



NGS introduction

Joana Meier

Sequencing costs have decreased massively





History of Illumina sequencing



High Throughput Sequencing (=Next Generation Sequencing)

- **Short-read sequencing technologies (2nd generation):**
 - Sequence millions of clonally amplified molecules
 - E.g. Illumina, Ion Torrent, SOLiD
- **Long-read technologies (3rd generation):**
 - Single molecules are sequenced in real-time, fast but expensive and high error rates
 - E.g. PacBio (bought by Illumina): ~12kb reads, single molecules are read multiple times to reduce error rate
 - E.g. Oxford Nanopore: up to 900 kb reads, high sequencing error rate (5-15%) and non-random errors, each DNA fragment can only be read 2x


Sanger Sequencing




- Manually check each sequence
- Resequenced failed sequences

Illumina flowcell: millions of DNA sequences



DNA fragments with Illumina adapters




Each lane contains a dense lawn of Illumina primers



Sequencing by synthesis by Illumina



Illumina HiSeq4000 and X Ten




Problem:
Illumina barcode switching
(Index hopping)

-> use double-indexing
Different P1 and P2 indices


Long read sequencing technologies

Nanopore




PacBio


Introduction to SMRT Sequencing



Each of the four nucleotides is labeled with a different colored fluorophore




Whole-genome sequencing (shotgun sequencing)



What kind of genomic data will/are you working on?

Answered: 18 Skipped: 1



RAD sequencing

Restriction Associated DNA sequencing

Restriction enzyme
(e.g. *SbfI*)



5'-TGCAGTGC GGTGGTCACCTGCA|GGCCGTGCGTGCTAGCAGTGC GGT...
3'-ACGTCACGCCACCAGTGG|ACGTCCGGCAGGCACGATCGTCACGCCA...

fPCRprimer-IlluminaPrimer-barcode-TGCA
fPCRprimer-IlluminaPrimer-barcode-P


5'-P1-barcode-TGCAGGTCCGTGCGTGCTAG...A P2-GAGAACA
3'-P1-barcode-ACGTCCAGGCACGCACGATC... T-P2-CACGATACGGCAGAAGACGAAC

PCR →

complement to reverse
PCR primer binding site


RAD sequencing

Restriction **A**ssociated **D**NA sequencing



Other «reduced-representation» techniques

- **CRoPS/ddRAD sequencing** (double-digest restriction enzyme and size selection instead of size selection)
- **GBS** (genotyping by sequencing): no shearing, sequencing select short fragments
- **UCE**: Selection of DNA fragments through sequencing on ultraconserved elements (conserved across species)
- **2b-RAD**: type IIB restriction enzymes that cut downstream of the restriction site
- **Transcriptome sequencing**: RNAseq, only cDNA to annotate, depth informative on expression
- **Targeted resequencing**: Sequence capture




Trade-offs: Splitting reads (i.e. costs) among:

- Number of sites to sequence
- Number of samples
- Depth of coverage
- Example: 1 HiSeq2500 flow cell (about 1000 Euro) ~250 mio read pairs of 125 bp each -> 75 Gb data
 - 5 whole-genomes of a species with 1 Gb genome size at 15x coverage
 - 50 whole-genomes of a species with 500 Mb genome size at 3x coverage
 - 30 Mbp sequenced for 100 samples with a reduced-representation technique at a sequencing depth of 25


Considerations in choosing the library preparation and sequencing techniques

- Research question and planned analyses
- Genome size
- Availability & quality of reference genome (no ref genome -> not wgs)
- Available budget
- Number of samples to sequence (tradeoff with sequencing depth)
- Amounts of DNA available
- Sequencing depth aimed at
- SNP density required
- Divergence between samples
- Heterozygosity of samples
- Phase required
- Accuracy of each single position (if high needed, avoid PCR-based methods)
- Importance of annotations
- Neutral dataset or specific regions wanted

Fastq format



Quality scores



$$\text{Phred} = -10 \log_{10} p$$

p = Probability call is incorrect

ASCII encoding

Quality Score	Probability of incorrect base call	Base call accuracy
10	1 in 10	90%
20	1 in 100	99%
30	1 in 1000	99.9%
40	1 in 10000	99.99%

40 : @

41 : A

42 : B

43 : C

44 : D

45 : E

... : ...

90 : Z

91 : [

92 : \

93 :]

94 : ^

95 : _

... : ...

141 : a

142 : b

143 : c

144 : d

145 : e

146 : f


... : ...

Read header


```
@EAS139:136:FC706VJ:2:2104:15343:197393 1:Y:18:ATCACG
```

EAS139	the unique instrument name
136	the run id
FC706VJ	the flowcell id
2	flowcell lane
2104	tile number within the flowcell lane
15343	'x'-coordinate of the cluster within the tile
197393	'y'-coordinate of the cluster within the tile
1	the member of a pair, 1 or 2 (<i>paired-end or mate-pair reads only</i>)
Y	Y if the read is filtered, N otherwise
18	0 when none of the control bits are on, otherwise it is an even number
ATCACG	index sequence

FastQC: Quality across bases (good example)




FastQC: Quality across bases (bad example)



Let's have a look at the first few sequences and check the sequencing quality with fastqc


RAD/GBS



Each read: will start with the barcode, then the restriction site, then a variable sequence


Issues with cluster identification

Due to low complexity at the beginning of the sequence,
Illumina cannot distinguish if a signal comes from one or two clusters




Phasing issues

“In Phase”





“Out of Phase”



The first 12 nucleotides are also used for «phasing», i.e. correcting for reads that are out of phase. The algorithm expects random nucleotide distribution!

-> Barcodes of the same length may lead to low quality overall

Per base sequence content




How to minimize the problem


- Use barcodes of different lengths to shift the restriction enzyme cut site
- Add PhiX virus DNA to the RAD libraries to increase the complexity of reads ('spiking')
- Reduce loading concentrations of Illumina plates
- Potentially: filter out bad reads

Quality scores across bases: Whole genome sequencing (PCR free library prep)


forward



reverse




Sequence duplication level



**Duplication level:
= Percentage of
reads that have x
copies**

Sequence duplication level


Example 1: 20 reads total
10 unique sequences + 5 sequences each present twice



Deduplicated sequences (=number of distinct copies)

15 distinct sequences are distributed as 10 singletons and 5 duplicates, $10/15=66\%$ and $5/15=33\%$ is the slope of the red line. Thus $15/20=75\%$ remaining after deduplication (distinct reads).

Example 2: 20 reads total
10 unique sequences + 1 sequence present 10x




Deduplicated sequences (=number of distinct copies)


11 total groups where $10/11=91\%$ are singletons and $1/11=9\%$ of the groups form at duplication rate of 10x. Therefore, $11/20 = 55\%$ distinct reads.

Per base sequence content


RAD1



RAD2




RAD/GBS



Each read: will start with the barcode, then the restriction site, then a variable sequence


Issues with cluster identification

Due to low complexity at the beginning of the sequence,
Illumina cannot distinguish if a signal comes from one or two clusters




Phasing issues

“In Phase”





“Out of Phase”



The first 12 nucleotides are also used for «phasing», i.e. correcting for reads that are out of phase. The algorithm expects random nucleotide distribution!

-> Barcodes of the same length may lead to low quality overall

Per base sequence content




How to minimize the problem


- Use barcodes of different lengths to shift the restriction enzyme cut site
- Add PhiX virus DNA to the RAD libraries to increase the complexity of reads ('spiking')
- Reduce loading concentrations of Illumina plates
- Potentially: filter out bad reads

GC distribution over all sequences



forward





reverse



GC distribution over all sequences





GC distribution over all sequences




Fastqc: Per sequence GC content

! Per sequence GC content




Sharp peak indicates specific motif. Adapters are the usual suspect.





Wider or multiple distributions suggest contamination.

Per base N content




Sequence Length Distribution




Sequence duplication level


wgs



RAD




Adapter content




In wgs datasets
not even shown

Kmer content

Most common barcodes + cut site



Most common barcodes + cut site



GBS

How many SNPs will I get?

Species	Genome Size (Mb)	Enzyme	Sample Size	No. SNPs
Maize	2,600	<i>ApeKI</i>	33,000	1,200K
Rice	400	<i>ApeKI</i>	850	60K
Grape	500	<i>ApeKI</i>	1000	200K
Willow*	460	<i>ApeKI</i>	459	23K
Pine*	16,000	<i>ApeKI</i>	12	63K
Vole*	3,400	<i>PstI</i>	283	53K
Fox*	2,400	<i>EcoT22I</i>	48	16K
Cow	3,000	<i>PstI</i>	48	64K
<i>Verticilliflorum</i> (fungus isolates)	40	<i>ApeKI</i>	2	10K

*No reference genome. UNEAK analysis pipeline used for analysis. To avoid homology/paralogy issues this pipeline calls SNPs very conservatively.