

Discrete Mathematics: Lecture 26

Paths and Isomorphism, Counting Paths, Euler Paths and Circuits

Xuming He

Associate Professor

School of Information Science and Technology
ShanghaiTech University

Spring Semester, 2022


Notes by Prof. Liangfeng Zhang

Vertex Connectivity

DEFINITION: A connected undirected graph $G = (V, E)$ is said to be **nonseparable**_{不可分的} if G has no cut vertex.

DEFINITION: Let $G = (V, E)$ be a connected simple graph.

- **vertex cut**_{点割集}: A subset $V' \subseteq V$ such that $G - V'$ is disconnected
- **vertex connectivity**_{点连通度} $\kappa(G)$: the minimum number of vertices whose removal disconnect G or results in K_1 ; equivalently,
 - if G is disconnected, $\kappa(G) = 0$; //additional definition
 - if $G = K_n$, $\kappa(G) = n - 1$ // K_n has no vertex cut
 - else, $\kappa(G)$ is the minimum size of a vertex cut of G



These graphs are all nonseparable


Edge Connectivity

DEFINITION: Let $G = (V, E)$ be a connected simple graph. $E' \subseteq E$ is an **edge cut** 边割集 of G if $G - E'$ is disconnected.

DEFINITION: Let $G = (V, E)$ be a simple graph.


The **edge connectivity** 边连通度 ($\lambda(G)$) of G is defined as below:

- G disconnected: $\lambda(G) = 0$
- G connected:
 - $|V| = 1$: $\lambda(G) = 0$
 - $|V| > 1$: $\lambda(G)$ is the minimum size of edge cuts of G .



Connectivity

THEOREM: Let $G = (V, E)$ be a simple graph. Then $\kappa(G) \leq \lambda(G) \leq \delta(G)$, where $\delta(G) = \min_{v \in V} \deg(v)$ is the least degree of G 's vertices.



- $\kappa(G) = 2$
- $\lambda(G) = 3$
- $\delta(G) = 4$

https://cp-algorithms.com/graph/edge_vertex_connectivity.html

<http://www.math.caltech.edu/~2014-15/2term/ma006b/05%20connectivity%201.pdf>


Paths and Isomorphism

(Simple graph)

Theorem

The existence of a simple circuit of length k , $k \geq 3$ is an isomorphism invariant for simple graphs.

3 edge
circuit




G_1

6 vertices, 8 edges

Degree sequence: 3, 3, 3, 3, 2, 2

4 edge
circuit

→ not iso-



G_2

Circuit

→ iso mo. Var

Paths and Isomorphism*

Theorem

The existence of a simple circuit of length k , $k \geq 3$ is an isomorphism invariant for simple graphs.

Proof: Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be isomorphic graphs: there is a bijective function $f : V_1 \rightarrow V_2$ respecting adjacency conditions.

Assume G_1 has a simple circuit of length k : $u_0, u_1, \dots, u_k = u_0$, with $u_i \in V_1$ for $0 \leq i \leq k$. Let's denote $v_i = f(u_i)$, for $0 \leq i \leq k$.

$(u_i, u_{i+1}) \in E_1 \Rightarrow (f(u_i), f(u_{i+1})) = (v_i, v_{i+1}) \in E_2$, for $0 \leq i \leq k-1$.

So v_0, \dots, v_k is a path of length k in G_2 .


It is a circuit because $v_k = f(u_k) = f(u_0) = v_0$.

It is simple: if not, at least one edge is traversed more than once, so it would mean that there exist $0 \leq i \neq j \leq k-1$ such that


$(v_i, v_{i+1}) = (v_j, v_{j+1})$. But this implies $(u_i, u_{i+1}) = (u_j, u_{j+1})$ by bijectivity of f . This is impossible because u_0, u_1, \dots, u_k is simple.

A way of finding BIJ

BIJ NOT only 1



G



H

5 vertices, 6 edges

Degree sequence: 3, 3, 2, 2, 2

1 simple circuit of length 3,

1 simple circuit of length 4,

1 simple circuit of length 5.

Isomorphic graphs ?

If there is an iso $f : V_G \rightarrow V_H$, the simple circuit of length 5

u_1, u_4, u_3, u_2, u_5 must be sent to the simple circuit of length 5 in H, respecting the degrees of vertices.

Check that $f(u_1) = v_1, f(u_4) = v_2, f(u_3) = v_3, f(u_2) = v_4, f(u_5) = v_5$ is an isomorphism by writing adjacency matrices.

Counting Paths Between Vertices

Theorem

Let G be a graph with adjacency matrix A with respect to the ordering of vertices v_1, \dots, v_n . The number of different paths of length $r \geq 1$ from v_i to v_j equals the (i, j) entry of the matrix A^r .

A^r

Proof: By induction

- $r = 1$: the number of paths of length 1 from v_i to v_j is equal to the (i, j) entry of A by definition of A , as it corresponds to the number of edges from v_i to v_j .

- Assume the (i, j) entry of the matrix A^r is the number of different paths of length r from v_i to v_j .

We can write $A^{r+1} = A^r A$

Let's denote $A^r = (b_{ij})_{1 \leq i, j \leq n}$, and $A = (a_{ij})_{1 \leq i, j \leq n}$. The (i, j) entry of A^{r+1} is given by:

$$\sum_{k=1}^n b_{ik} a_{kj} = b_{i1} a_{1j} + b_{i2} a_{2j} + \cdots + b_{in} a_{nj} \quad (1)$$


By hypothesis: b_{ik} equals the number of paths of length r from v_i to v_k .

"Path of length $r + 1$ from v_i to v_j = path of length r from v_i to any vertex v_k + an edge from v_k to v_j ."

This is equal to the sum (1).

Example

How many paths of length four are there from a to d in the simple graph G



with ordering of vertices (a, b, c, d, \cancel{e}) :

$$A_G = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix}$$

$r=1$ path

$r=2$ path. G $a \rightarrow b \rightarrow a$
 $a \rightarrow c \rightarrow a$

$$A_G^2 = \begin{pmatrix} 2 & 0 & 0 & 2 \\ 0 & 2 & 2 & 0 \\ 0 & 2 & 2 & 0 \\ 2 & 0 & 0 & 2 \end{pmatrix}$$

$$A_G^3 = \begin{pmatrix} 0 & 4 & 4 & 0 \\ 4 & 0 & 0 & 4 \\ 4 & 0 & 0 & 4 \\ 0 & 4 & 4 & 0 \end{pmatrix}$$




$$A_G^4 = \begin{pmatrix} 8 & 0 & 0 & 8 \\ 0 & 8 & 8 & 0 \\ 0 & 8 & 8 & 0 \\ 8 & 0 & 0 & 8 \end{pmatrix}$$

Euler Paths and Circuits

一次 回路



DEFINITION: Let $G = (V, E)$ be a graph.

- **Euler Path** 欧拉路径: a simple path that traverses every edge of G .
- **Euler Circuit** 欧拉回路: a simple circuit that traverses every edge of G .




path

not simple



circuit



Remark: When G has multiple edges, these edges will be given different names and considered as different. This is implicit in the textbook.

Circuit Euler Circuits

所有顶点

THEOREM: Let $G = (V, E)$ be a connected multigraph of order ≥ 2 . Then G has an Euler circuit iff $2 \mid \deg(x)$ for every $x \in V$.

- \Rightarrow : Let $P: \{x_0, x_1\}, \dots, \{x_{i-1}, x_i\}, \dots, \{x_{n-1}, x_n\}$ be an Euler circuit, $x_0 = x_n$
 - Every occurrence of x_i in P contributes 2 to $\deg(x_i)$
 - Every vertex x_i has an even degree
 - \Leftarrow : Let $P: \{x_0, x_1\}, \dots, \{x_{n-1}, x_n\}$ be a longest simple path in G .
 - Let $H = G[P]$, the subgraph of G induced by all edges in P
 - If $x_n \neq x_0$, then $\deg_H(x_n)$ is odd and so P cannot be longest.
 - $x_n = x_0$, P is a simple circuit, and $2 \mid \deg_H(x_i)$ for all i .
 - If $\exists i \in \{0, 1, \dots, n-1\}$ such that $\deg_H(x_i) < \deg_G(x_i)$,
 - then $\exists y \in V$ such that $\{x_i, y\} \notin P$
 - $y, x_i, x_{i+1}, \dots, x_n, x_1, \dots, x_{i-1}, x_i$ is longer than P
- Hence, $\deg_H(x_i) = \deg_G(x_i)$ for all $i \in \{0, 1, \dots, n-1\}$.
- $V = \{x_0, x_1, \dots, x_{n-1}\}$ and $H = G$.
 - P is an Euler circuit.

Remark: H contains all vertices of G . Otherwise, P can be extended.

maximum

path

\Downarrow

circuit




Construction

How to find?

ALGORITHM (Hierholzer):


- **Input:** $G = (V, E)$, a connected multigraph, $2 \mid \deg(x), \forall x \in V$
- **Output:** an Euler circuit
 - **circuit:** = a circuit in G
 - $H := G - \text{circuit} - \text{isolated vertices}$
 - while H has edges do
 - **subcircuit:** = a circuit in H that intersects **circuit**
 - $H := H - \text{subcircuit} - \text{isolated vertices}$
 - **circuit:** = **circuit** \cup **subcircuit**
 - return **circuit**

Example







small circuit

circuit = a, b, e, a



H

Example







circuit = a, b, e, a

H

subcircuit = a, c, f, a

Example



circuit 1





circuit 2

circuit = a, b, e, a

subcircuit = a, c, f, a

H




Example




circuit = a, b, e, a, c, f, a

H

Example






circuit = a, b, e, a, c, f, a



H

subcircuit = c, d, e, f, b, c

Example






"Insert"

circuit = $a, b, e, a, \textcircled{c}, f, a$


H

subcircuit = c, d, e, f, b, c

Example



circuit = $a, b, e, a, c, d, e, f, b, c, f, a$





Euler Paths

THEOREM: Let $G = (V, E)$ be a connected multigraph of order ≥ 2 . Then G has an Euler path (not Euler circuit) iff G has exactly 2 vertices of odd degree.

ALGORITHM:

- Input:** $G = (V, E)$, a connected multigraph, $x, y \in V$ have odd degrees
- Output:** an Euler path
 - $H := G + \{x, y\}$
 - find an Euler circuit using Hierholzer's algorithm
 - remove the edge $\{x, y\}$ from the circuit



circuit - additional edge of 2 odd = path

$a, c, d, e, f, b, a, e, b, c, f, a$

$a, c, d, e, f, b, a, e, b, c, f, a$


$a, e, b, c, f, a, c, d, e, f, b$




Hamilton Paths and Circuits

DEFINITION: Let $G = (V, E)$ be a graph.


- **Hamilton Path:** A simple path that passes through every vertex exactly once.
- **Hamilton Circuit:** A simple circuit that passes through every vertex exactly once.




✓ Hamilton path
✓ Hamilton circuit



✓ Hamilton path
✓ Hamilton circuit



✓ Hamilton path
× Hamilton circuit



× Hamilton path
× Hamilton circuit

Hamilton Circuits

NP-C

Determine if there is a Hamilton circuit in a given graph G ?

- This problem is NP-Complete. //that means very difficult

Necessary conditions on Hamilton circuit.

- If G has a vertex of degree 1, then G cannot have a Hamilton circuit.
- If G has a vertex of degree 2, then a Hamilton circuit of G traverses both edges.

Sufficient conditions on Hamilton circuit.



不直接连 u, v

- **Ore's Theorem:** Let $G = (V, E)$ be a simple graph of order $n \geq 3$. If $\deg(u) + \deg(v) \geq n$ for all $\{u, v\} \notin E$, then G has a Hamilton circuit.
- **Dirac's Theorem:** Let $G = (V, E)$ be a simple graph of order $n \geq 3$. If $\deg(u) \geq n/2$ for every $u \in V$, then G has a Hamilton circuit.
 - This is a corollary of Ore's Theorem
 - $\forall u \in V, \deg(u) \geq n/2 \Rightarrow \forall u, v \in V, \deg(u) + \deg(v) \geq n$





4/11/18

Hamilton Circuits

- Examples (sufficient condition)



Hamilton Circuits



Remark: Dirac's and Ore's Theorems do not give a necessary condition for the existence of a Hamilton circuit!


Dirac, Ore just 充分

Shortest Path Problem

Definition

A **weighted graph** is a graph $G = (V, E)$ such that each edge is assigned with a strictly positive number.

The **length** of a path in weighted graph is the sum of the weights of the edges of this path.



a, b, c is a path of length 7 and b, e, d, c is a path of length 6

Remark: Observe that in a non-weighted graph the length of a path is the number of edges in the path!