

Discrete Mathematics: Lecture 18

logically equivalent, rule of replacement, tautological implications

Xuming He

Associate Professor

School of Information Science and Technology
ShanghaiTech University

Spring Semester, 2022

Notes by Prof. Liangfeng Zhang


(but low)

(Continue from last lec)

Logically Equivalent

THEOREM: Let $A^{-1}(\mathbf{T})$ be the set of truth assignments such that A is true. Then $A \equiv B$ if and only if $A^{-1}(\mathbf{T}) = B^{-1}(\mathbf{T})$.

- $A \equiv B$ if and only if $A^{-1}(\mathbf{F}) = B^{-1}(\mathbf{F})$ → Just a sign



Proving $A \equiv B$

EXAMPLE: $P \wedge Q \equiv Q \wedge P$

//commutative law

- Idea: Show that $A^{-1}(\mathbf{T}) = B^{-1}(\mathbf{T})$.
- $A = P \wedge Q; B = Q \wedge P$
 - $A = \mathbf{T}$ if and only if $(P, Q) = (\mathbf{T}, \mathbf{T})$
 - $A^{-1}(\mathbf{T}) = \{(\mathbf{T}, \mathbf{T})\}$
 - $B = \mathbf{T}$ if and only if $(Q, P) = (\mathbf{T}, \mathbf{T})$
 - $B^{-1}(\mathbf{T}) = \{(\mathbf{T}, \mathbf{T})\}$
- $A^{-1}(\mathbf{T}) = B^{-1}(\mathbf{T})$
- $A \equiv B$

REMARK: $P \wedge (Q \wedge R) \equiv (P \wedge Q) \wedge R$ can be shown similarly.

- **Associative law**

Proving $A \equiv B$

EXAMPLE: $P \vee Q \equiv Q \vee P$

//commutative law

- Idea: Show that $A^{-1}(\mathbf{F}) = B^{-1}(\mathbf{F})$.
- $A = P \vee Q; B = Q \vee P$ *prove with \mathbf{F} better*
 - $A = \mathbf{F}$ if and only if $(P, Q) = (\mathbf{F}, \mathbf{F})$
 - $A^{-1}(\mathbf{F}) = \{(\mathbf{F}, \mathbf{F})\}$
 - $B = \mathbf{F}$ if and only if $(Q, P) = (\mathbf{F}, \mathbf{F})$
 - $B^{-1}(\mathbf{F}) = \{(\mathbf{F}, \mathbf{F})\}$
- $A^{-1}(\mathbf{F}) = B^{-1}(\mathbf{F})$
- $A \equiv B$

REMARK: $P \vee (Q \vee R) \equiv (P \vee Q) \vee R$ can be shown similarly.

- **Associative law**

Tautological Implications

DEFINITION: Let A and B be WFFs in propositional variables p_1, \dots, p_n .

- A **tautologically implies** (重言蕴涵) B if every truth assignment that causes A to be true causes B to be true.

- Notation: $A \Rightarrow B$, called a **tautological implication**

- $A^{-1}(\mathbf{T}) \subseteq B^{-1}(\mathbf{T}); B^{-1}(\mathbf{F}) \subseteq A^{-1}(\mathbf{F})$

THEOREM: $A \Rightarrow B$ iff $A \rightarrow B$ is a tautology.

- $A \Rightarrow B$ iff $A^{-1}(\mathbf{T}) \subseteq B^{-1}(\mathbf{T})$ iff $A \rightarrow B$ is a **tautology**

THEOREM: $A \Rightarrow B$ iff $A \wedge \neg B$ is a contradiction.

- $A \rightarrow B \equiv \neg A \vee B \equiv \neg(A \wedge \neg B)$

Proving $A \Rightarrow B$: (1) $A^{-1}(\mathbf{T}) \subseteq B^{-1}(\mathbf{T})$; (2) $B^{-1}(\mathbf{F}) \subseteq A^{-1}(\mathbf{F})$;
(3) $A \rightarrow B$ is a tautology; (4) $A \wedge \neg B$ is a contradiction

tautologically imply
tautology

diff

\Rightarrow

\rightarrow

\neg for all truth
assign

\nwarrow
 \nearrow

Proving $A \Rightarrow B$

$B = q$

since no matter P .

EXAMPLE: Show the tautological implication " $p \wedge (p \rightarrow q) \Rightarrow q$ ".

- Let $A = p \wedge (p \rightarrow q)$; $B = q$. Need to show that " $A \Rightarrow B$ "
- $A^{-1}(T) = \{(T, T)\}$; $B^{-1}(T) = \{(T, T), (\underline{F}, T)\}$: $A^{-1}(T) \subseteq B^{-1}(T)$.

p	q	$p \rightarrow q$	A	B
T	T	T	T	T
T	F	F	F	F
F	T	T	F	T
F	F	T	F	F

- $A \rightarrow B \equiv \neg(p \wedge (p \rightarrow q)) \vee q$
 iff $\equiv (\neg p \vee \neg(p \rightarrow q)) \vee q$
 $A \rightarrow B$ $\equiv (\neg p \vee q) \vee \neg(p \rightarrow q)$
 is T $\equiv (p \rightarrow q) \vee \neg(p \rightarrow q)$
 $\equiv T$
- $A \wedge \neg B \equiv (p \wedge (p \rightarrow q)) \wedge \neg q$
 iff $\equiv (\neg q \wedge p) \wedge (p \rightarrow q)$
 $A \wedge \neg B$ $\equiv \neg(p \rightarrow q) \wedge (p \rightarrow q)$
 is F $\equiv F$

① $A = \dots$ $B = \dots$

② $A \rightarrow B$ T / $A \wedge \neg B$ F / {t, f}

Tautology

Tautological Implications

Name	Tautological Implication	NO.
Conjunction(合取)	$(P) \wedge (Q) \Rightarrow P \wedge Q$	1
Simplification(化简)	$P \wedge Q \Rightarrow P$	2
Addition(附加)	$P \Rightarrow P \vee Q$	3
Modus ponens(假言推理)	$P \wedge (P \rightarrow Q) \Rightarrow Q$	4
Modus tollens(拒取)	$\neg Q \wedge (P \rightarrow Q) \Rightarrow \neg P$	5
Disjunctive syllogism(析取三段论)	$\neg P \wedge (P \vee Q) \Rightarrow Q$	6
Hypothetical syllogism(假言三段论)	$(P \rightarrow Q) \wedge (Q \rightarrow R) \Rightarrow (P \rightarrow R)$	7
Resolution (归结)	$(P \vee Q) \wedge (\neg P \vee R) \Rightarrow Q \vee R$	8

Proofs for 5 and 6

EXAMPLE: $\neg Q \wedge (P \rightarrow Q) \Rightarrow \neg P$

- $A = \neg Q \wedge (P \rightarrow Q), B = \neg P.$
- $$\begin{aligned} A \rightarrow B &\equiv \neg(\neg Q \wedge (P \rightarrow Q)) \vee \neg P \\ &\equiv (Q \vee \neg(P \rightarrow Q)) \vee \neg P \\ &\equiv (\neg P \vee Q) \vee \neg(P \rightarrow Q) \\ &\equiv \mathbf{T} \end{aligned}$$
 $\xrightarrow{\neg(17 \vee Q)}$

EXAMPLE: $\neg P \wedge (P \vee Q) \Rightarrow Q$

- $A = \neg P \wedge (P \vee Q), B = Q.$
- $$\begin{aligned} A \rightarrow B &\equiv \neg(\neg P \wedge (P \vee Q)) \vee Q \\ &\equiv (P \vee \neg(P \vee Q)) \vee Q \\ &\equiv (\neg(P \vee Q) \vee P) \vee Q \\ &\equiv \neg(P \vee Q) \vee (P \vee Q) \\ &\equiv \mathbf{T} \end{aligned}$$

Proofs for 7 and 8

EXAMPLE: $(P \rightarrow Q) \wedge (Q \rightarrow R) \Rightarrow P \rightarrow R$

- $A = (P \rightarrow Q) \wedge (Q \rightarrow R); B = (P \rightarrow R).$
- $A \wedge \neg B \equiv ((\neg P \vee Q) \wedge (\neg Q \vee R)) \wedge (P \wedge \neg R)$
 $\equiv ((\neg P \vee Q) \wedge P) \wedge ((\neg Q \vee R) \wedge \neg R)$
 $\equiv ((\neg P \wedge P) \vee (Q \wedge P)) \wedge ((\neg Q \wedge \neg R) \vee (R \wedge \neg R))$
 $\equiv (Q \wedge P) \wedge (\neg Q \wedge \neg R)$
 $\equiv \mathbf{F}$

EXAMPLE: $(P \vee Q) \wedge (\neg P \vee R) \Rightarrow Q \vee R$

- $A = (P \vee Q) \wedge (\neg P \vee R); B = (Q \vee R).$
- $A \wedge \neg B \equiv (P \vee Q) \wedge (\neg P \vee R) \wedge (\neg Q \wedge \neg R)$
 $\equiv ((P \vee Q) \wedge \neg Q) \wedge ((\neg P \vee R) \wedge \neg R)$
 $\equiv (P \wedge \neg Q) \wedge (\neg P \wedge \neg R)$
 $\equiv \mathbf{F}$

More Examples

CNF

EXAMPLE: $(P \leftrightarrow Q) \wedge (Q \leftrightarrow R) \Rightarrow (P \leftrightarrow R)$

- $A = (P \leftrightarrow Q) \wedge (Q \leftrightarrow R); B = (P \leftrightarrow R).$
- $A = \mathbf{T}$ iff $(P \leftrightarrow Q) = \mathbf{T}$ and $(Q \leftrightarrow R) = \mathbf{T}$ iff $P = Q$ and $Q = R$
 - $A^{-1}(\mathbf{T}) = \{(\mathbf{T}, \mathbf{T}, \mathbf{T}), (\mathbf{F}, \mathbf{F}, \mathbf{F})\}$
- $B = \mathbf{T}$ iff $P = R$
 - $B^{-1}(\mathbf{T}) = \{(\mathbf{T}, \mathbf{T}, \mathbf{T}), (\mathbf{T}, \mathbf{F}, \mathbf{T}), (\mathbf{F}, \mathbf{T}, \mathbf{F}), (\mathbf{F}, \mathbf{F}, \mathbf{F})\}$
- $A^{-1}(\mathbf{T}) \subseteq B^{-1}(\mathbf{T})$

T, F

EXAMPLE: $(Q \rightarrow R) \Rightarrow ((P \vee Q) \rightarrow (P \vee R))$

- $A = Q \rightarrow R; B = ((P \vee Q) \rightarrow (P \vee R)).$
- $A = \mathbf{F}$ iff $(Q, R) = (\mathbf{T}, \mathbf{F})$
 - $A^{-1}(\mathbf{F}) = \{(\mathbf{T}, \mathbf{T}, \mathbf{F}), (\mathbf{F}, \mathbf{T}, \mathbf{F})\}$
- $B = \mathbf{F}$ iff $(P \vee Q, P \vee R) = (\mathbf{T}, \mathbf{F})$ iff $(P, Q) \neq (\mathbf{F}, \mathbf{F})$ and $(P, R) = (\mathbf{F}, \mathbf{F})$
 - $B^{-1}(\mathbf{F}) = \{(\mathbf{F}, \mathbf{T}, \mathbf{F})\}$
- $A^{-1}(\mathbf{F}) \supseteq B^{-1}(\mathbf{F})$

inverse!

More Examples

EXAMPLE: $(P \rightarrow R) \wedge (Q \rightarrow S) \wedge (P \vee Q) \Rightarrow R \vee S$

- $A = (P \rightarrow R) \wedge (Q \rightarrow S) \wedge (P \vee Q); B = R \vee S$
- $A \wedge \neg B \equiv (P \rightarrow R) \wedge (Q \rightarrow S) \wedge (P \vee Q) \wedge \neg(R \vee S)$
$$\begin{aligned} &\equiv (\neg P \vee R) \wedge (\neg Q \vee S) \wedge (P \vee Q) \wedge (\neg R \wedge \neg S) \\ &\equiv ((\neg P \vee R) \wedge \neg R) \wedge ((\neg Q \vee S) \wedge \neg S) \wedge (P \vee Q) \\ &\equiv ((\neg P \wedge \neg R) \vee (R \wedge \neg R)) \wedge ((\neg Q \wedge \neg S) \vee (S \wedge \neg S)) \wedge (P \vee Q) \\ &\equiv ((\neg P \wedge \neg R) \vee \mathbf{F}) \wedge ((\neg Q \wedge \neg S) \vee \mathbf{F}) \wedge (P \vee Q) \\ &\equiv (\neg P \wedge \neg R) \wedge (\neg Q \wedge \neg S) \wedge (P \vee Q) \\ &\equiv \neg R \wedge (\neg Q \wedge \neg S) \wedge (\neg P \wedge (P \vee Q)) \\ &\equiv \neg R \wedge (\neg Q \wedge \neg S) \wedge ((\neg P \wedge P) \vee (\neg P \wedge Q)) \\ &\equiv \neg R \wedge (\neg Q \wedge \neg S) \wedge (\mathbf{F} \vee (\neg P \wedge Q)) \\ &\equiv \neg R \wedge (\neg Q \wedge \neg S) \wedge (\neg P \wedge Q) \\ &\equiv \neg R \wedge \neg S \wedge \neg P \wedge (\neg Q \wedge Q) \\ &\equiv \neg R \wedge \neg S \wedge \neg P \wedge \mathbf{F} \\ &\equiv \mathbf{F} \end{aligned}$$