

Discrete Mathematics

Lecture 2

Liangfeng Zhang

School of Information Science and Technology

ShanghaiTech University

Summary of Lecture 1

Divide, Divisor, Multiple, Prime, Composite

Fundamental Theorem of Arithmetic: $n = p_1^{e_1} \cdots p_r^{e_r}$

The Well-Ordering Property: $\emptyset \neq S \subseteq \mathbb{N} \Rightarrow \min S \in S$

Division Algorithm: $a = bq + r; 0 \leq r < b$ for unique q, r

Ideal of \mathbb{Z} : A nonempty set $I \subseteq \mathbb{Z}$ such that

- $a, b \in I \Rightarrow a + b \in I; a \in I, r \in \mathbb{Z} \Rightarrow ra \in I$

THEOREM: I is an ideal of $\mathbb{Z} \Leftrightarrow I = d\mathbb{Z}$

Sum of Ideals: $I_1 + I_2 = \{x + y : x \in I_1, y \in I_2\}$

THEOREM: I_1, I_2 are ideals of $\mathbb{Z} \Rightarrow I_1 + I_2$ is an ideal of \mathbb{Z}

QUESTION: $a\mathbb{Z} + b\mathbb{Z} = ?$

Greatest Common Divisor

DEFINITION: Let $a, b \in \mathbb{Z}$ and at least one of them is nonzero.

- **common divisor:** an integer d such that $d|a, d|b$
- **greatest common divisor** $\gcd(a, b)$: the largest common divisor
 - **relatively prime:** $\gcd(a, b) = 1$

THEOREM: Let $a, b \in \mathbb{Z}$ and at least one of them is nonzero.

Then $a\mathbb{Z} + b\mathbb{Z} = \gcd(a, b)\mathbb{Z}$.

gcd theorem

已经决定 d 为 gcd 了.

- $\{a, b\} \neq \{0\} \Rightarrow a\mathbb{Z} + b\mathbb{Z} \neq \{0\}$
- There exists $d \in \mathbb{Z} \setminus \{0\}$ such that $a\mathbb{Z} + b\mathbb{Z} = d\mathbb{Z}$. W.l.o.g., $d > 0$.
 - d is a common divisor of a, b : $a \cdot 1 + b \cdot 0 \in d\mathbb{Z}$
 - d is greatest: Suppose that d' is a common divisor of a, b
 - $d'|a, d'|b$
 - $a\mathbb{Z} + b\mathbb{Z} = d\mathbb{Z} \Rightarrow d = as + bt$ for some integers s, t
 - $d'|d$ and thus $d' \leq d$

without loss of generality

proof divisor

proof greatest

THEOREM: There exist $s, t \in \mathbb{Z}$ such that $\gcd(a, b) = as + bt$.

FTA Proof

$c, a \in \mathbb{Z}$

THEOREM: If $a, b, c \in \mathbb{Z}, c|ab$ and $\gcd(c, a) = 1$, then $c|b$.

- There exist s, t such that $1 = \gcd(a, c) = as + ct$. 目标: $c|b$
- $b = \overbrace{bas}^{ba \cdot s} + bct$ 质技巧
- $c|ab, c|ct \Rightarrow c|(bas + bct) \Rightarrow c|b$

THEOREM: If p is a prime and $p|ab$, then $p|a$ or $p|b$.

- $p|a$: done
- $p \nmid a \Rightarrow \gcd(p, a) = 1$
 - $(\gcd(p, a) = 1) \wedge (p|ab) \Rightarrow p|b$ $n = p_1 \cdots p_r$

Fundamental Theorem of Arithmetic: proof of uniqueness

- Suppose that $n = p_1 \cdots p_r = q_1 \cdots q_s$, where p_i, q_j are all primes
 - $p_1|n \Rightarrow p_1|q_1 \cdots q_s \Rightarrow p_1|q_j$ for some $j \Rightarrow p_1 = q_j$
 - W.l.o.g., we suppose that $j = 1$. Then $p_2 \cdots p_r = q_2 \cdots q_s$
 - The theorem is true by induction.

$p_2 = 2 \cdots$

思想

FTA Applications

THEOREM: Suppose that $a = p_1^{\alpha_1} \cdots p_r^{\alpha_r}$, $b = p_1^{\beta_1} \cdots p_r^{\beta_r}$. Then

greatest: $d := p_1^{\min(\{\alpha_1, \beta_1\})} \cdots p_r^{\min(\{\alpha_r, \beta_r\})} = \gcd(a, b).$

- d is a common divisor of a, b
- d is largest among the common divisors
 - Suppose that d' is a common divisor of a, b
 - $d' = p_1^{e_1} \cdots p_r^{e_r}$ *tech*
 - $d' | a \Rightarrow e_i \leq \alpha_i$ for all $i \in [r]$; $d' | b \Rightarrow e_i \leq \beta_i$ for all $i \in [r]$
 - $e_i \leq \min\{\alpha_i, \beta_i\}$ for all $i \in [r]$

THEOREM: There are infinitely many primes.

- Suppose there are only n primes: p_1, \dots, p_n
- By FTA, $N = p_1 \cdots p_n + 1$ must be the product of primes
- $\exists i \in [n]$ such that $p_i | N$
- But $p_i \nmid N$

$$p_1 \cdots p_n \cdot p_i = p_1 \cdots p_n + 1$$

Handwritten notes: p_i is circled in green. $N = p_1 \cdots p_n + 1$ is written in red. $N \nmid p_i$ is written in green.

pl impossible

集合

Equivalence Relation

R是什么

运算表达 $A \times B = \{(a, b) \mid a \in A, b \in B\}$ = 二元关系

DEFINITION: Let A, B be two sets. A **binary relation** from A to B is a subset $R \subseteq A \times B$. // aRb means $(a, b) \in R$

EXAMPLE: $R = \{(a, a) \mid a \in \mathbb{Z}^+\}$ is a binary relation from \mathbb{Z}^+ to \mathbb{Z}^+
 • aRb means that $a = b$; R is " $=$ " (a, a) 的表达式决定这里是 "="

DEFINITION: Let A be a set. An **equivalence relation** R on A is a binary relation R from A to A such that

- Reflexive:** aRa for all $a \in A$ eg $(a, a) \in R$
- Symmetric:** $aRb \Rightarrow bRa$ for all $a, b \in A$ $(a, b) \in R, (b, a) \in R$
- Transitive:** $aRb, bRc \Rightarrow aRc$ for all $a, b, c \in A$ $(a, b) \in R, (b, c) \in R \Rightarrow (a, c) \in R$

DEFINITION: The **equivalence class** of $a \in A$ is the set

$$[a]_R = \{x \in A : xRa\}$$

↓ 可用于证明 eg relation

这是 eg relation 的特点, 不同于某个 R

$$[a]_R = \{a\}$$

不一定只是 " $=$ "

Congruence 同余

某个 n

THEOREM: Let $n \in \mathbb{Z}^+$. Then $R = \{(a, b) \in \mathbb{Z}^2 : n | (a - b)\}$ is an equivalence relation on \mathbb{Z} (from \mathbb{Z} to \mathbb{Z}).

- R is a binary relation from \mathbb{Z} to \mathbb{Z}
 - Reflexive: $n | (a - a) \Rightarrow aRa$
 - Symmetric: $aRb \Rightarrow n | (a - b) \Rightarrow n | (b - a) \Rightarrow bRa$
 - Transitive: $aRb, bRc \Rightarrow n | (a - b), n | (b - c) \Rightarrow n | (a - c) \Rightarrow aRc$

DEFINITION: Let $n \in \mathbb{Z}^+$ and $R = \{(a, b) \in \mathbb{Z}^2 : n | (a - b)\}$.

- The notation $a \equiv b \pmod{n}$ means that aRb .
 - $a \equiv b \pmod{n}$ is called a **congruence**
 - Read as: a is **congruent** to b (modulo n)
 - n is called the **modulus** of the congruence
 - $a \not\equiv b \pmod{n}$: $(a, b) \notin R$, or equivalently $n \nmid (a - b)$
 - Read as: a is not congruent to b modulo n

$\rightarrow R$ here: $a \equiv b \pmod{n}$

Topic
interested 在等价关系下找等价类

Congruence

$$a = kn + r$$

THEOREM: Let $n \in \mathbb{Z}^+$. For any $a \in \mathbb{Z}$, there is a unique integer r such that $0 \leq r < n$ and $a \equiv r \pmod{n}$.

- **Existence:** by division algorithm, $\exists q, r \in \mathbb{Z}$ s.t. $0 \leq r < n, a = qn + r$
 - $a \equiv r \pmod{n}$
- **Uniqueness:** suppose that $0 \leq r' < n$ and $a \equiv r' \pmod{n}$
 - $|r - r'| < n$ and $r \equiv r' \pmod{n}$
 - $|r - r'| < n$ and $n \mid (r - r')$
 - $r = r'$

DEFINITION: Let $a, n \in \mathbb{Z}$ and $n > 0$. Then there are unique integers q, r such that $0 \leq r < n$ and $a = nq + r$.

- We define $a \bmod n$ as r .

opt, 取余

Residue Class

残余

余集

DEFINITION: Let $\alpha \in \mathbb{R}$.

- $\lfloor \alpha \rfloor$: **floor** of α , the largest integer $\leq \alpha$
- $\lceil \alpha \rceil$: **ceiling** of α , the smallest integer $\geq \alpha$
 - If $a = bq + r$, then $q = \lfloor a/b \rfloor$ and $r = a - bq$

DEFINITION: Let $a \in \mathbb{Z}, n \in \mathbb{Z}^+$. We denote the equivalence class of a under the (equivalence relation mod n) with $[a]_n$ and call it the **residue class of a mod n** .

余数相等

$$[a]_n = a + n\mathbb{Z} = \{a + nx : x \in \mathbb{Z}\}$$

- any element of $[a]_n$ is a **representative** of $[a]_n$, since have same charac

EXAMPLE: $[0]_6 = \{0, \pm 6, \pm 12, \dots\}$; $[1]_6 = \{\dots, -11, -5, 1, 7, 13, \dots\}$; ...

↑
 $6q+1$

Residue Class

THEOREM: Let $n \in \mathbb{Z}^+$, $a, b \in \mathbb{Z}$. Then - 同全同

$$[a]_n \cap [b]_n = \emptyset \text{ or } [a]_n = [b]_n.$$

- $[a]_n \cap [b]_n = \emptyset$: done
- $[a]_n \cap [b]_n \neq \emptyset$
 - $\exists c \in [a]_n \cap [b]_n$
 - $c \equiv a \pmod{n}, c \equiv b \pmod{n}$
 - $a \equiv b \pmod{n}$ transitive
 - $\exists t \in \mathbb{Z}$ such that $a = b + nt$ \hookrightarrow mod 显式化
 - $[a]_n = \{a + nx : x \in \mathbb{Z}\} = \{b + nt + nx : x \in \mathbb{Z}\} = [b]_n$

推论

COROLLARY: $[a]_n = [b]_n$ iff $a \equiv b \pmod{n}$.

COROLLARY: $\{[0]_n, [1]_n, \dots, [n-1]_n\}$ is a partition of \mathbb{Z} .

- $[a]_n \cap [b]_n = \emptyset$ for all $a, b \in \{0, 1, \dots, n-1\}$
- $\mathbb{Z} = [0]_n \cup [1]_n \cup \dots \cup [n-1]_n$

$$\mathbb{Z}_n$$

DEFINITION: Let n be any positive integer. We define \mathbb{Z}_n to be set of all residue classes modulo n .

- $\mathbb{Z}_n = \{[0]_n, [1]_n, \dots, [n-1]_n\}$

- $\mathbb{Z}_n = \{0, 1, \dots, n-1\};$

- $\mathbb{Z}_n = \{[1]_n, [2]_n, \dots, [n]_n\}$

- $\mathbb{Z}_n = \{1, 2, \dots, n\}$

$\rightarrow = [0]_n$

cover all residue possibility

$(0, 1, \dots, n-1)$

EXAMPLE: Two representations of the set \mathbb{Z}_6

- $\mathbb{Z}_6 = \{[0]_6, [1]_6, [2]_6, [3]_6, [4]_6, [5]_6\}$

$= \{0, 1, 2, 3, 4, 5\}$

- $\mathbb{Z}_6 = \{[-3]_6, [-2]_6, [-1]_6, [0]_6, [1]_6, [2]_6\}$

$= \{-3, -2, -1, 0, 1, 2\}$

> Q: diff?

一种简写法

$$\mathbb{Z}_n$$

DEFINITION: Let $n \in \mathbb{Z}^+$. For all $[a]_n, [b]_n \in \mathbb{Z}_n$, define

- **addition:** $[a]_n + [b]_n = [a + b]_n$ *residue class*
- **subtraction:** $[a]_n - [b]_n = [a - b]_n$ *operation*
- **multiplication:** $[a]_n \cdot [b]_n = [a \cdot b]_n$

Well-defined? If $a \equiv a' \pmod{n}$ and $b \equiv b' \pmod{n}$, then $a \pm b \equiv a' \pm b' \pmod{n}$ and $ab \equiv a'b' \pmod{n}$.

- Hence, $[a]_n \pm [b]_n = [a']_n \pm [b']_n$; $[a]_n \cdot [b]_n = [a']_n \cdot [b']_n$
 - $a \equiv a' \pmod{n} \Rightarrow n | (a - a') \Rightarrow \exists x$ such that $a - a' = nx$
 - $b \equiv b' \pmod{n} \Rightarrow n | (b - b') \Rightarrow \exists y$ such that $b - b' = ny$
 - $(a + b) - (a' + b') = nx + ny$ *(mod n) 相关证明:*
 - $(a - b) - (a' - b') = nx - ny$ *结果 n 因子化*
 - $ab - a'b' = a(b - b') + b'(a - a') = any + b'n x$

$$\mathbb{Z}_n^*$$

倒数集

DEFINITION: Let $n \in \mathbb{Z}^+$ and $[a]_n \in \mathbb{Z}_n$. $[s]_n \in \mathbb{Z}_n$ is called an **inverse** of $[a]_n$ if $[a]_n[s]_n = [1]_n$.

- **division:** If $[a]_n[s]_n = [1]_n$, define $\frac{[b]_n}{[a]_n} = [b]_n \cdot [s]_n$ $\exists s.t. as+nt=1$

THEOREM: Let $n \in \mathbb{Z}^+$. $[a]_n \in \mathbb{Z}_n$ has an inverse iff $\gcd(a, n) = 1$.

- Only if: $\exists s$ s.t. $[a]_n[s]_n \equiv [1]_n$; $\exists t, as - 1 = nt$; $\gcd(a, n) = 1$
- If: $\exists s, t$ s.t. $as + nt = 1$; $as \equiv 1 \pmod{n}$ $as=nt+1$

DEFINITION: Let $n \in \mathbb{Z}^+$. Define $\mathbb{Z}_n^* = \{[a]_n \in \mathbb{Z}_n : \gcd(a, n) = 1\}$

- If n is prime, then $\mathbb{Z}_n^* = \{1, 2, \dots, n-1\}$
- If n is composite, then $\mathbb{Z}_n^* \subset \mathbb{Z}_n$

EXAMPLE: $\mathbb{Z}_5^* = \{1, 2, 3, 4\}$; $\mathbb{Z}_6^* = \{1, 5\}$; $\mathbb{Z}_8^* = \{1, 3, 5, 7\}$

2, 3, 4
gcd $\neq 1$

Euler's Phi Function

QUESTION: How many elements are there in \mathbb{Z}_n^* ?

- $|\mathbb{Z}_n^*|$ is the number of integers $a \in [n]$ such that $\gcd(a, n) = 1$

DEFINITION: (Euler's Phi Function) $\phi(n) = |\mathbb{Z}_n^*|, \forall n \in \mathbb{Z}^+$

- $\phi(n)$ is the number of integers $a \in [n]$ such that $\gcd(a, n) = 1$

THEOREM: Let p be a prime. Then $\forall e \in \mathbb{Z}^+, \phi(p^e) = p^{e-1}(p-1)$.

- Let $x \in [p^e]$.
- $\gcd(x, p^e) \neq 1$ iff $p|x$
 iff $x = p, 2p, \dots, p^{e-1} \cdot p$
- $\phi(p^e) = p^e - p^{e-1} = p^{e-1}(p-1)$

EXAMPLE: $\phi(3^2) = 3(3-1) = 6$

- $\mathbb{Z}_9^* = \{1, 2, 3, 4, 5, 6, 7, 8, 9\}$

EXAMPLE: $\phi(p) = p-1$

- $\mathbb{Z}_p^* = \{1, 2, \dots, p-1\}$

Euler's Phi Function

$$p_1^{e_1-1}(p_1-1) p_2^{e_2-1}(p_2-1) \dots \rightarrow p_1^{e_1-1} \dots p_k^{e_k-1} \left(1 - \frac{1}{p_1}\right) \dots \left(1 - \frac{1}{p_k}\right)$$

QUESTION: Formula of $\phi(n)$ for general integer n ?

THEOREM: If $n = p_1^{e_1} \dots p_k^{e_k}$ for distinct primes p_1, \dots, p_k and integers $e_1, \dots, e_k \geq 1$, then $\phi(n) = \phi(p_1^{e_1}) \dots \phi(p_k^{e_k})$.

Hence, $\phi(n) = n(1 - p_1^{-1}) \dots (1 - p_k^{-1})$.

- There are many proofs. We will see in the future.

COROLLARY: If $n = pq$ for two different primes p and q , then $\phi(n) = (p - 1)(q - 1)$.

EXAMPLE: $\phi(10) = (2 - 1)(5 - 1) = 4$; $n = 10$; $p = 2, q = 5$

- $\mathbb{Z}_{10}^* = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$

Euler's Theorem

THEOREM (Euler) Let $n \geq 1$ and $\alpha \in \mathbb{Z}_n^*$. Then $\alpha^{\phi(n)} = 1$.

- $\alpha^{\phi(n)}, 1$ are both residue classes modulo n
- Suppose that $\alpha = [a]_n$ for $a \in \mathbb{Z}$. Then $\alpha^{\phi(n)} = 1$ is $([a]_n)^{\phi(n)} = [1]_n$
- How to prove?
 - Consider the map $f: \mathbb{Z}_n^* \rightarrow \mathbb{Z}_n^* \quad [x]_n \mapsto [a]_n \cdot [x]_n$
 - We show that f is injective 单射
 - $f([x]_n) = f([y]_n)$
 - $[a]_n \cdot [x]_n = [a]_n \cdot [y]_n$
 - $[ax]_n = [ay]_n$
 - $n | a(x - y)$
 - $n | (x - y)$, this is because $\gcd(n, a) = 1$
 - $[x]_n = [y]_n$

Euler's Theorem

THEOREM (Euler) Let $n \geq 1$ and $\alpha \in \mathbb{Z}_n^*$. Then $\alpha^{\phi(n)} = 1$.

- $\alpha^{\phi(n)}, 1$ are both residue classes modulo n
- Suppose that $\alpha = [a]_n$ for $a \in \mathbb{Z}$. Then $\alpha^{\phi(n)} = 1$ is $([a]_n)^{\phi(n)} = [1]_n$
- How to prove?
 - Consider the map $f: \mathbb{Z}_n^* \rightarrow \mathbb{Z}_n^* \quad [x]_n \mapsto [a]_n \cdot [x]_n$
 - Suppose that $\mathbb{Z}_n^* = \{[x_1]_n, \dots, [x_{\phi(n)}]_n\}$.
 - $f([x_1]_n) \cdots f([x_{\phi(n)}]_n) = [x_1]_n \cdots [x_{\phi(n)}]_n$
 - $[ax_1]_n \cdots [ax_{\phi(n)}]_n = [x_1]_n \cdots [x_{\phi(n)}]_n$
 - $[a^{\phi(n)} x_1 \cdots x_{\phi(n)}]_n = [x_1 \cdots x_{\phi(n)}]_n$
 - $n \mid (a^{\phi(n)} - 1)x_1 \cdots x_{\phi(n)}$
 - $n \mid (a^{\phi(n)} - 1)$, this is because $\gcd(n, x_1 \cdots x_{\phi(n)}) = 1$
 - $[a^{\phi(n)}]_n = [1]_n$, i. e., $([a]_n)^{\phi(n)} = [1]_n$

Fermat's Little Theorem

EXAMPLE: Understand Euler's theorem with $\mathbb{Z}_{10}^* = \{1, 3, 7, 9\}$.

- $n = 10, \phi(n) = 4,$
- $1^4 \equiv 1 \pmod{10} \Rightarrow ([1]_{10})^4 = [1]_{10}$
- $3^4 = 81 \equiv 1 \pmod{10} \Rightarrow ([3]_{10})^4 = [1]_{10}$
- $7^4 = 2401 \equiv 1 \pmod{10} \Rightarrow ([7]_{10})^4 = [1]_{10}$
- $9^4 = 6561 \equiv 1 \pmod{10} \Rightarrow ([9]_{10})^4 = [1]_{10}$
 - Consider the map $f: \mathbb{Z}_{10}^* \rightarrow \mathbb{Z}_{10}^* \quad [x]_n \mapsto [9]_n \cdot [x]_n$
 - $f([1]_{10}) = [9]_{10} \cdot [1]_{10} = [9]_{10}; f([3]_{10}) = [7]_{10}; f([7]_{10}) = [3]_{10}; f([9]_{10}) = [1]_{10}$
 - f is injective
 - $f([1]_{10})f([3]_{10})f([7]_{10})f([9]_{10}) = [9]_{10}[7]_{10}[3]_{10}[1]_{10}$

Fermat's Little Theorem: If p is a prime and $\alpha \in \mathbb{Z}_p$.

Then $\alpha^p = \alpha$.

- This is a corollary of Euler's theorem for $n = p$
- By Euler's theorem, $\alpha^{p-1} = 1$
 - $\alpha^p = \alpha$

$p \geq 1 \quad \phi(p) = p-1 \quad \alpha \in \mathbb{Z}_p$

$\alpha \in \mathbb{Z}_p^*$

since p prime

$\alpha^{p-1} = 1 \rightarrow \alpha^p = \alpha$