Exercise 02 C

Cristian Manuel Abrante Dorta

March 2020

We consider an square image I whose size is $\mathcal{N}x\mathcal{N}$, and three types of structuring elements.

First type is structuring element \mathcal{B} , which is a 2-dimensional structuring element of size $\mathcal{M}x\mathcal{M}$.

$$\mathcal{B} = \begin{pmatrix} b_{1,1} & b_{1,2} & \cdots & b_{1,m} \\ b_{2,1} & b_{2,2} & \cdots & b_{2,m} \\ \vdots & \vdots & \ddots & \vdots \\ b_{m,1} & b_{m,2} & \cdots & b_{m,m} \end{pmatrix}$$
(1)

Second type is \mathcal{C} which is a 1 dimensional horizontal structuring element:

$$C = \begin{pmatrix} c_1 & b_2 & \cdots & c_m \end{pmatrix} \tag{2}$$

Third type is a \mathcal{D} wich is a 1 dimensional vertical structuring element:

$$\mathcal{D} = \begin{pmatrix} d_1 \\ d_2 \\ \vdots \\ c_m \end{pmatrix} \tag{3}$$

We can assume those two properties of the elements:

$$\mathcal{B} = \delta_{\mathcal{C}}(\mathcal{D}) = \delta_{\mathcal{D}}(\mathcal{C}) \tag{4}$$

Calculate the number of max operations that must be computed in order to process a $\mathcal{N}x\mathcal{N}$ input image using the following alternatives:

- First alternative: $\delta_{\mathcal{B}}(I)$
- Second alternative: $\delta_{\mathcal{C}}\delta_{\mathcal{D}}(I)$

1 Considerations

We are going to compute the number of elementary max operations that we have to do in order to compute the max of a list of elements.

Considering only two elements:

$$max(x_1, x_2) \to 1 \ operation$$
 (5)

Considering three elements:

$$max(x_1, x_2, x_3) = max(x_1, max(x_2, x_3)) \rightarrow 2 \text{ operations}$$
 (6)

Considering four elements:

$$max(x_1, x_2, x_3, x_4) = max(x_1, max(x_2, max(x_3, x_4))) \rightarrow 3 \text{ operations}$$
 (7)

Finally if we consider n elements:

$$max(x_1, x_2, \cdots, x_n) = max(x_1, max(x_2, \cdots, max(x_{n-1}, x_n))) \rightarrow n-1 \text{ operations}$$
(8)

2 First alternative: $\delta_{\mathcal{B}}(I)$

For computing the operations of the structuring element, whose size is $\mathcal{M}x\mathcal{M}$, we have M^2 elements. Following the reasoning of previous equation:

$$\mathcal{M}x\mathcal{M} \to \mathcal{M}^2 - 1 \ operations$$
 (9)

As the image have $\mathcal{N}x\mathcal{N}$ elements, or \mathcal{N}^2 , we can compute the total number of operations $(NumOp_1)$:

$$NumOp_1 = \mathcal{N}^2(\mathcal{M}^2 - 1) \tag{10}$$

3 Second alternative: $\delta_{\mathcal{C}}\delta_{\mathcal{D}}(I)$

For each structuring element as their size is \mathcal{M} , we can say that the number of elementary operations is M-1, following the preovious reasoning.

As the image have $\mathcal{N}x\mathcal{N}$ elements, or \mathcal{N}^2 , we can compute the total number of operations, with only one structuring element:

$$\mathcal{N}^2(\mathcal{M} - 1) \tag{11}$$

But, we have two elements, so we have to multiply the number of operations by two:

$$NumOp_2 = 2\mathcal{N}^2(\mathcal{M} - 1) \tag{12}$$

4 Conclusion

The number of operations for each alternative is:

$$NumOp_1 = \mathcal{N}^2(\mathcal{M}^2 - 1)$$

$$NumOp_2 = 2\mathcal{N}^2(\mathcal{M} - 1)$$
(13)

We can confirm that the number of operations in the second alternative is lower than the first one.