Ontology of plant genealogy

What does this set of words mean?



OG – ontology for plant genealogy

- It IS a description of the breeding history of 'germplasm'
- It IS NOT a description of 'germplasm' phenotype
- It IS NOT a description of 'germplasm' genotype



OG – a definition


A formalised logical description of chapter 2 "THE GENEALOGY MANAGEMENT SYSTEM" of the ICIS manual in such a way that both biologists (plant breeders, plant geneticists) and database developers (bio-informaticists and programmers) can understand each other.

		MGRP MCODE M			maintenance m			MREF MPF	RGN MFPRG	MATT	R GENEQ	MUID	LMID MD	ATE	
					Methods for storing l	istorical pe	digrees with incomplete informat								
1	GEN	S UGM U	NKNOWN GENERA	TIVE	Unknown generative meth	od for storin	g historic pedigrees for self	0	0	0	0	1	2 19980	0610	
	52.1	Annex 2.2M	F™Cultivars, l	Popula	ations, Breeding	strategi	ies, Generative, Deriv	ative an	d Maint	nanc	e Metho	ds for]
2	GEN	fertilised, Q			Viegetatively Re	eroduc	edietro pes pecies: ross	0	0	0	0	1	0 19980	0610	
3	GEN	Breeding System	Cultivars and Popu	ılations	Breeding Strategies		Generative Methods	Derivative	methods		Maintenanc	e methods	s		
3	GEN	Out Breeding	Out Breeding Cultivars		Half Mass Selection		Single cross cf		Induced mutation pop		Seed increase open pollination				1
4	GEN	Cross breeding Cross fertilising	Open Pollinated Variety Synthetic CV Tah		Full Mass Selection Recurrent Half-sib S	Selection	Three-way cross cf Double cross cf	Double haploid pop Rouging cf		Seed increase full diallel bulked Seed increase half diallel bulked					
-	GEN	Lious fertinising			D	chection Choutline Fof a general model f		C:11			C-1:				
5	GEN		Hybrid CV Obsolete CV		Selfed Families		Full diallel gross bulked	Restorer se	lection				cross		
6	GEN	Ħ	Populations	The b	pases tor the classification of the seed, which can be	contained e a single	in this model is that the source each prosend bulked from a sing	pirthe temal le plant a se	gleametes is lected set of	determi	ned by the ragandom	nation of	·f		
		Ц		set of p	plants.Reprover rafths	relleneneed	s parbardetemminedsbut can be fi	iomathrasan	seplantananotl	er plan	Breekestest	ed cf	.1		
7	GEN		Landrace population Landrace cultivar	on set	t of plantseononrandom se Reciprocal Recurrer					depende	Pouthetation Certified se				
8	GEN	\forall	Collection populat		Stelatri Stoarceng Se		PopulatativBadlethoods:f	Landrace c	ult iveris fative						
		Ш	Collection wild sp population	P e Source	Families Population Back Cr	oss	Population Backcross recessive cf	Collection Collection	population of wild spp		Import Seed	i			
9	GEN		Collection weedy s	Pingle	Same Single Plant Plus all the methods	40(G) Se	filingerspecific cross of Selected pollen cross of		S05(O) Sing	le Seed		1			
31	DER	H	population Genetic Stock	Plant	below for selection			Labyleten Calleten	Mant Mentin	ation	Selection	 	OF 50		
			Mechanical Mixtu	re	fixed lines.	Ta	Narrow based tester, pop cf	iplasm	5100 QC	nati	om sto	red ii	n GMS.		
32	DER	Ħ			Different Single Blant		Broad based tester, pop er	210(S)	Maintainer !	Selection	n				or any germplasm recorded. Th
33	DER	H			Different Single Plant Modified Bulk Sele	1692(S),40	Note the second of the second								f seed that exist or did exist yo
دد	DEK				Bulk Selection Single Seed Descen	103(I),40	3 (Rahlkoubhra Gress) ul Opien lpoblinossion of		e GEN a				a me pack	zeis 0	seed that exist of the exist yo
		7			Double Haploid	405(Q) ₁ F	unopenipornomon ci ultanallation esperaliked aktomaticani sees cf						rmnlaar	and a	etails are stored in the method
40	CEN	귀			Back Cross Re-selection		alfandingen seess of alfantial of usolis belikusts of	table (une meti	2 P/	The me	s or ge	ndigatos y	anu (whath	er the germplasm was
40	GEN				Ke-selection	408(O) P	ar Randoan Halfosi b Fam								
	<u> </u>	닉				409(O) P	arsishedish Helfosib Fanked								ive process, and then how the
						424(O) C 425(O) P	onvergent Cross Natural mutation of Induced mutation population	germpi	asiii was	proc	ucpable	zezalisk Posta	ceproduc ti v	WISYSt	which change with each
60	MAN					107(S) B	Induced mutation population	applica	iiion are	store		linated	miormatic	Inbree	eins in Crop Hains in State Letton 2.2.4) eing, self pollinated or self fertilised s produce Rhy, Baryerative
61	MAN	H				109(S) In	Office State Control of the Contro	identif	ication c	oae i	or last ir	istance	e or germj	Plasn	ABIOMILLE ARE BEGIETATIVE
		Ш			0.1 . 10.11	416(O) N	Allo-polyploid of arrow based tester, line CF								ved Cotto De geohpea, Sorghum
62	MAN				Selected Bulk	1104(S) F	made Complex Top Cross				I the ot	reent 11 Pollinate	nstance is	Outbre	uced by a generative process.
	l	1				416(8) N	arted gollep Closs SF		ication c		or the in	media	ate source	Z ora Examp	rling cross pollinated on cross fertilistes: Maize, Pearl Millet, Cucurbits
							road based tester, Line CF		lasm. (N		14 45H P	domnati	HA SCIIII N	Sellin	compatible species
					Bulk	104(S) F	arrow Based Tester, POP CF emale Complex Top Cross —			ode o					neon the generis of the paya
							andom Pollen Cross SF	germpl			Vegeta				ntively or clonally propagated species.
							andom Half Sib Families elected Half Sib Families	Identif	ier of bre	eding	g of off	erryoc:			oles: Potatoes, Casava, Yam, Taro, Supple, Strawberries
				Select ed	Single Plant		Tale Complex Top Cross	Data	f generat	ion	Anomi	ctic5			
				ea Bulk		416(O) N	larrow based tester line CF		•	-		1 *	reservatio	Examp	tlic (produce seed by asexual means) s bles: Green Panic, Buffel grass
					Same Selected Bulk	420(Q) P	REFER	Identat	Purification	liogr	aphic re	ed for it	n the ICIS N	lethod here	s Table the germplasm data was g methods usually follow those for Se
						421 (0) R EN	CE	re204(S),	Selected Bul	gi ng 5¹ k		iply cate	eřed for as b	oreedin	g methods úsuálly follow those for Se
									Full Mass S		crops.	thly cate	ered for by r	method	ds suitable for Open Pollinated crops.
					Different Selected Bulk		omplex Cross pen Pollenated SF								most maintenance methods catered for
						404(O) F	ull diallel cross								sed in their mode of sexual reproduction
							ull diallel cross bulked Ialf diallel cross				⁵ Proba	ably cate	ered for by r	method	ls suitable for Vegetatively Propagated
						407(O) H	tati dianel cross								

Н

Ontology

- First bit
 - □ A taxonomy of concepts
 - hierarchical classification of definitions
- Second bit?
 - Defining relationships
 - Exclusive
 - Is a parent of
 - Is an off-spring of
 - **?**
- A program that converts a human version (english?) to a logical framework





Germplasm is

- Propagules which are
 - □ double haploid
 - genetically modified
 - □ predominately cross fertilise Derivative Name
 - □ predominately self fertilised on MaintenanceName
 - vegetatively reproduced

Asserted Hierarchy





- owl:Thing
- Germplasm
 - DoubleHaploid
 - C) GMO
 - C PredominatelyCrossFertilised
 - C) PredominatelySelfFertilised
 - VegetativelyReproduced
- Germplasmldentifer
 - - GenerativeName
- - 🕒 GermplasmMethod
 - Derivative
 - Generative
 - Maintenance



Germplasm is named serted Hierarchy

Name of propagules






- 🔻 🕒 Germplasm
 - C DoubleHaploid
 - C GMO
- □ derived from a derivative method redominately Cross Fertilised
- derived from a generative method described des
- L C Germplasmidentife
- □ derived from a management methodideName
 - MaintenanceName
 - GermplasmMethod
 - 🕨 🥲 Derivative
 - Generative
 - 🕨 🔘 Maintenance



Germplasm is created


- Propagules produced
 - □ to decrease genetic variation (ננים selection)
 - □ to increase genetic variation (using the limit of the limit of
 - □ to maintain genetic variation (USUAlly Seed increase)





Germplasm is

- Propagules which are
 - □ Double haploid
 - □ Genetically modified



Germplasm is predominantly cross fertilised

Propagules which are

predominately cross fertilised

and are randomly mated

□ in professional breeding

□ as a traditional variety not produced by professional breeding C Random C Selected

For Project: genealogy_07-01-05_a

PredominatelyCrossFertilised

C ReciproculRecurrentSelectionCycle

C RecurrentSelectionCycle

C OpenPollenatedVariety

C DoubleHaploid

▼ (C) RandomlyMated C Breeding

C Wild

C Hybrid

C Landrace

Asserted Hierarchy

C GMO

owl: Thing C Germplasm


□ in a reciprocal recurrent selection program C VegetativelyReproduced

in a recurrent selection program

in the wild (not maintained by farmers)




- Propagules which are
 - predominantly cross fertilised
 - and are reproduced by selective mating
 - □ of a few selected (by progeny testing) fixed lines...
 - ☐ of selected (by phenotype) individuals (mass selection)
 - □ of many selected (by progeny testing) fixed
 - and are reproduced by selfing
 - of randomly chosen individuals
 - □ of selected individuals



Germplasm is predominately wizards code self fertilised

- Propagules which are
 - predominately self fertilised
 - derived from a single plant from an advanced inbred population
 - and used for breeding
 - and released as a cultivar
 - □ and derived from a traditional variety hot professionally bred
 - and collected off farm
 - are from an early inbred generation
 - vegetatively reproduced
 - from a single plant
 - from tissue culture derived from a single plant clone





Germplasm is created (derivative)

- Propagules produced
 - □ to decrease genetic variation (usually selection)
 - by collection (a landrace or wild population which may be an individual or a population)
 - by selection of
 - and bulking
 - selected plants
 - unselected plants
 - □ single plants
 - mutants
 - by doubling haploids

Н

Germplasm is created (generative)

- Propagules produced
 - □ to increase genetic variation (usually a cross)
 - by producing haploids from heterozygotes
 - by mutation
 - caused by a mutagenic agent
 - recognized as a natural variant
 - by tissue culture methods
 - by crossing
 - individual plants
 - populations
 - individuals to populations
 - by genetically transforming a plant (the result of a conversion event)



Germplasm is created (management)

- Propagules produced
 - □ to maintain genetic variation (usually seed increase)
 - By forming a cultivar for release
 - for exporting to another organisation or project
 - by importation from elsewhere (the process of recording the accession)
 - by extraction of a fixed line from an advanced inbred population
 - by creating a population (usually by random mating)
 - by maintaining an existing population (usually by random mating)
 - by seed increase
 - by seed storage



Germplasm is named (derivative)

- Name of propagules
 - ☐ from a derivative method
 - is a breeder (or breeding program) name
 - is a selection history
 - □ of a (modified) pedigree program
 - □ of a recurrent selection program

М

Germplasm is named (generative)

- Name of propagules
 - ☐ from a generative method
 - due to genetic engineering event
 - due to a cross
 - □ is a sequential name (usually numeric) in order of crossing (overall, in a year or a cycle)
 - □ is a name
 - □ is a string representing parentage
 - by a non Purdy string
 - by a Purdy string
 - due to haploid doubling
 - due to mutation

Н

Germplasm is named (management)

- Name of propagules
 - □ is an accession number
 - externally given
 - locally given
 - □ is a name for a commercial variety
 - Which is heterozygous
 - propogated asexually
 - is a homogenous collection of a single heterozygous genotype
 - is a traditional clonal variety
 - propagated by crossing
 - of a small number of inbred lines
 - by farmer selection
 - by mass selection
 - Many inbred lines
 - Which is homozygous
 - derived from an a single plant of a highly inbred population
 - is a traditional variety
 - □ is a deliberate mixture of fixed lines

- □ is the name of a (collection of) homozygous individual(s)
 - from a breeding population
 - collected
 - from a bulk of many plants
 - is a weed
 - not a weed
 - □ from a single plant
 - is a weed
 - not a weed
- □ is the name of a population (collection of heterogenous/heterozygous individuals)
 - used for breeding
 - is a recurrent selection cycle name
 - is a recognised genetic stock
 - is a deliberate mixture of populations
 - named after place of collection
 - collected off farm