The experience of developing an Earth System Model in Brazil

Vinicius Capistrano, Paulo Nobre, Silvio Figueroa, Fernando Pesquero, Manoel Baptista

Center for Weather Forecast and Climate Studies (CPTEC)
National Institute for Space Research (INPE)
Cachoeira Paulista-SP, Brazil

Introduction

The Brazilian Earth System Model, coupled ocean-atmosphere (BESM-OA2.5) is the coupling of the CPTEC/INPE Brazilian Atmospheric Model (BAM) and the GFDL/NOAA Modular Ocean Model (MOM4p1 and MOM5) (Nobre et al., 2013; Capistrano et al., 2016).

BESM development strategy

- 1) full use of CPTEC's experience and sub-models
- 2 collaboration with advanced climate change centers abroad
 - Use GFDL/FMS coupler to add components:

Dynamic vegetation with carbon cycle;


Continental hydrology-ocean coupling;

Ocean carbon cycle;


Enhanced sea ice and pack ice;


Atmospheric chemistry.


BESM component models





Results














Pesquero et al. (2016).

Acknowledgements

This work is a contribution to the Brazilian Earth System Model (BESM) funded by CAPES/ANA (No. 88887.115872/2015-01), Rede-Clima (550022/2014-7) and CNPq/IBAS/MOST (No. 490237/2011-8).

ESGF@INPE/CPTEC

Home About Us Contact Us



Reference

- [1] Nobre, P. et~al.. Climate simulation and change in the Brazilian climate model. J~Clim, vol. 26(17):6716-6732,~2013.
- [2] Capistrano, V. B. et al.. Climate Sensitivity BESM version 2.5. To be submitted to Journal of Climate, 2016.
- [3] Pesquero, F. et al.. The equilibrium response of the Brazilian Climate Model BESM2.5 to 4xCO2 atmospheric concentration forcing. Submitted to Theoretical and Applied Climatology, 2016.















