A Formal Proof of the Conservation of the Laplace–Runge–Lenz Vector, Kepler's First Law and the Hidden $\mathfrak{so}(4)$ Symmetry in Lean 4

Nichlas Löfdahl Claude (Anthropic)*

October 2026

Abstract

We present a machine-checked formalization, in the Lean 4 proof assistant on top of the Physlib library, of the classical Kepler problem: a particle in the inverse-square potential $V(r) = -k/r$. We define the Laplace–Runge–Lenz (LRL) vector on phase space and prove that its Poisson bracket with the Hamiltonian vanishes away from the centre of force. We prove the algebraic identities $\langle \mathbf{A}, \mathbf{L} \rangle = 0$, $\langle \mathbf{A}, \mathbf{q} \rangle = \|\mathbf{L}\|^2 - mk\|\mathbf{q}\|$ and $\|\mathbf{A}\|^2 = (mk)^2 + 2mH\|\mathbf{L}\|^2$, and derive that bound orbits ($H < 0$) have eccentricity less than one. Finally, we lift these phase-space statements to trajectories: along every differentiable solution of Hamilton's equations avoiding the origin, the LRL vector and $\|\mathbf{L}\|^2$ are constant, so the position satisfies the polar equation of a conic with a focus at the centre of force (Kepler's first law). Finally, we formalize the hidden symmetry of the Kepler problem: the Poisson algebra $\{L_a, L_b\} = L_{a\times b}$, $\{A_a, L_b\} = A_{a\times b}$, $\{A_a, A_b\} = -2mH\, L_{a\times b}$, the decomposition of the algebra generated by $\mathbf{L}$ and $\mathbf{M} = \mathbf{A}/\sqrt{-2mH}$ on the bound region into two commuting copies of $\mathfrak{so}(3)$, and the Casimir relation $\|\mathbf{J}_\pm\|^2 = -mk^2/(8H)$, a classical counterpart of Pauli's algebraic derivation of the hydrogen spectrum. The development is about 860 lines and uses no axioms beyond Lean's standard three.

1 Introduction

Formal verification of physics has gained momentum with the growth of Lean's mathematical library Mathlib [6] and the physics library Physlib (formerly HepLean/PhysLean) [7]. Physlib now contains, among much else, a quantum-mechanical treatment of the hydrogen atom including commutation relations of the quantum LRL operator, and, as of October 2026, basic infrastructure for classical central-force problems: the Hamiltonian $H(\mathbf{p}, \mathbf{q}) = \langle \mathbf{p}, \mathbf{p} \rangle / 2m + V(\|\mathbf{q}\|)$, its gradients, and the conservation of angular momentum in Poisson-bracket form.

What was missing is the defining classical result of the Kepler problem: the existence of the additional conserved vector $\mathbf{A}$ that explains why Kepler orbits close, and the derivation of the conic orbit equation from it. Before starting we searched Physlib's sources, its open pull requests and issues, and the web; we are not aware of an existing formalization of the classical LRL vector or of the conic orbit theorem in Lean. This note fills that gap.

*AI system (model Claude Opus 5.5), developed by Anthropic. Claude wrote the Lean proofs and drafted this manuscript in collaboration with the first author.

Contributions.

- A definition of the Kepler problem as an instance of Physlib's `CentralForce` structure, and of the LRL vector as a function on phase space (§3.1).
- Algebraic identities: the triple-product expansion of $\mathbf{A}$, orthogonality to $\mathbf{L}$, the orbit equation, the norm identity, and the eccentricity–energy criterion (§3.2).
- Conservation: explicit phase-space gradients of each component of $\mathbf{A}$ and the vanishing of $\{A_i, H\}$ (§3.3).
- Trajectory results: a general lemma that a Poisson-commuting observable is constant along solutions of Hamilton's equations, and Kepler's first law (§3.4).
- The hidden symmetry: the Poisson algebra of $\mathbf{L}$, $\mathbf{A}$ and H, the isomorphism $\mathfrak{so}(4) \cong \mathfrak{so}(3) \oplus \mathfrak{so}(3)$ on the bound region, and its Casimirs (§4).

The source is available in the `kepler-lrl` folder of `https://github.com/Crotalus/ai-formalized`; it builds against Physlib commit `3bfc7af` with Lean `v4.35.0`.

2 Mathematical background

Consider a particle of mass $m > 0$ at position $\mathbf{q} \in \mathbb{R}^3$ with momentum $\mathbf{p}$ in the potential $V(r) = -k/r$, with Hamiltonian

$$H(\mathbf{p}, \mathbf{q}) = \frac{\langle \mathbf{p}, \mathbf{p} \rangle}{2m} - \frac{k}{\|\mathbf{q}\|}.$$

The angular momentum is $\mathbf{L} = \mathbf{q} \times \mathbf{p}$ and the Laplace–Runge–Lenz vector is

$$\mathbf{A} = \mathbf{p} \times \mathbf{L} - \frac{mk}{\|\mathbf{q}\|}\,\mathbf{q} = \langle \mathbf{p}, \mathbf{p} \rangle\, \mathbf{q} - \langle \mathbf{q}, \mathbf{p} \rangle\, \mathbf{p} - \frac{mk}{\|\mathbf{q}\|}\,\mathbf{q},$$

where the second form uses $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = \langle \mathbf{a}, \mathbf{c} \rangle \mathbf{b} - \langle \mathbf{a}, \mathbf{b} \rangle \mathbf{c}$ [2, §3.9]. Taking the inner product with $\mathbf{q}$ and using Lagrange's identity $\|\mathbf{L}\|^2 = \langle \mathbf{q}, \mathbf{q} \rangle \langle \mathbf{p}, \mathbf{p} \rangle - \langle \mathbf{q}, \mathbf{p} \rangle^2$ gives

$$\langle \mathbf{A}, \mathbf{q} \rangle = \|\mathbf{L}\|^2 - mk\|\mathbf{q}\|. \tag{1}$$

If $\mathbf{A}$ and $\mathbf{L}$ are constants of motion and θ denotes the angle between $\mathbf{A}$ and $\mathbf{q}$, (1) reads $\|\mathbf{q}\|(mk + \|\mathbf{A}\| \cos\theta) = \|\mathbf{L}\|^2$, that is $\|\mathbf{q}\| = \frac{\|\mathbf{L}\|^2/mk}{1 + e\cos\theta}$ with eccentricity $e = \|\mathbf{A}\|/mk$: a conic with focus at the origin. A direct expansion also gives

$$\|\mathbf{A}\|^2 = (mk)^2 + 2mH\|\mathbf{L}\|^2, \tag{2}$$

so for $k > 0$ and $\mathbf{L} \neq 0$, $e < 1$ if and only if $H < 0$. Elementary derivations of these facts can be found in [2, 3].

3 Formalization

3.1 Setting and conventions

Physlib models phase space for a particle in three dimensions as pairs $(\mathbf{p}, \mathbf{q})$ of elements of `EuclideanSpace` $\mathbb{R}$ `(Fin 3)`, and observables as curried functions `p` $\to$ `q` $\to \mathbb{R}$. The Poisson bracket is

$$\{f, g\} = \langle \nabla_{\mathbf{q}} f, \nabla_{\mathbf{p}} g \rangle - \langle \nabla_{\mathbf{p}} f, \nabla_{\mathbf{q}} g \rangle,$$

and a central-force system is a structure carrying a mass $m > 0$ and a potential `V : ℝ → ℝ` smooth on $(0, \infty)$. We follow these conventions exactly so that the development can be contributed upstream. The Kepler problem and the LRL vector are:

```
noncomputable def kepler (m k : ℝ) (hm : 0 < m) : CentralForce where
  m := m
  V := fun r => -k / r
  m_pos := hm
  contDiffOn_V := contDiffOn_const.div contDiffOn_id
    fun _ hr => (Set.mem_Ioi.mp hr).ne'

noncomputable def lrlVector (m k : ℝ) (p q : EuclideanSpace ℝ (Fin 3)) :
    EuclideanSpace ℝ (Fin 3) :=
  WithLp.toLp 2 (p.ofLp ×₃ (angularMomentum p q).ofLp) - (m * k / ‖q‖) • q
```

The definition of `lrlVector` mirrors the textbook formula $\mathbf{p}\times\mathbf{L}-(mk/\|\mathbf{q}\|)\mathbf{q}$, reusing Physlib's `angularMomentum` and Mathlib's cross product on `Fin 3 → ℝ`. The `toLp/ofLp` coercions are needed because the cross product lives on the plain function type, whereas norms and inner products live on `EuclideanSpace`.

Remark 1. Lean's convention $x/0 = 0$ makes `lrlVector` total. A consequence we exploited is that the orbit equation (1) holds *without* the hypothesis $\mathbf{q} \neq 0$: at $\mathbf{q} = 0$ both sides vanish. In contrast, (2) genuinely requires $\mathbf{q} \neq 0$.

3.2 Algebraic identities

```
lemma lrlVector_eq (m k : ℝ) (p q : EuclideanSpace ℝ (Fin 3)) :
    lrlVector m k p q = ⟪p, p⟫_ℝ • q - ⟪q, p⟫_ℝ • p - (m * k / ‖q‖) • q

lemma inner_lrlVector_angularMomentum (m k : ℝ) (p q : EuclideanSpace ℝ (Fin 3)) :
    ⟪lrlVector m k p q, angularMomentum p q⟫_ℝ = 0

lemma inner_lrlVector_position (m k : ℝ) (p q : EuclideanSpace ℝ (Fin 3)) :
    ⟪lrlVector m k p q, q⟫_ℝ = angularMomentumSq p q - m * k * ‖q‖

lemma norm_lrlVector_sq (m k : ℝ) (hm : 0 < m) (p : EuclideanSpace ℝ (Fin 3))
    {q : EuclideanSpace ℝ (Fin 3)} (hq : q ≠ 0) :
    ‖lrlVector m k p q‖ ^ 2 =
      (m * k) ^ 2 + 2 * m * (kepler m k hm).hamiltonian p q * angularMomentumSq p q

lemma norm_lrlVector_lt_iff (m k : ℝ) (hm : 0 < m) (hk : 0 < k)
    (p : EuclideanSpace ℝ (Fin 3)) {q : EuclideanSpace ℝ (Fin 3)} (hq : q ≠ 0)
    (hL : angularMomentumSq p q ≠ 0) :
    ‖lrlVector m k p q‖ < m * k ↔ (kepler m k hm).hamiltonian p q < 0
```

The triple-product expansion and the orthogonality to $\mathbf{L}$ are proved componentwise: after `ext i; fin_cases i` the goals become polynomial identities in the six coordinates (with $mk/\|\mathbf{q}\|$ as an opaque atom), closed by `simp` and `ring`. All later results are stated in terms of inner products and norms and use `lrlVector_eq` to avoid coordinates. For (2) we expand $\langle \mathbf{A}, \mathbf{A}\rangle$ bilinearly, rewrite with Physlib's Lagrange identity `angularMomentumSq_eq`, clear the denominator $\|\mathbf{q}\|$ with `field_simp`, and close with `ring`. One subtlety: rewriting $\langle x, x\rangle = \|x\|^2$ must happen *after* bilinear expansion, otherwise it fires on $\langle c\,\mathbf{q}, c\,\mathbf{q}\rangle$ and blocks the expansion. The eccentricity criterion follows from (2) by comparing squares of non-negative reals and a single `nlinarith` call with the hint $m\|\mathbf{L}\|^2 > 0$.

3.3 Conservation of the LRL vector

For each component A_i we compute the partial gradients explicitly:

$$\nabla_{\mathbf{p}} A_i = 2q_i\,\mathbf{p} - p_i\,\mathbf{q} - \langle \mathbf{q}, \mathbf{p}\rangle\,\mathbf{e}_i,$$
$$\nabla_{\mathbf{q}} A_i = \langle \mathbf{p}, \mathbf{p}\rangle\,\mathbf{e}_i - p_i\,\mathbf{p} - \frac{mk}{\|\mathbf{q}\|}\,\mathbf{e}_i + \frac{mk\,q_i}{\|\mathbf{q}\|^3}\,\mathbf{q} \qquad (\mathbf{q} \neq 0).$$

In Lean these are `HasGradientAt` statements, assembled from Physlib's gradient calculus (`hasGradientAt_inner_self`, `hasGradientAt_coord`, `hasGradientAt_norm` and a chain rule `HasDerivAt.comp_hasGradientAt`), followed by `module` or `field_simp` to normalize the resulting linear combination. Using $V'(r) = k/r^2$ (lemma `kepler_deriv_V`) and Physlib's gradients of the Hamiltonian, $\nabla_{\mathbf{p}} H = \mathbf{p}/m$ and $\nabla_{\mathbf{q}} H = (k/\|\mathbf{q}\|^3)\,\mathbf{q}$, the Poisson bracket evaluates to

$$\{A_i, H\} = -k\Big(\frac{p_i}{\|\mathbf{q}\|} - \frac{q_i\langle \mathbf{q}, \mathbf{p}\rangle}{\|\mathbf{q}\|^3}\Big) - \frac{k}{\|\mathbf{q}\|^3}\big(q_i\langle \mathbf{q}, \mathbf{p}\rangle - p_i\|\mathbf{q}\|^2\big) = 0.$$

```
lemma poissonBracket_lrlVector_hamiltonian (m k : ℝ) (hm : 0 < m) (i : Fin 3)
    (p : EuclideanSpace ℝ (Fin 3)) {q : EuclideanSpace ℝ (Fin 3)} (hq : q ≠ 0) :
    poissonBracket (fun p' q' => lrlVector m k p' q' i)
      (kepler m k hm).hamiltonian p q = 0
```

The proof is eleven lines: rewrite the bracket via `poissonBracket_eq_of_hasGradientAt` with the four gradients, expand inner products, and close with `field_simp; ring`.

3.4 From phase space to trajectories

A vanishing Poisson bracket is the infinitesimal statement of conservation. To obtain Kepler's first law we need the integrated statement along an actual motion. Physlib provides Hamilton's equations as `hamiltonEqOp H p q = 0` for curves `p q : Time → EuclideanSpace ℝ (Fin 3)`, and the theorem `deriv_comp_eq_poissonBracket` stating $\frac{d}{dt} f(\mathbf{p}(t), \mathbf{q}(t)) = \{f, H\}(\mathbf{p}(t), \mathbf{q}(t))$ along solutions. We proved a general lemma:

```
lemma eq_of_poissonBracket_eq_zero {H f : E → E → ℝ} {p q : Time → E}
    (heq : hamiltonEqOp (fun _ => H) p q = 0)
    (hp : Differentiable ℝ p) (hq : Differentiable ℝ q)
    (hf : ∀ t, DifferentiableAt ℝ (Function.uncurry f) (p t, q t))
    (hpb : ∀ t, poissonBracket f H (p t) (q t) = 0) (t s : Time) :
    f (p t) (q t) = f (p s) (q s)
```

(here `E` abbreviates `EuclideanSpace ℝ (Fin 3)`). Physlib's `Time` is a structure wrapping a real number rather than $\mathbb{R}$ itself; the proof transports the derivative to $\mathbb{R}$ with the bridge lemma `hasDerivAt_mk_of_differentiableAt` and applies Mathlib's `is_const_of_deriv_eq_zero`. Together with differentiability of A_i away from the origin and the existing conservation of $\|\mathbf{L}\|^2$, this yields:

Theorem 2 (Kepler's first law)**.** *Let $(\mathbf{p}(t), \mathbf{q}(t))$ be a differentiable solution of Hamilton's equations for the Kepler Hamiltonian with $\mathbf{q}(t) \neq 0$ for all t. Then for all times t_0, t,*

$$\langle \mathbf{A}(t_0), \mathbf{q}(t)\rangle = \|\mathbf{L}(t_0)\|^2 - mk\,\|\mathbf{q}(t)\|.$$

```
lemma inner_lrlVector_position_of_hamiltonEqOp (m k : ℝ) (hm : 0 < m)
    {p q : Time → EuclideanSpace ℝ (Fin 3)}
    (heq : hamiltonEqOp (fun _ => (kepler m k hm).hamiltonian) p q = 0)
    (hp : Differentiable ℝ p) (hq : Differentiable ℝ q) (hq0 : ∀ t, q t ≠ 0)
    (t₀ t : Time) :
    ⟪lrlVector m k (p t₀) (q t₀), q t⟫_ℝ =
      angularMomentumSq (p t₀) (q t₀) - m * k * ‖q t‖
```

Combined with `norm_lrlVector_lt_iff`, this shows that a bound Kepler motion with non-zero angular momentum traces an ellipse. Note that we assume, rather than construct, a solution of Hamilton's equations; existence and uniqueness of solutions are not needed for these statements and are not addressed.

4 The hidden $\mathfrak{so}(4)$ symmetry

The closed orbits of the Kepler problem reflect a symmetry larger than rotations. Together with $\mathbf{L}$, the components of $\mathbf{A}$ close under the Poisson bracket, and on each negative energy surface they generate $\mathfrak{so}(4)$ [2, §§9.7–9.8]. This is the classical shadow of Pauli's derivation of the hydrogen spectrum [5] and of the $O(4)$ symmetry of hydrogen [1]. Physlib formalizes the quantum version: an abstract Pauli system of operators with the hydrogen commutation relations, from which the Bohr levels follow. We prove the classical statements. They live in a separate file, `KeplerLRL/SO4.lean`.

4.1 Statements without the Levi-Civita symbol

Rather than introducing ε_{ijk}, we state every relation for components along arbitrary fixed vectors $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3$, writing $X_{\mathbf{a}} = \langle \mathbf{a}, \mathbf{X} \rangle$. Bilinearity then gives the usual index form by choosing basis vectors, for example $\{L_i, L_j\} = L_{\mathbf{e}_i \times \mathbf{e}_j} = \varepsilon_{ijk} L_k$. With `euclideanCross a b` denoting $\mathbf{a} \times \mathbf{b}$ on `EuclideanSpace ℝ (Fin 3)`, the three structure relations read:

```
lemma poissonBracket_angularMomentum_angularMomentum (a b p q : E) :
    poissonBracket (fun p q => ⟪a, angularMomentum p q⟫_ℝ)
      (fun p q => ⟪b, angularMomentum p q⟫_ℝ) p q =
      ⟪euclideanCross a b, angularMomentum p q⟫_ℝ

lemma poissonBracket_lrlVector_angularMomentum (m k : ℝ) (a b p : E) {q : E}
    (hq : q ≠ 0) :
    poissonBracket (fun p q => ⟪a, lrlVector m k p q⟫_ℝ)
      (fun p q => ⟪b, angularMomentum p q⟫_ℝ) p q =
      ⟪euclideanCross a b, lrlVector m k p q⟫_ℝ

lemma poissonBracket_lrlVector_lrlVector (m k : ℝ) (hm : 0 < m) (a b p : E) {q : E}
    (hq : q ≠ 0) :
    poissonBracket (fun p q => ⟪a, lrlVector m k p q⟫_ℝ)
      (fun p q => ⟪b, lrlVector m k p q⟫_ℝ) p q =
      -2 * m * (kepler m k hm).hamiltonian p q *
        ⟪euclideanCross a b, angularMomentum p q⟫_ℝ
```

We also prove $\{L_{\mathbf{a}}, H\} = 0$ for an arbitrary central force (Physlib previously had this only for $\|\mathbf{L}\|^2$ and L_z) and $\{A_{\mathbf{a}}, H\} = 0$ for the Kepler problem.

Proofs. The partial gradients of $L_{\mathbf{a}} = \langle \mathbf{a}, \mathbf{q} \times \mathbf{p} \rangle$ are $\nabla_{\mathbf{p}} L_{\mathbf{a}} = \mathbf{a} \times \mathbf{q}$ and $\nabla_{\mathbf{q}} L_{\mathbf{a}} = \mathbf{p} \times \mathbf{a}$, and those of $A_{\mathbf{a}}$ are the formulas of §3.3 with $\mathbf{e}_i$ replaced by $\mathbf{a}$. The first two brackets are then polynomial identities in coordinates, with $\|\mathbf{q}\|$ treated as an opaque atom. This works because the only term carrying $\|\mathbf{q}\|^{-3}$ is multiplied by $\langle \mathbf{q}, \mathbf{b} \times \mathbf{q} \rangle$, which vanishes identically. The third bracket is different: its $\|\mathbf{q}\|^{-3}$ terms combine to $\langle \mathbf{q}, \mathbf{q} \rangle \, \|\mathbf{q}\|^{-3}$ and must be simplified with $\langle \mathbf{q}, \mathbf{q} \rangle = \|\mathbf{q}\|^2$, so a coordinate expansion fails. Instead we rewrite the right-hand side with the Binet–Cauchy identity $\langle \mathbf{a} \times \mathbf{b}, \mathbf{q} \times \mathbf{p} \rangle = \langle \mathbf{a}, \mathbf{q} \rangle \langle \mathbf{b}, \mathbf{p} \rangle - \langle \mathbf{a}, \mathbf{p} \rangle \langle \mathbf{b}, \mathbf{q} \rangle$ and keep everything in terms of inner products. Since `simp` uses ordered rewriting for permutation lemmas, passing it `real_inner_comm` puts every inner product in a canonical argument order, after which `field_simp; ring` closes the goal.

4.2 The rescaled vector and $\mathfrak{so}(3) \oplus \mathfrak{so}(3)$

On the bound region $H < 0$ define $\mathbf{M} = \mathbf{A}/\sqrt{-2mH}$ and $\mathbf{J}_{\pm} = (\mathbf{L} \pm \mathbf{M})/2$. In Lean, `scaledLrlVector` multiplies $\mathbf{A}$ by $c(H)$ with $c(E) = (\sqrt{-2mE})^{-1}$. The definition is total, but the theorems assume $H < 0$. Writing $M_{\mathbf{a}} = c(H)\, A_{\mathbf{a}}$, the bracket $\{M_{\mathbf{a}}, M_{\mathbf{b}}\}$ expands by Physlib's Leibniz rules into $c^2 \{A_{\mathbf{a}}, A_{\mathbf{b}}\}$ plus terms containing $\{c(H), A_{\mathbf{b}}\}$ and $\{A_{\mathbf{a}}, c(H)\}$. A small chain-rule lemma we add,

$$\{\varphi \circ f, g\} = \varphi'(f)\, \{f, g\},$$

reduces these to multiples of $\{H, A\} = 0$. Together with $c^2 \cdot (-2mH) = 1$ this gives $\{M_{\mathbf{a}}, M_{\mathbf{b}}\} = L_{\mathbf{a} \times \mathbf{b}}$. In the same way, $\{M_{\mathbf{a}}, L_{\mathbf{b}}\} = \{L_{\mathbf{a}}, M_{\mathbf{b}}\} = M_{\mathbf{a} \times \mathbf{b}}$. A single bilinear-expansion lemma then handles all three brackets of $\mathbf{J}_{\pm}$ at once. For $s, t \in \mathbb{R}$,

$$\left\{ \tfrac{1}{2}(L_{\mathbf{a}} + s M_{\mathbf{a}}), \tfrac{1}{2}(L_{\mathbf{b}} + t M_{\mathbf{b}}) \right\} = \tfrac{1}{4}\big((1 + st)\, L_{\mathbf{a} \times \mathbf{b}} + (s + t)\, M_{\mathbf{a} \times \mathbf{b}}\big),$$

and $(s, t) = (1, 1), (-1, -1), (1, -1)$ give:

```
lemma poissonBracket_jPlus_jPlus (m k : ℝ) (hm : 0 < m) (a b p : E) {q : E}
    (hq : q ≠ 0) (hH : (kepler m k hm).hamiltonian p q < 0) :
    poissonBracket (fun p q => ⟪a, jPlus m k hm p q⟫_ℝ)
      (fun p q => ⟪b, jPlus m k hm p q⟫_ℝ) p q =
      ⟪euclideanCross a b, jPlus m k hm p q⟫_ℝ

lemma poissonBracket_jPlus_jMinus (m k : ℝ) (hm : 0 < m) (a b p : E) {q : E}
    (hq : q ≠ 0) (hH : (kepler m k hm).hamiltonian p q < 0) :
    poissonBracket (fun p q => ⟪a, jPlus m k hm p q⟫_ℝ)
      (fun p q => ⟪b, jMinus m k hm p q⟫_ℝ) p q = 0
```

together with the analogue `poissonBracket_jMinus_jMinus`. So $\mathbf{J}_+$ and $\mathbf{J}_-$ each satisfy the $\mathfrak{so}(3)$ relations and commute with each other, which is the decomposition $\mathfrak{so}(4) \cong \mathfrak{so}(3) \oplus \mathfrak{so}(3)$.

4.3 The Casimirs and the energy

The two Casimirs of $\mathfrak{so}(4)$ are $\langle \mathbf{L}, \mathbf{M} \rangle$ and $\|\mathbf{L}\|^2 + \|\mathbf{M}\|^2$. The first vanishes by $\langle \mathbf{A}, \mathbf{L} \rangle = 0$. For the second, (2) gives $\|\mathbf{M}\|^2 = ((mk)^2 + 2mH\|\mathbf{L}\|^2)/(-2mH)$, so $\|\mathbf{L}\|^2 + \|\mathbf{M}\|^2 = -mk^2/(2H)$. Hence both $\mathfrak{so}(3)$ factors have the same Casimir, and the energy is a function of it:

Theorem 3. *On the bound region $H < 0$, away from the origin, $\|\mathbf{J}_+\|^2 = \|\mathbf{J}_-\|^2 = -\dfrac{mk^2}{8H}$. If moreover $k \neq 0$, then $H = -\dfrac{mk^2}{8\|\mathbf{J}_+\|^2}$.*

```
lemma norm_jPlus_sq (m k : ℝ) (hm : 0 < m) (p : E) {q : E} (hq : q ≠ 0)
    (hH : (kepler m k hm).hamiltonian p q < 0) :
    ‖jPlus m k hm p q‖ ^ 2 = -(m * k ^ 2) / (8 * (kepler m k hm).hamiltonian p q)
```

In the quantum case the corresponding Casimir is $\mathbf{J}^2 = -(\hbar^2 + mk^2/2E)/4$. Imposing $\mathbf{J}^2 = \hbar^2 j(j+1)$ gives the Bohr levels $E = -mk^2/(2\hbar^2 n^2)$ with $n = 2j+1$ [5]. Setting $\hbar = 0$ recovers exactly our classical relation. Classically $\|\mathbf{J}_+\|$ can take any positive value, and the relation expresses the energy through the $\mathfrak{so}(4)$ Casimir. Since every observable here Poisson-commutes with H, the lemma `eq_of_poissonBracket_eq_zero` of §3.4 also makes $\mathbf{M}$ and $\mathbf{J}_\pm$ constant along bound trajectories. We have not stated that corollary separately.

5 Discussion

Size and checking. The development consists of 7 definitions and 50 lemmas in about 860 lines, including documentation: about 300 lines for §§3.1–3.4 and 550 for §4. The longest proof has 18 lines. `#print axioms` on the main results reports only `propext`, `Classical.choice` and `Quot.sound`. The file for §§3.1–3.4 passes Physlib's module-documentation, alpha-library and style linters. The $\mathfrak{so}(4)$ file has not yet been run through them.

What was easy. Once the right normal form (`lrlVector_eq`) was available, every identity of the Kepler problem reduced to `field_simp` and `ring`. Physlib's existing gradient API and its Poisson-bracket lemma `poissonBracket_eq_of_hasGradientAt` made the conservation proof essentially a transcription of the pen-and-paper computation.

What required care. The friction points were elaboration issues rather than mathematics: the higher-order unification of `DifferentiableAt.comp` and of the chain rule for gradients needed explicit named arguments; `fun_prop` could not discharge differentiability of $\mathbf{q} \mapsto q_i$ on `EuclideanSpace` without hand-supplied facts; and moving between `Time` and $\mathbb{R}$ required Physlib's bridge lemmas. In §4 the main decision was where to work in coordinates and where in inner products: coordinates suit identities that are polynomial once $\|\mathbf{q}\|$ is treated as an atom, and inner products suit identities that need $\langle \mathbf{q}, \mathbf{q} \rangle = \|\mathbf{q}\|^2$.

Future work. Natural continuations are Kepler's second law (constant areal velocity, essentially the conservation of $\mathbf{L}$ along trajectories), Kepler's third law relating the period to the semi-major axis, and the global form of the hidden symmetry: integrating the $\mathfrak{so}(4)$ action to an $SO(4)$ action on each negative energy surface, as in Moser's regularization of the Kepler flow to the geodesic flow on S^3 [4]. Bertrand's theorem, which singles out the Kepler and harmonic potentials by the closure of bounded orbits, is formalized in a companion development. For an upstream contribution to Physlib, the general lemma `eq_of_poissonBracket_eq_zero` should move to the Poisson-bracket module, and the work should be split into several smaller pull requests as required by the project's guidelines.

Author contributions. N.L. chose the problem, directed the work and reviewed the results. Claude, an AI system developed by Anthropic, surveyed the existing literature and libraries, designed the statements, wrote the Lean proofs and drafted the manuscript. All statements were checked by the Lean kernel. As an AI system cannot take responsibility for a publication, N.L. is accountable for its content.

References

[1] M. Bander and C. Itzykson. Group theory and the hydrogen atom (I). *Reviews of Modern Physics*, 38:330–345, 1966.

[2] Herbert Goldstein, Charles Poole, and John Safko. *Classical Mechanics*. Addison-Wesley, 3rd edition, 2002.

[3] M. Moriconi. Elementary solution of Kepler problem (and a few other problems), 2021.

[4] Jürgen Moser. Regularization of Kepler's problem and the averaging method on a manifold. *Communications on Pure and Applied Mathematics*, 23:609–636, 1970.

[5] Wolfgang Pauli. Über das Wasserstoffspektrum vom Standpunkt der neuen Quantenmechanik. *Zeitschrift für Physik*, 36:336–363, 1926.

[6] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, pages 367–381. ACM, 2020.

[7] Joseph Tooby-Smith. HepLean: Digitalising high energy physics, 2024.