



**CERTIFICATE  
ISOTOPIC REFERENCE MATERIAL (SPIKE) IRMM-618**

$$(1.121\ 3 \pm 0.001\ 7) \cdot 10^{-4} \text{ mol } ^{87}\text{Rb} \cdot \text{kg}^{-1} \text{ of solution}$$

The Isotopic Reference Material (Spike) is supplied with a molar concentration of  $^{87}\text{Rb}$  certified as above.

Other rubidium isotopes present are related to the  $^{87}\text{Rb}$  concentration through the following certified molar ratio :

$$n(^{85}\text{Rb})/n(^{87}\text{Rb}) : 0.020\ 498 \quad \pm 0.000\ 024$$

This corresponds to an isotopic composition with following abundances :

	Amount %	Mass %	Uncertainty
$^{85}\text{Rb}$	2.008 7	1.963 4	$\pm 0.002\ 3$
$^{87}\text{Rb}$	97.991 3	98.036 6	$\pm 0.002\ 3$

The molar mass of the rubidium is  $(86.869\ 066 \pm 0.000\ 045) \text{ g} \cdot \text{mol}^{-1}$

From the certified values, the following element concentrations are derived :

$$\begin{aligned} (9.743 \quad \pm 0.015 \quad ) \cdot 10^{-6} \text{ kg } ^{87}\text{Rb} \cdot \text{kg}^{-1} \text{ of solution} \\ (9.940 \quad \pm 0.015 \quad ) \cdot 10^{-6} \text{ kg } \text{Rb} \cdot \text{kg}^{-1} \text{ of solution} \\ (1.144\ 3 \quad \pm 0.001\ 7 \quad ) \cdot 10^{-4} \text{ mol } \text{Rb} \cdot \text{kg}^{-1} \text{ of solution} \end{aligned}$$

## NOTES

1. All uncertainties indicated are levels of possible inaccuracies, computed on a 2s basis for all components. These include : measurement reproducibility, reproducibilities of measurements of correction factors for known systematic errors and possible uncertainties of systematic nature estimated on a 2s equivalence basis.
2. IRMM-618 consists of flame-sealed quartz glass ampoules containing about 5 ml of a chemically stable solution of rubidium in nitric acid. The molality is about 0.5 m HNO<sub>3</sub> (i.e. 0.5 mol HNO<sub>3</sub> · kg<sup>-1</sup> of solvent); the molarity is about 0.5 M HNO<sub>3</sub> (i.e. 0.5 mol HNO<sub>3</sub> · L<sup>-1</sup> of solution).
3. The Avogadro constant used is (6.022 136 ± 0.000 012) · 10<sup>23</sup> mol<sup>-1</sup>.
4. The molar masses, used in the calculations, are  

$$^{85}\text{Rb} \quad : \quad (84.914\,390 \pm 0.000\,008) \text{ g} \cdot \text{mol}^{-1}$$

$$^{87}\text{Rb} : (86.911\,794 \pm 0.000\,006) \text{ g} \cdot \text{mol}^{-1}$$
5. Using this Spike Reference Material (Spike), <sup>85</sup>Rb and Rb concentrations in unknown samples can be determined by Isotope Dilution Mass Spectrometry, through a measurement of the molar isotope dilution ratio  $R_B = n(^{85}\text{Rb})/n(^{87}\text{Rb})$  in the blend. They should be computed with the aid of the following formula which allows an easy identification of the sources of the uncertainties in the procedure :

$$c(^{85}\text{Rb})_X = \frac{R_Y - R_B}{R_B - R_X} \cdot R_X \cdot \frac{m_Y}{m_X} \cdot c(^{87}\text{Rb})_Y$$

$$c(\text{Rb})_X = \frac{R_Y - R_B}{R_B - R_X} \cdot \frac{1 + R_X}{1 + R_Y} \cdot \frac{m_Y}{m_X} \cdot c(\text{Rb})_Y$$

where

$R_X$	=	molar ratio $n(^{85}\text{Rb})/n(^{87}\text{Rb})$ in the unknown sample material
$R_Y$	=	molar abundance ratio $n(^{85}\text{Rb})/n(^{87}\text{Rb})$ in the spike material
$m_X$	=	mass of the unknown sample
$m_Y$	=	mass of the sample of spike solution used
$c(^{85}\text{Rb})_Y$	=	number of moles <sup>85</sup> Rb · kg <sup>-1</sup> sample material
$c(^{87}\text{Rb})_X$	=	number of moles <sup>87</sup> Rb · kg <sup>-1</sup> spike solution
$c(\text{Rb})_X$	=	number of moles Rb · kg <sup>-1</sup> sample material
$c(\text{Rb})_Y$	=	number of moles Rb · kg <sup>-1</sup> spike solution.

6. This Isotopic Reference Material (Spike) is traceable to the international SI unit for amount of substance - the mole - in the shortest possible way. Measurements calibrated against these Isotopic Reference Materials will, therefore, also be traceable to the SI unit system.

The isotopic measurements by Thermal Ionisation Mass Spectrometry were performed by G. Lapitajs.

Chemical preparation of the samples for isotopic measurements was performed by G. Lapitajs and A. Verbruggen.

The purity of  $^{87}\text{Rb}$  was determined by Spark Source Mass Spectrometry by W. Lycke.

Metrological aspects involved in the preparation and certification were performed by F. Hendrickx. The ampoulation of this Isotopic Reference Material (Spike) was accomplished by G. Lapitajs, A. Alonso-Muñoz and A. Verbruggen.

The overall co-ordination leading to the establishment, certification and issuance of this Isotopic Reference Material (Spike), was performed by A. Verbruggen.

B-2440 Geel  
April 1989  
Revised July 1993  
Revised December 1996

P. DE BIEVRE  
Head  
EC-IRMM Stable Isotope Measurements