

Week 24.11 - 28.11

- continue writing
- compute manifolds with B_m corrected
- compute 1-D line function for comparing distance between peak
- compute manifolds for new jellyfish shot


$$\Delta \mathbf{b}_m^{\text{na}} \simeq -(\Delta \mathbf{X}_c \cos(\theta_m) + \Delta \mathbf{Y}_c \sin(\theta_m)) \mathbf{I}_c \frac{\partial \mathbf{B}_{mc}}{\partial \mathbf{r}_m} - (\Delta \mathbf{Z}_c^{\text{oy}} \cos(\theta_m) + \Delta \mathbf{Z}_c^{\text{ox}} \sin(\theta_m)) \mathbf{I}_c \frac{\partial \mathbf{B}_{mc}}{\partial \mathbf{z}_m}$$



$$\mathbf{b}_m^{\text{corr}} = \mathbf{b}_m - \Delta \mathbf{b}_m^{\text{na}}$$



$$\Delta \mathbf{b}^{\text{corr}} = \sum_i |\mathbf{b}_3^{\text{corr}} - \mathbf{b}_{11}^{\text{corr}}|_i = 0.0438$$

$$\Delta \mathbf{b} = \sum_i |\mathbf{b}_3 - \mathbf{b}_{11}|_i = 0.0947$$

- then re-compute LIUQE with corrected Bm







to do:

- write my master's thesis
- compute trace of fixed point in “legged” vs “non legged” shots
- Get fastcam image
- Compute manifold for X-point target and shot 61'848
- compute ratio of peak distance MF/ MANTIS to approximate mismatch of B_z