



Pieri and Murnaghan–Nakayama Type Rules for Chern classes of Schubert Cells

Neil J.Y. Fan¹, Peter L. Guo², and Rui Xiong³

¹Sichuan University



²Nankai University



³University of Ottawa



1 Introduction

We develop **Pieri type** as well as **Murnaghan–Nakayama (MN) type** formulas for equivariant **Chern–Schwartz–MacPherson** classes of Schubert cells in the classical flag variety. These formulas include as special cases many previously known multiplication formulas for Chern–Schwartz–MacPherson classes or Schubert classes. We apply the equivariant Murnaghan–Nakayama formula to the enumeration of **rim hook tableaux**.

2 Classic Pieri Rules

Historically, the classical Pieri formula provides a solution to the multiplication of a **Schur polynomial** by an **elementary** or a **complete homogeneous symmetric polynomial**. For example,


$$\begin{array}{|c|c|} \hline & \\ \hline \end{array} \cdot e_3 = \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 2 & \\ \hline 3 & & \end{array} + \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 2 & \\ \hline 3 & & \end{array} + \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 2 & \\ \hline 3 & & \end{array} + \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 2 & \\ \hline 3 & & \end{array}$$

This formula is generalized to **Schubert polynomials** times hook shape Schur polynomial by Sottile [9]. The formula is given by chasing over **k-Bruhat graph**. For example,

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \cdot s_{(2,1)}(x_1, x_2, x_3) = \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} + \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}$$

23154 45123 35142


The graph is (black arrows)



Geometrically, elementary or complete homogeneous symmetric polynomial serves as not only Schubert classes but also the **Chern classes** or **Segre classes** of the tautological bundle over flag variety. We generalize the classic Pieri rule to Chern–Schwartz–MacPherson classes.

3 CSM Pieri rules

The **Chern–Schwartz–MacPherson (CSM)** classes are one way to extend the Chern classes of complex manifolds to complex varieties (possibly singular or noncompact). The structure of (equivariant) CSM classes for Schubert cells in flag varieties has received much attention in recent years and devolved into a rich area of research, see [1, 2]. Our formula is given by chasing over the **extended k-Bruhat graph**. For example,



One of the key ingredients we developed is the **Rigidity Theorem**, which means that the structure constants for equivariant rules can be controlled by the structure constants in their nonequivariant situations.

When restricting to Grassmannian, we can get a description in terms of partition combinatorics. For example, in $H^\bullet(Gr(3, 7))$, we have

$$\begin{array}{|c|c|} \hline & \\ \hline \end{array} \cdot e_3 = \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 2 & \\ \hline 3 & & \end{array} + \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 2 & \\ \hline 3 & & \end{array} + \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 2 & \\ \hline 3 & & \end{array} + \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 2 & \\ \hline 3 & & \end{array}$$

Note that all partitions cannot exceed the rectangle (4, 4, 4).

4 Hook formulas


Starting from the work of Ikeda and Naruse [5], it was gradually realized that **equivariant Chevalley type** formulas could be utilized to deduce **hook formulas**, see Naruse [6] and the recent work of Morales, Pak and Panova [8] and Mihalcea, Naruse and Su [7].

We generalize the idea of [5] to higher degrees by viewing the equivariant MN formula as a **higher degree analogue** of the equivariant Chevalley formula. It amazingly applies to the enumeration of **rim hook tableaux**. We reveal the enumeration formulas for standard rim hook tableaux due to Alexandersson, Pfannerer, Rubey and Uhlin [3] and Fomin and Lulov [4] in a relatively uniform manner.

$$\# \left\{ \begin{array}{|c|c|c|c|} \hline 1 & 2 & 1 & 3 \\ \hline 3 & 4 & 2 & 4 \\ \hline 1 & 2 & 3 & 4 \\ \hline 3 & 4 & 1 & 2 \\ \hline \end{array} \right\} = \frac{8!!}{4 \times 2 \times 4 \times 2}.$$

$$\# \left\{ \begin{array}{|c|c|c|} \hline 1 & 2 & 1 \\ \hline 3 & 4 & 2 \\ \hline 1 & 2 & 3 \\ \hline 3 & 4 & 1 \\ \hline \end{array} \right\} = \frac{8!!}{6 \times 4 \times 4 \times 2}.$$

where the denominator is the product of all **even hook lengths**.



References

- [1] P. Aluffi and L. Mihalcea, Chern–Schwartz–MacPherson classes for Schubert cells in flag manifolds. *Compos. Math.* 152(12), 2603–2625 (2016)
- [2] P. Aluffi, L. Mihalcea, Jörg Schürmann, and C. Su, Shadows of characteristic cycles, Verma modules, and positivity of Chern–Schwartz–MacPherson classes of Schubert cells. *arXiv:1709.08697*, 2017.
- [3] P. Alexandersson, S. Pfannerer, M. Rubey and J. Uhlin, Skew characters and cyclic sieving, *Forum Math. Sigma* 9 (2021), e41.
- [4] S. Fomin and N. Lulov, On the number of rim hook tableaux *J. Math. Sci. (New York)*, 87 (1997), pp. 4118–4123
- [5] T. Ikeda and H. Naruse, Excited Young diagrams and equivariant Schubert calculus, *Trans. Amer. Math. Soc.* 361 (2009), 5193–5221.
- [6] H. Naruse, Schubert calculus and hook formula, talk slides at 73rd Sem. Lothar. Combin., Strobl, Austria, 2014, available at <https://www.emis.de/journals/SLC/wpapers/s73vortrag/naruse.pdf>.
- [7] L. Mihalcea, H. Naruse and Changjian Su, Hook formulae from Segre–MacPherson classes, *arXiv:2203.16461v1*.
- [8] A.H. Morales, I. Pak and G. Panova, Hook formulas for skew shapes IV. increasing tableaux and factorial Grothendieck polynomials, *J. Math. Sci. (N.Y.)* 261 (2022), 630–657.
- [9] F. Sottile, Pieri’s formula for flag manifolds and Schubert polynomials, *Annales de l’Institut Fourier* 46(1996), pp.89–110