

Bumpless pipe dreams meet Puzzles


(joint with Neil J.Y. Fan and Peter L. Guo)

Rui Xiong

December 23, 2023



meet



Linear algebra

Denote the *standard opposite flag*

$$F^0 = \mathbb{C}^n \geq \dots \geq F^{n-2} = \langle \mathbf{e}_n, \mathbf{e}_{n-1} \rangle \geq F^{n-1} = \langle \mathbf{e}_n \rangle \geq F^n = 0.$$

For each $V \leq \mathbb{C}^n$ of dimension k , we have a decreasing flag

$$V = F^0 \cap V \geq \dots \geq F^{n-2} \cap V \geq F^{n-1} \cap V \geq F^n \cap V = 0.$$

We can assign the set of “jumping indices” λ , i.e.

$$\lambda_i = 1 \iff \dim(F^{i-1} \cap V) > \dim(F^i \cap V)$$

$$\lambda_i = 0 \iff \dim(F^{i-1} \cap V) = \dim(F^i \cap V)$$

Grassmannians

Denote

$$\mathrm{Gr}(k, n) = \{ V \leq \mathbb{C}^n \mid \dim V = k \}.$$

Let us denote *Schubert cell* for $\lambda \in \binom{[n]}{k}$

$$\Sigma_\lambda^\circ = \left\{ V \in \mathrm{Gr}(k, n) \mid \text{jumping indices of } V = \lambda \right\}.$$

$$\Sigma_\lambda = \text{closure of } \Sigma_\lambda^\circ, \quad \sigma_\lambda = [\Sigma_\lambda^\circ] \in H^\bullet(\mathrm{Gr}(k, n)).$$











It is known that

$$H^\bullet(\mathrm{Gr}(k, n)) = \bigoplus_{\lambda \in \binom{[n]}{k}} \mathbb{Q} \cdot \sigma_\lambda \quad (\text{as a vector space})$$

Example

Let us identify

$$\text{Gr}(2, 4) = \left\{ \text{lines in } \mathbb{P}^3 \right\}.$$

ALL LINES		
1100 	1010 	0110 
		A LINE
1001 	0101 	0011 


Littlewood–Richardson coefficients

Assume


$$\sigma_\lambda \cdot \sigma_\mu = \sum_{\nu \in \binom{[n]}{k}} c_{\lambda\mu}^\nu \cdot \sigma_\nu.$$

The coefficients $c_{\lambda\mu}^\nu \in \mathbb{Z}_{\geq 0}$ are known as *Littlewood–Richardson (LR) coefficients*.


It also appears in the study of representation theory and symmetric functions. These coefficients admit a lot of combinatorial models like



Robinson–Schensted
correspondence



jeu de taquin



crystal
Schur operators

Geometric meaning

Let us denote


$$c_{\lambda\mu\nu} = c_{\lambda\mu}^{\nu^{op}}$$

Then for generic $x, y, z \in GL_n$



$$c_{\lambda\mu\nu} = \# \left\{ \nu \in \text{Gr}(k, n) \mid xV \in \Sigma_\lambda, yV \in \Sigma_\mu, zV \in \Sigma_\nu \right\}.$$

If it is empty or infinite, then it is understood as zero.

Examples



We can compute:



$$c_{1010,1010,0110} = 1 \quad c_{1010,1010,1001} = 1$$

As a result,

$$\sigma_{1010} \cdot \sigma_{1010} = \sigma_{0110} + \sigma_{1001}.$$


Puzzles


Let us use the following convention

red = 1, blue = 0.


Theorem (A. Knutson, T. Tao, and C. Woodward)

The number $c_{\lambda\mu}^{\nu}$ is the number of puzzles



Warning:  are not allowed (we cannot reflect puzzles).

Examples



$$\sigma_{1010} \cdot \sigma_{1010} = \sigma_{0110} + \sigma_{1001}.$$

Generalization A

$$H^\bullet(\mathrm{Gr}(k, n)) \rightsquigarrow K(\mathrm{Gr}(k, n))$$

Let us denote

$$\mathcal{O}_\lambda = [\mathcal{O}_{\Sigma_\lambda}] = \text{structure sheaf for } \Sigma_\lambda.$$

$$\mathcal{I}_\lambda = [\mathcal{O}_{\Sigma_\lambda}(-\partial\Sigma_\lambda)] = \text{ideal sheaf for } \partial\Sigma_\lambda = \Sigma_\lambda \setminus \Sigma_\lambda^\circ.$$

It is known that they are dual basis under the Poincaré pairing.


Similarly, we have

$$K(\mathrm{Gr}(k, n)) = \bigoplus_{\lambda \in \binom{[n]}{k}} \mathbb{Q} \cdot \mathcal{O}_\lambda = \bigoplus_{\lambda \in \binom{[n]}{k}} \mathbb{Q} \cdot \mathcal{I}_\lambda.$$

We call the coefficients of their expansion the *structure constants*.


Theorem (Vakil)

The structure constant for \mathcal{O}_λ is the number of puzzles




Theorem (Wheeler and Zinn-Justin)


The structure constant for \mathcal{I}_λ is the number of puzzles



Examples



$$\mathcal{O}_{1010} \cdot \mathcal{O}_{1010} = \mathcal{O}_{0110} + \mathcal{O}_{1001} + \mathcal{O}_{0101}$$



$$\mathcal{I}_{1010} \cdot \mathcal{I}_{1010} = \mathcal{I}_{0110} + \mathcal{I}_{1001} + \mathcal{I}_{0101} + \mathcal{I}_{0011}.$$

Generalization B

$$H^\bullet(\mathrm{Gr}(k, n)) \rightsquigarrow H_T^\bullet(\mathrm{Gr}(k, n))$$

Here we are considering the *toric equivariant cohomology*. We have


$$H_T^\bullet(\mathrm{Gr}(k, n)) = \bigoplus_{\lambda \in \binom{[n]}{k}} \mathbb{Q}[t_1, \dots, t_n] \cdot \sigma_\lambda$$

Similarly, we have *toric equivariant K-theory*

$$\begin{aligned} K_T(\mathrm{Gr}(k, n)) &= \bigoplus_{\lambda \in \binom{[n]}{k}} \mathbb{Q}[\tau_1^{\pm 1}, \dots, \tau_n^{\pm 1}] \cdot \mathcal{O}_\lambda, \\ &= \bigoplus_{\lambda \in \binom{[n]}{k}} \mathbb{Q}[\tau_1^{\pm 1}, \dots, \tau_n^{\pm 1}] \cdot \mathcal{I}_\lambda. \end{aligned}$$


Theorem (Knutson and Tao)

The structure constant for $H_T^\bullet(\mathrm{Gr}(k, n))$ can be computed by






Theorem (Pechenik and Yong, Wheeler and Zinn-Justin)

The structure constant for $K_T(\mathrm{Gr}(k, n))$ can be computed by












Examples



$$\sigma_{1010} \cdot \sigma_{1010} = (t_3 - t_2) \cdot \sigma_{1010} + \sigma_{0110} + \sigma_{1001}.$$

Summary

tiles	K-tiles	equivariant tiles
		



Flag varieties

Now we turn to *flag varieties*

$$\mathrm{Fl}(n) = \{0 = V_0 < V_1 < \cdots < V_n = \mathbb{C}^n\}.$$

For each flag $V_\bullet \in \mathrm{Fl}(n)$, we can similarly assign a permutation w such that

$$w(i) = j \iff \dim \frac{F^{i-1} \cap V_j + F^i}{F^{i-1} \cap V_{j-1} + F^i} = 1.$$

We can similarly define

$$\Sigma_w^\circ = \{V_\bullet \in \mathrm{Fl}(k, n) \mid \text{permutations of } V = w\}.$$

$$\Sigma_w = \text{closure of } \Sigma_w^\circ, \quad \sigma_w = [\Sigma_w] \in H^\bullet(\mathrm{Fl}(n)).$$

Littlewood–Richardson coefficients

It is known that

$$H^\bullet(\mathrm{Fl}(n)) = \bigoplus_{w \in S_n} \mathbb{Q} \cdot \sigma_w \quad (\text{as a vector space})$$

The central problem in Schubert calculus is to compute the coefficients c_{uv}^w in the expression

$$\sigma_u \cdot \sigma_v = \sum_{w \in S_n} c_{uv}^w \cdot \sigma_w.$$

There is no general combinatorial model for c_{uv}^w up to now.

Schubert polynomials

To study it, we define *Schubert polynomials*. For $w \in S_\infty$


$$\begin{aligned}\mathfrak{S}_{n \cdots 21} &= x_1^{n-1} x_2^{n-2} \cdots x_{n-1}, \\ \mathfrak{S}_{w(i,i+1)} &= \frac{\mathfrak{S}_w - \mathfrak{S}_{w|_{x_i \leftrightarrow x_{i+1}}}}{x_i - x_{i+1}}, \quad w_i < w_{i+1}.\end{aligned}$$

It turns out the structure constant can be computed by

$$\mathfrak{S}_u \cdot \mathfrak{S}_v = \sum_{w \in S_\infty} c_{uv}^w \cdot \mathfrak{S}_w.$$

Thus we translate a geometric problem to an algebraic problem.

Examples



We have


$$\sigma_{213} \cdot \sigma_{132} = \sigma_{231} + \sigma_{312}.$$

Bumpless pipe dream

There is an amazing combinatorial model for Schubert polynomials called *bumpless pipe dream*.

Theorem (Lam, Lee, and Shimozono)

Schubert polynomial \mathfrak{S}_w is the weighted sum of



using



such that each pair of pipes crosses at most once

Examples

$$\mathfrak{S}_{321} = x_1^2 x_2$$

$$\mathfrak{S}_{231} = x_1 x_2$$

$$\mathfrak{S}_{312} = x_1^2$$

$$\mathfrak{S}_{213} = x_1$$

$$\mathfrak{S}_{132} = x_1 + x_2$$

$$\mathfrak{S}_{123} = 1$$

Generalization A

$$H^\bullet(\mathrm{Fl}(n)) \leadsto K(\mathrm{Fl}(n))$$

Similarly,

$$K(\mathrm{Fl}(n)) = \bigoplus_{w \in S_n} \mathbb{Q} \cdot \mathcal{O}_w \quad (\text{as a vector space}).$$


The structure constant of \mathcal{O}_w is the same as the the structure constant of *Grothendieck polynomials*:

$$\mathfrak{G}_{n \dots 21} = x_1^{n-1} x_2^{n-2} \cdots x_{n-1},$$

$$\mathfrak{G}_{w(i,i+1)} = \frac{(1 + \beta x_{i+1}) \mathfrak{G}_w - (1 + \beta x_i) \mathfrak{G}_{w|_{x_i \leftrightarrow x_{i+1}}}}{x_i - x_{i+1}}, \quad w_i < w_{i+1}.$$

Theorem (Weigandt)


Grothendieck polynomial \mathfrak{G}_w is the weighted sum of



using

						
1				β	$1 + \beta x_i$	x_i

such that

each pair of pipes crosses at most once
in each , the J-pipe $>$ the Γ -pipe

Generalization B

$$H^\bullet(\mathrm{Fl}(k, n)) \leadsto H_T^\bullet(\mathrm{Fl}(n))$$

We have

$$H_T^\bullet(\mathrm{Fl}(n)) = \bigoplus_{w \in S_n} \mathbb{Q}[t_1, \dots, t_n] \cdot \sigma_w$$

$$K_T(\mathrm{Fl}(n)) = \bigoplus_{w \in S_n} \mathbb{Q}[\tau_1^{\pm 1}, \dots, t_n^{\pm 1}] \cdot \mathcal{O}_w$$

The corresponding polynomial is known as *double Schubert/Grothendieck polynomial*.

Theorem (Lam, Lee, and Shimozono)

Double Schubert polynomial \mathfrak{S}_w is the weighted sum of bumpless pipe dreams but with double weight:

	1	
		$x_i - t_j$

Theorem (Weigandt)

Double Grothendieck polynomial \mathfrak{G}_w is the weighted sum of bumpless pipe dreams but with double weight:

	1		β		$1 + \beta(x_i - t_j)$		$x_i - t_j$
---	-----	---	---------	---	------------------------	--	-------------

Examples




$$(x_1 - t_1)(x_1 - t_2)(x_2 - t_1)$$



$$(x_1 - t_1)$$



$$(x_1 - t_1)(x_2 - t_1)$$



$$(x_2 - t_2) + (x_1 - t_1)$$



$$(x_1 - t_1)(x_1 - t_2)$$



$$1$$

Seperated descents

Assume $u, v \in S_n$ have seperated descents

$$\max(\text{des}(u)) \leq k \leq \min(\text{des}(v)).$$

There is a very recent combinatorial rule by Knutson and Zinn-Justin for the expansion of

$$\mathcal{O}_u \cdot \mathcal{O}_v = \sum_w c_{uv}^w(t) \cdot \mathcal{O}_w,$$

We generalize it to the *triple version*.

Our main result

single Schubert calculus	double Schubert calculus	triple Schubert calculus
non-equivariant $\mathfrak{G}_u(x)\mathfrak{G}_v(x)$	equivariant $\mathfrak{G}_u(x, t)\mathfrak{G}_v(x, t)$	★ $\mathfrak{G}_u(x, t)\mathfrak{G}_v(x, y)$

We can view triple Schubert calculus as the *universal rule* for

$$\mathfrak{G}_u(x, t) \cdot \mathfrak{G}_v(x, wt)$$

which geometrically corresponds to the intersection of Schubert varieties of different transversality.


Theorem (FGX)

There is a combinatorial rule for $c_{uv}^w(y, t)$ in the expansion

$$\mathfrak{G}_u(x, y) \cdot \mathfrak{G}_v(x, t) = \sum_{w \in S_\infty} c_{uv}^w(y, t) \cdot \mathfrak{G}_w(x, t).$$


Pipe Puzzles

Let us first state the rule for cohomology, i.e. $\beta = 0$.



using



in each , the hori. pipe $<$ the vert. pipe

For K-theory, it can be computed by using one more piece .

Example



Recall

$$\mathfrak{S}_{213}(x, y) = x_1 - y_1$$

$$\mathfrak{S}_{132}(x, t) = x_1 + x_2 - t_1 - t_2$$

$$\mathfrak{S}_{231}(x, t) = (x_1 - t_1)(x_2 - t_1) \quad \mathfrak{S}_{312}(x, t) = (x_1 - t_1)(x_1 - t_2)$$

$$k = 1, \quad u = 2 \mid 13, \quad v = 1 \mid 32.$$



$$\mathfrak{S}_{213}(x, y) \cdot \mathfrak{S}_{132}(x, t) = (t_1 - y_1) \mathfrak{S}_{132}(x, t) + \mathfrak{S}_{231}(x, t) + \mathfrak{S}_{312}(x, t).$$

On the proof

Our proof is based on the classical *6-vertex model*, and is significantly simple! What we need is to prove

I. induction on y II. induction on t III. initial cases.




Historically, people realized that equivariant cohomology is usually easier than usual cohomology.


$$\boxed{\text{single}} \implies \boxed{\text{double}}$$

It turns out the same happens for

$$\boxed{\text{double}} \implies \boxed{\text{triple}}$$

Specialization A — separated descents puzzles

If we set $y_i = t_i$, then on the diagonal  has weight 0. So it suffices to count those with  or  on the diagonal; so all pipes must go straight down under the diagonal. So we only need the upper triangle. This specializes to Knutson and Zinn-Justin's puzzle.



$$\mathfrak{S}_{213}(x, t) \cdot \mathfrak{S}_{132}(x, t) = \mathfrak{S}_{231}(x, t) + \mathfrak{S}_{312}(x, t).$$

Specialization B — bumpless pipe dream

If we set $k = n$, then $v = \text{id}$. Taking $x = t$ on both sides of

$$\mathfrak{G}_u(x, \textcolor{red}{y}) \cdot \mathfrak{G}_v(x, t) = \sum_{w \in S_\infty} c_{uv}^w(\textcolor{red}{y}, t) \cdot \mathfrak{G}_w(x, t),$$

we will get


$$\mathfrak{G}_u(t, \textcolor{red}{y}) = c_{u \text{id}}^{\text{id}}(\textcolor{red}{y}, t).$$

By reflecting against the diagonal and changing the labels, we recover the Weigandt's model of bumpless pipe dream.




Specialization C — classical puzzles

When u and v are both k -Grassmannian (i.e. at most one descent at k), we can recover the Grassmannian puzzles introduced in the first part. First, let us color pipes $\leq k$ by **red** and $\geq k$ by **blue**. Then we replace



Then rotate 120° anticlockwise.



Thanks

