The 17 taxa reported decreased in stool in all three conditions: 9 genera and 8 higher-rank nodes

7 of them had more decreased than increased stool studies in every condition. Keeping only genus and species ranks gives mean d_J = 0.478, close to the primary 0.479.

A. Mean pairwise d_J under each set definition



B. Stool studies reporting each core taxon ↓ or ↑

		PD	HIV	CRC	net ↓ in all 3
Roseburia	genus	20↓ 4↑	2↓ 0↑	6↓ 0↑	✓
Blautia	genus	13↓ 2↑	1↓ 3↑	6↓ 1↑	
Clostridium	genus	7↓ 3↑	5↓ 2↑	6↓ 3↑	✓
Bacteroides	genus	6↓ 3↑	8↓ 0↑	2↓ 2↑	
Lachnospira	genus	5↓ 2↑	3↓ 0↑	6↓ 1↑	✓
Ruminococcus	genus	6↓ 2↑	4↓ 0↑	4↓ 3↑	✓
Anaerostipes	genus	2↓ 1↑	3↓ 0↑	3↓ 1↑	✓
Eubacterium	genus	1↓ 1↑	1↓ 1↑	3↓ 2↑	
Butyrivibrio	genus	2↓ 2↑	1↓ 2↑	1↓ 0↑	
Lachnospiraceae	family	20↓ 7↑	5↓ 3↑	9↓ 5↑	✓
Bacillota	phylum	8↓ 5↑	4↓ 2↑	4↓ 1↑	✓
Clostridia	class	5↓ 7↑	4↓ 0↑	7↓ 1↑	
Oscillospiraceae	family	8↓ 13↑	4↓ 0↑	2↓ 1↑	
Clostridiaceae	family	8↓ 2↑	3↓ 0↑	2↓ 2↑	
Eubacteriales	order	5↓ 6↑	4↓ 1↑	4↓ 1↑	
Bacteroidia	class	5↓ 2↑	2↓ 0↑	1↓ 4↑	
Bacteroidota	phylum	3↓ 0↑	3↓ 1↑	1↓ 4↑	

Dot: blue when ↓ studies outnumber ↑, red when ↑ outnumber ↓, grey when equal. Upper block: genera; lower block: family, order, class and phylum nodes, whose annotations are rolled up from descendant taxa. Net ↓ = more decreased than increased studies in every condition.

A taxon is in a condition's set when at least one decreased stool signature reports it; the counts in B show that some core taxa are also reported increased, sometimes more often (for example Oscillospiraceae in PD and Bacteroidota in CRC). PD, Parkinson's disease; HIV, HIV infection; CRC, colorectal cancer. BugSigDB 2026-04-24_00:41_UTC; human profiles; condition and body site from each experiment's own record.